

### Tail color signals performance in blue tit nestlings

Journal:	<i>Journal of Evolutionary Biology</i>
Manuscript ID	JEB-2018-00555.R3
Manuscript Type:	Research Papers
Keywords:	Sexual selection, Coloration, Good genes, G matrix, Heritability, <i>Cyanistes caeruleus</i> , Wild, Immune response

SCHOLARONE™  
Manuscripts

## 1 Tail color signals performance in blue tit nestlings

2

3 **Abstract**

4 Indirect sexual selection arises when reproductive individuals choose their mates based on  
5 heritable ornaments that are genetically correlated to fitness. Evidence for genetic associations  
6 between ornamental coloration and fitness remain scarce. In this study we investigate the  
7 quantitative genetic relationship between different aspects of tail structural coloration  
8 (brightness, hue and UV chroma) and performance (cell mediated immunity, body mass and  
9 wing length) in blue tit (*Cyanistes caeruleus*) nestlings. In line with previous studies, we find  
10 low heritability for structural coloration and moderate heritability for performance measures.  
11 Multivariate animal models show positive genetic correlations between the three measures of  
12 performance, indicating quantitative genetic variation for overall performance while tail  
13 brightness and UV chroma, two genetically independent color measures, are genetically  
14 correlated with performance (positively and negatively respectively). Our results suggest that  
15 mate choice based on independent aspects of tail coloration can have fitness payoffs in blue tits  
16 and provide support for the indirect benefits hypothesis. However, low heritability of tail  
17 structural coloration implies that indirect sexual selection on mate choice for this ornament will  
18 be a weak evolutionary force.

19

20 **Keywords:** Sexual selection, coloration, good genes, genetic correlation, G matrix,  
21 heritability, *Cyanistes caeruleus*, wild, immune response

## 22 **Introduction**

23 Sexual selection theory has provided a powerful framework to understand the existence and  
24 function of ornaments. Reproductive individuals are predicted to choose their mates based on  
25 potential direct benefits (“good parent”) or indirect benefits (“good genes”) that they provide  
26 (Kirkpatrick & Barton 1997). In particular, the good genes model implies that genes increasing  
27 fitness can be phenotypically signaled by individuals, for example through ornamentation.  
28 While direct benefits models of mate choice have been well supported empirically, the indirect  
29 benefits models requires that ornaments and fitness are heritable and positively genetically  
30 correlated (Kirkpatrick & Barton 1997), which thus far is lacking evidence (Jones & Ratterman  
31 2009).

32 Birds often exhibit colorful plumage patches, which have been extensively studied in relation  
33 to mate choice and intra-sexual competition (Amundsen 2000). In particular, box-breeding  
34 birds are ideal study systems to test for indirect benefits of ornaments because they allow long-  
35 term monitoring and cross-fostering experiment. Indeed, the combination of pedigree data  
36 obtained through long-term monitoring, and cross-fostering can be effectively used to  
37 decompose environmental from genetic resemblance between relatives (Kruuk & Hadfield  
38 2007). Cross-fostering experiments are used to separately estimate the effects of foster and  
39 genetic parents’ traits on offspring performance (e.g. growth, mass, immunity) or viability, the  
40 former capturing direct effects (through parental care) and the latter capturing indirect (genetic)  
41 effects (assuming the absence of early environmental and maternal effects). In great tits (*Parus*  
42 *major*) for example, such experiments have provided support for direct and indirect benefits of  
43 carotenoid and melanin-based coloration in males and females. Indeed, great tits with a larger  
44 black breast stripe were found to produce more viable or heavier young (Norris 1993, Remeš  
45 & Matysiokova 2013) while yellower males and females with a blacker breast stripe were found  
46 to raise heavier nestlings (Pickett et al. 2013).

47 In combination with cross-fostering, quantitative genetics, which use information on the  
48 relatedness between individuals derived from the population pedigree, are particularly relevant  
49 to study sexual selection (Reid 2014). Indeed, this approach allows disentangling early  
50 environmental or maternal effects from additive genetic effects causing resemblance between  
51 relatives (Kruuk & Hadfield 2007), which cannot be done using only cross-fostering. In  
52 practice, a (co)variance partitioning approach implemented using (multivariate) mixed models,  
53 can be used to estimate the heritability of ornaments, fitness and their genetic correlation along  
54 with other sources of variance in and covariance between them (e.g. year, mother, nest of  
55 rearing). Earlier quantitative genetic studies find low heritability for structural and carotenoid-  
56 based ornaments (Hadfield et al. 2006, Evans & Sheldon 2012, Drobniak et al. 2013,  
57 Charmantier et al. 2017), as well as recruitment (Hadfield et al. 2006) in blue tits (*Cyanistes*  
58 *caeruleus*). In contrast, performance traits which are correlates of fitness, such as body  
59 condition or growth in nestlings, are found to be moderately heritable in collared flycatchers  
60 (*Ficedula albicollis*, Merilä et al. 2001, Pitala et al. 2007) and blue tits (Hadfield et al. 2007).  
61 To our knowledge, only two studies estimated the genetic correlation between a structural color  
62 ornament and fitness, and did not find support for the indirect benefits hypothesis (Hadfield et  
63 al. 2006, Qvarnström et al. 2006).

64 Under a particular case of the indirect benefits hypothesis, the “parasite-mediated sexual  
65 selection” (PMSS) hypothesis (cf. Hamilton & Zuk 1982), individuals’ sexual ornaments are  
66 predicted to signal immunocompetence. This hypothesis is based on the idea that cycles of  
67 coadaptation between hosts and parasites, which maintain additive genetic variation in  
68 resistance and thus fitness in hosts, can promote sexual selection for displays that signal  
69 resistance. Within species exposed to various chronic parasites, showiness is hence predicted  
70 to signal resistance to a wide variety of parasites. One common proxy of immunocompetence  
71 is the T-cell mediated immune response, assayed using phytohaemagglutinin (PHA). Although

72 PHA was shown to be condition-dependent (Alonso-Alvarez & Tella 2001, Thompson et al.  
73 2014), several studies showed that it is heritable in various bird populations and species (Pitala  
74 et al. 2007, Bonneaud et al. 2009, Kinnard & Westneat 2009, Kim et al. 2013, Sakaluk et al.  
75 2014). To date, the few cross-fostering studies testing the association between males' or  
76 females' ornaments and nestlings' PHA find mixed evidence for the presence of such  
77 relationship. In the blue-footed booby (*Sula nebouxii*), Velando et al. (2005) found no  
78 correlation between nestling PHA and father's foot coloration while in great tits, Pickett et al.  
79 (2013) found a positive correlation between nestlings' PHA and genetic father's but not  
80 mother's yellow plumage brightness. In the same species, Remeš & Matysiokova (2013) found  
81 that females with more immaculate white cheeks produce offspring with a higher PHA. In  
82 addition, only two studies estimated genetic correlations between coloration and PHA in adult  
83 male zebra finches (*Taeniopygia guttata*, Birkhead et al. 2006) and in common kestrel nestlings  
84 (*Falco tinnunculus*, Kim et al. 2013) and did not support the PMSS hypothesis.

85 Therefore, empirical evidence for the indirect fitness benefits and PMSS hypotheses remains  
86 equivocal, and more quantitative genetic studies are needed to gain insights into the evolution  
87 of color ornaments through sexual selection. In the present study, we investigate the quantitative  
88 genetic association between blue ornamentation and performance in blue tit nestlings. In this  
89 species, structural colors, in particular the blue cap of adults, have been extensively studied.  
90 This ornament has been shown to be sexually dimorphic (Hunt et al. 1998) and involved in  
91 mate choice (Andersson et al. 1998, Hunt et al. 1999) and intra-sexual competition (Alonso-  
92 Alvarez et al. 2004, Remy et al. 2010). In addition to their cap, blue tits show sexually  
93 dimorphic blue coloration of their tail feathers, which is already detectable at the nestling stage  
94 (Johnsen et al. 2003). Because most birds do not molt their tail feathers during their first year  
95 (Peters et al. 2007, Svensson 1992), the structural coloration of tail feathers may be involved in  
96 sexual selection. Here, we estimate genetic correlations between different aspects of tail

97 structural color (brightness, hue, UV chroma) and different performance measures (PHA, body  
98 mass, and wing length) using a multivariate animal model. This study provides rare estimates  
99 of genetic correlations between color and performance and supports the indirect benefits  
100 hypothesis.

101

## 102 **Material & Methods**

### 103 *Measures of blue tit nestlings*

104 Blue tits were studied in a nest-box breeding population in south-west Finland (Tammisaari,  
105 60°01' N, 23°31' E). Hatch date of a brood (day 0) was established by daily checks. When the  
106 offspring were 2 days old (2005-2009), approximately half of each brood (on average 42±9%)  
107 was reciprocally swapped between a pair of nests with same aged and similar sized offspring  
108 (average brood mass at day 0). Nestlings were weighed and individually marked by clipping  
109 their nails. Whether the heaviest nestling was cross-fostered or not was decided at random and  
110 this action was subsequently alternated down the mass ranking of offspring (for more details  
111 see Brommer & Klun 2012). When nestlings were 13 days old, the thickness of their wing web  
112 after feather removal was measured (to the nearest 0.01mm) two times using a spessimetre  
113 (Mitutoya 700-117SU, modified by the removal of a spring). Nestlings were injected with 0.04  
114 ml of a solution of 5mg ml<sup>-1</sup> Phytohaemagglutinin (Sigma code L-8754) in saline. After 24 hours,  
115 the thickness of the wing web was measured (to the nearest 0.01mm) three times and PHA,  
116 which is the responsive swelling, was calculated as the difference between the average thickness  
117 before and after injection. This procedure was carried out from 2003 to 2007 (except in 2004).  
118 A higher PHA reflects a stronger T-cell mediated response and is a measure of innate immunity  
119 (Smits et al. 1999). When nestlings were 16 days old, their tarsus length was measured with a  
120 sliding caliper (0.1mm accuracy) and they were weighed using a spring balance (accuracy 0.1g).

121 Their wing length was measured using a ruler (1mm accuracy) and then one middle tail feather  
122 was pulled. Nestlings were sexed using molecular markers (see Brommer & Kluen 2012).

### 123 *Spectrometry*

124 Reflectance was measured in the lab using a spectrometer (Avantes AvaSpec-2048-SPU2) and  
125 a deuterium-halogen light source (AvaLight-DH-S). The light source and the probe were  
126 maintained at a 90° angle. Reflectance of the blue in the tail feather of offspring was measured  
127 just where the feather vane comes out of the blood shaft of the feather. Because this spot is  
128 rather small in nestlings and can be missed by the incident light beam, we measured its  
129 reflectance five times. Each spectra was smoothed using a  $\pm 10$ nm running average and we  
130 discarded measurements where reflectance did not decrease between 320 and 600nm, which is  
131 the general pattern observed for tail structural coloration (see Johnsen et al. 2003 and Figure  
132 S1). Feathers collected in 2005 and 2006 were measured in 2006-2007 (University of Jyväskylä,  
133 spectrophotometer 1) while the rest of the feathers (2003, 2007-2009) was measured in 2009-  
134 2010 (University of Turku) on a different spectrometer of the same model (spectrophotometer  
135 2). Spectra obtained from these two spectrometers have different averages (Figure S1), which  
136 was accounted for statistically in further analyses. We used reflectance values measured  
137 between 320 and 600 nm in the calculations of the following metrics of coloration: (1)  
138 Brightness was quantified as the total reflectance of the feather, (2) hue was calculated as the  
139 wavelength of maximum reflectance and (3) UV chroma as the proportion of the total  
140 reflectance comprised between 320 and 400nm. Sample sizes (number of individuals and  
141 broods), means and standard deviations for each trait are provided in Table 1 and their  
142 distribution is plotted in Figure S2. The length of the vane of the feather was measured using a  
143 sliding caliper and was used as a covariate in all analyses.

### 144 *Quantitative genetic analyses*

145 The pruned pedigree (including all phenotyped individuals and up to 3 generations of  
 146 unphenotyped links between them) was obtained using the function trimPed from the R package  
 147 “pedigree” (Coster 2012) and was analyzed using the R package “pedantics” (Morrissey 2018).  
 148 It holds records for 3652 individuals, of which 671 are founders. Mean maternal and paternal  
 149 sibship sizes are 10.1 and 10.5 respectively, with 2950 dams and 2709 sires. This pedigree was  
 150 collected over multiple generations, with a maximum pedigree depth of 5 generations, and 2592  
 151 grandparents. All nestlings hatched in a nest were assumed to be sired by their social father.  
 152 Extra-pair paternity in this population is not known but is probably within the range of what  
 153 was found in other populations (7-25%, Brommer et al. 2010). This level of extra-pair paternity  
 154 is likely to cause little error in the estimation of quantitative genetic parameters (Charmantier  
 155 & Réale 2005). There was phenotypic data for 430 genetic broods and on average 69% of them  
 156 (65%-89%) were reciprocally cross-fostered each year during 2005–2009. Broods from 2003  
 157 were not cross-fostered but represent less than 5% of all broods, which is unlikely to affect our  
 158 estimates of quantitative genetic (co) variances.

159 We first estimated additive genetic variance for tail brightness, hue, UV chroma, PHA, body  
 160 mass, and wing length separately using univariate animal models. An animal model is a form  
 161 of mixed model which allows partitioning phenotypic variance into variance due to additive  
 162 genetic effects and other sources of variation, using information on the relatedness between  
 163 individuals derived from the population pedigree (Wilson et al. 2009). This model is noted:

$$164 \quad y = X\beta + Z_A\mathbf{u}_A + Z_{CE}\mathbf{u}_{CE} + \boldsymbol{\varepsilon} \quad (1)$$

165 Where  $\mathbf{y}$  is a vector containing all observations on all individuals for each trait,  $\beta$  is a vector of  
 166 fixed effects and  $X$  the design matrix relating fixed effects to each individual observation. The  
 167 vector  $\mathbf{u}_A$ , fitted as a random effect, is the vector of additive genetic effects, and its covariance  
 168 structure is assumed to be proportional to the relatedness matrix  $Z_A$ . To account for common

169 environmental effects occurring when individuals share the same nest,  $\mathbf{u}_{CE}$  (and its design  
 170 matrix  $Z_{CE}$ ) was fitted as an additional random effect. Finally,  $\boldsymbol{\varepsilon}$  is a vector of residual errors  
 171 capturing differences between individuals that are unexplained by fixed, additive genetic, and  
 172 common environment effects. Here, the phenotypic variance ( $V_P$ ) in performance traits is  
 173 decomposed into additive genetic variance ( $V_A$ ), common environment variance ( $V_{CE}$ ) and  
 174 residual variance ( $V_R$ ) as:

$$175 \quad V_P = V_A + V_{CE} + V_R \quad (2)$$

176 Because each individual's feather was measured multiple times, an additional individual  
 177 random effect  $\mathbf{Z}_I \mathbf{u}_I$  was fitted to capture among-individual differences in color measures. In  
 178 these models, the residual component thus captures variation between measurements of the  
 179 same feather (measurement error). Therefore, the phenotypic variance in color measures is here  
 180 decomposed into:

$$181 \quad V_P = V_A + V_{CE} + V_I + V_{ME} \quad (3)$$

182 Where  $V_I$  is the variance between individuals (equivalent to  $V_R$  in 2) and  $V_{ME}$  the variance in  
 183 measurement error.

184 Residuals of all animal models were approximately normally distributed (Shapiro-Wilk test  
 185 values  $>0.92$ ; Figure S3). Because of visible differences between reflectance measurements  
 186 made by the two spectrometers (Figure S1 and Table S1),  $V_{CE}$ ,  $V_I$  and  $V_{ME}$  of color measures  
 187 were allowed to vary between these two spectrometers. This was done to assure that our  
 188 inference of the average additive genetic  $\mathbf{G}$  matrix for all nestling traits during the entire study  
 189 period was accurate.

190 In all models, sex and year were fitted as categorical fixed effects to account for sexual  
 191 dimorphism and between-year average differences. Sexes were pooled because cross-sex

192 genetic correlations for all six traits are high and sex-specific  $\mathbf{G}$  matrices are qualitatively  
193 similar (see Tables S2-S10 and Figure S4). The length of the feather vane was fitted as fixed  
194 effect covariate for all color measures, and tarsus length was included as a covariate in analyses  
195 of nestling's body mass to correct for body size. The animal model was solved using Restricted  
196 Maximum Likelihood (REML) and implemented in R (R Development Core Team 2018) using  
197 the package "asreml" (Butler et al. 2009). The statistical significance of  $V_A$  was tested by  
198 comparing each model with a model where  $V_A$  was not estimated, using likelihood ratio tests  
199 (LRT) with one degree of freedom. Heritability was calculated as the ratio  $V_A/V_P$  where  $V_P$  did  
200 not include  $V_{ME}$  for color measures. Uncertainty of this ratio was calculated using the delta  
201 method (Fischer et al. 2004).

202 To partition the covariances between traits into different components, we used a multivariate  
203 animal model, in which performance traits and individual averages of each color measure were  
204 all fitted as response variables. Using individual averages for the color traits only reduces  $V_{ME}$   
205 and does not alter the estimation of  $V_A$ . Random effects fitted in this model therefore only  
206 included additive genetic effects, common environment effects and residual errors. Fixed  
207 effects were similar to the ones used in univariate animal models, some being trait-specific.  
208 Covariances on the additive genetic level were tested individually by comparing the  
209 unconstrained models to models where these covariances were fixed to zero using LRT, with 1  
210 degree of freedom. The estimated  $\mathbf{G}$  matrix provides 15 estimates of genetic correlations  
211 between all six traits. R code for quantitative genetic analyses is provided in Text S1.

### 212 *Structural equation models*

213 We performed structural equation models (SEM) to reduce the dimensionality of the  $\mathbf{G}$  matrix  
214 and investigate the general relationship between each color trait and performance. To do so,  $\mathbf{G}$   
215 matrix estimated by the six-trait animal model was first transformed into a correlation matrix

216 and used as input data into different SEMs. All performance traits were reduced to a latent  
217 factor named “performance” on which they loaded positively. In these models, the variances of  
218 “performance” and of each color trait were fixed to 1. Because a correlation matrix was used  
219 as input data, the residual variance of each indicator (here, each performance trait), which is the  
220 variance unexplained by the latent factor, was fixed to 1 minus its squared factor loading. Each  
221 SEM was fitted in R using the package “lavaan” (Rosseel 2012). Sample size in these models  
222 was set at 306 as this was the number of nestlings with at least 1 trait measured ( $n=3240$ ) divided  
223 by the average paternal sibship size (10.6) and thus approximates the number of families. The  
224 sample size in a SEM will not affect the inferred loadings or correlations between latent  
225 variables but impacts their uncertainty. In order to take forward the uncertainty of the **G** matrix  
226 estimates into the SEM estimates, we calculated 95% confidence intervals (CI) of these  
227 correlations using simulation. We first simulated phenotypic data based on the population  
228 pedigree and the estimated **G** matrix using the R package “pedantics”. Each simulated data set  
229 was analyzed using a multivariate animal model and the estimated **G** matrix was used as input  
230 data in a SEM. Only SEM models in which all loadings of performance traits were  $> 0.05$  were  
231 kept and simulations were run until obtaining 1000 estimate of correlations between each color  
232 trait and performance. This procedure excluded models in which either one loading of a  
233 performance trait on “performance” or the correlation between color and performance were  
234 abnormally high ( $>10$ ) while the (other) loadings were zero. This situation occurred under  
235 certain values of the **G** matrix (e.g. when low correlations switch signs) and thus, SEM  
236 estimates based on these values were discarded. As a result, the 95% CI of the average  
237 correlation between color traits and performance may not have fully incorporated the  
238 uncertainty of the **G** matrix and should not be interpreted as a test for statistical significance. R  
239 code for performing SEMs and simulations are provided in Text S2 and Text S3.

## 240 **Results**

241 All tail color and performance traits were measured in over 2800 nestlings, except the PHA  
242 response, which was measured on fewer nestlings (Table 1). Measurement repeatability of tail  
243 color was significant, although relatively low (40-50% for brightness and UV chroma; around  
244 30% for hue, Table S1) compared to measurement repeatability of adult color in other blue tit  
245 populations (e.g. Doutrelant et al. 2008, Figuerola et al. 1999). There was significant sexual  
246 dimorphism in these nestlings for all tail color measures but also for mass and wing length  
247 (Table S11-S16).

248 Heritability of tail color measures was low (1-12%, Table 2). These color measures were  
249 negatively but not significantly correlated with each other on the genetic level (trivariate model:  
250 UV chroma-brightness:  $r_G=-0.17$ ,  $se=0.36$ ),  $\chi_1^2=0.07$ ,  $p=0.78$ ; hue-brightness:  $r_G=-0.75$ ,  
251  $se=0.45$ ,  $\chi_1^2=2.50$ ,  $p=0.11$ ; hue-UV chroma:  $r_G=-0.34$ ,  $se=0.54$ ,  $\chi_1^2=0.37$ ,  $p=0.54$ ; Table 3). We  
252 hence studied color traits independently in further analyses. In contrast, heritability estimates  
253 of performance traits (body mass, wing length, and PHA response) were all moderate (19-28%,  
254 Table 2). Heritability of PHA using repeated measures of nestlings' wing web before and after  
255 injection was also calculated (excluding measurement error) using a bivariate animal model and  
256 was close to the estimate based on individual averages ( $h^2=0.16$ ,  $se=0.04$ ; Table S17).

257 All performance traits were positively correlated on the genetic level (trivariate model: PHA-  
258 body mass:  $r_G=0.61$ ,  $se=0.04$ ,  $\chi_1^2=53.1$ ,  $p<0.001$ ; wing-PHA:  $r_G=0.36$ ,  $se=0.07$ ,  $\chi_1^2=30.5$ ,  
259  $p<0.001$ ; body mass-wing:  $r_G=0.46$ ,  $se=0.07$ ,  $\chi_1^2=408.0$ ,  $p<0.001$ ; Table 3). These correlations  
260 were not affected by the nestlings' wing web thickness before injection (PHA- body mass:  
261  $r_G=0.59$ ,  $se=0.04$ ; wing-PHA:  $r_G=0.36$ ,  $se=0.07$ ; body mass-wing:  $r_G=0.43$ ,  $se=0.06$ ).  
262 (Co)Variance matrices are all reported in the electronic supplementary material (Tables S18-  
263 S22).

264 Regarding genetic correlations between coloration and performance traits, inspection of the **G**  
265 matrix (highlighted part in Table 3) shows that UV chroma is consistently negatively correlated  
266 with performance traits. In contrast, the signs of the genetic correlations between the two other  
267 color measures and performance traits are inconsistent. Separate SEMs were run for each color  
268 trait independently, to estimate its average correlation with performance. These analyses  
269 indicated that tail brightness was (largely) positively (coefficient = 0.39; 95% CI= -0.09–1.00),  
270 and UV chroma negatively (coefficient = -0.69; 95% CI=-1.36–-0.15) correlated with  
271 performance. For hue, this correlation was weakly negative (coefficient = -0.12; 95% CI=-  
272 0.81–0.40, Figure S5).

## 273 **Discussion**

274 In this study, we used long-term nestling data, pedigree information and a cross-fostering design  
275 to estimate additive genetic variation in coloration measures, performance traits and genetic  
276 correlations between them, and find evidence for indirect benefits of blue tail coloration in wild  
277 blue tits.

278 Tail color measures had a low heritability which is in line with other heritability estimates of  
279 carotenoid and structural coloration in blue tits (Evans & Sheldon 2012, Hadfield et al. 2006,  
280 Drobniak et al. 2013, Charmantier et al. 2017). In contrast, heritability of performance traits  
281 (body mass, wing length, and PHA response) was generally moderate and within the range of  
282 what was found in other populations or species (see e.g. Merilä & Sheldon 2000, Merilä et al.  
283 2001, Hadfield et al. 2007 for morphometric traits, and Pitala et al. 2007, Kim et al. 2013 for  
284 PHA response). In addition all performance traits were positively correlated on the genetic  
285 level, which is consistent with previous estimates of genetic correlations between mass and  
286 wing length (e.g. Björklund et al. 2013) and between mass and PHA response (Kim et al. 2013)  
287 and indicates the presence of additive genetic variation for individual performance. Using

288 SEMs, we found that the average genetic correlation between each color measure and a latent  
289 factor “performance”, was positive for tail brightness, negative for UV chroma, and weakly  
290 negative for hue.

291 Hence, our findings suggest indirect fitness benefits of choosing mates with tails that are  
292 brighter or reflecting less in the UV. Indeed, all three performance traits are known to be  
293 positively associated with fitness. While the positive effects of mass at fledging on survival  
294 have been well documented (e.g. Perrins 1965, Tinbergen & Boerlijst 1990, Lindén et al. 1992,  
295 Radersma et al. 2015), wing length was shown to determine fledging date (Radersma et al.  
296 2011), and impact recruitment probability (Verboven & Visser 1998), possibly through  
297 competitive advantage or decreased predation risk. PHA can also be considered an important  
298 functional trait as it was shown to increase survival and recruitment probability of nestlings in  
299 various species (Hörak et al. 1999, Moreno et al. 2005, Cichoń & Dubiec 2005, López-Rull et  
300 al. 2011), although this relationship may be due to its condition-dependence (Alonso-Alvarez  
301 & Tella 2001, Thompson et al. 2014). Nevertheless, PHA, body mass and wing length were  
302 heritable and genetically correlated, which suggests that their genetic architecture “captures the  
303 so-called good genes” and allow sexual selection for indirect benefits to occur in this  
304 population. Because tail color indicates immune response through its genetic relationship with  
305 performance, our results also support the PMSS hypothesis.

306 Previous studies showed that bluer plumage ornaments are condition-dependent in juveniles  
307 (Johnsen et al. 2003, Jacot & Kempenaers 2007, Peters et al. 2007 in blue tits, Siefferman et al.  
308 2008, Siefferman & Hill 2007 in other species) and in adults (Doutrelant et al. 2012, Beck et  
309 al. 2015, Galván 2011), which contrasts with the negative genetic association that we found  
310 between hue, UV chroma, and performance traits. Importantly, correlations between hue, UV  
311 chroma and performance traits were also negative on the phenotypic level (Table S23). This  
312 negative relationship implies that developing bluer tails early in life has costs which can impact

313 individuals' probability to recruit. However, these fitness costs may be offset on the first  
314 reproductive year, if yearlings with a bluer tail are sexually more attractive than duller yearlings.  
315 On the other hand, genetic correlations between hue, UV chroma and performance should be  
316 interpreted with caution given the very low heritability of both color measures and the high  
317 uncertainty of their additive genetic variance estimates.

318 In the context of sexual selection based on indirect benefits, key parameters are the (square  
319 root) heritability of the sexually selected trait, additive genetic variance in fitness and genetic  
320 correlation between this trait and fitness (Kirkpatrick and Barton 1997). This means that, for  
321 mate choice based on indirect benefits of an ornament to evolve, this ornament has to provide  
322 sufficient information regarding an individual's genotype. Because we found low heritability  
323 for tail color measures, our results imply that indirect sexual selection on mate choice for this  
324 ornament will be a weak evolutionary force (cf. Møller & Alatalo 1999, Qvarnström et al.  
325 2006).

326 Despite broad interest in understanding the evolution of animal ornaments, studies estimating  
327 heritability of coloration and its genetic correlation with performance to test the indirect benefits  
328 hypothesis remain rare. This study provides evidence for additive genetic variation in tail  
329 coloration and performance and for a genetic association between them. More specifically, our  
330 findings suggest that mate choice based on tail brightness and UV chroma can have fitness  
331 payoffs in blue tits and hence support indirect benefits of mate choice for this ornament.  
332 However, because of the low heritability of tail structural coloration, this mechanism may not  
333 be a major driver of its evolution.

334 **References**

- 335 Alonso-Alvarez, C., & Tella, J. L. (2001). Effects of experimental food restriction and body-  
336 mass changes on the avian T-cell-mediated immune response. *Canadian Journal of*  
337 *Zoology*, 79(1), 101–105. <https://doi.org/10.1139/z00-190>
- 338 Alonso-Alvarez, C., Doutrelant, C., & Sorci, G. (2004). Ultraviolet reflectance affects male-  
339 male interactions in the blue tit (*Parus caeruleus ultramarinus*). *Behavioral Ecology*,  
340 15(5), 805–809. <https://doi.org/10.1093/beheco/arh083>
- 341 Amundsen, T. (2000). Why are female birds ornamented? *Trends in Ecology & Evolution*,  
342 15(4), 149–155.
- 343 Andersson, S., Ornborg, J., & Andersson, M. (1998). Ultraviolet sexual dimorphism and  
344 assortative mating in blue tits. *Proceedings of the Royal Society of London B: Biological*  
345 *Sciences*, 265(1395), 445–450. <https://doi.org/10.1098/rspb.1998.0315>
- 346 Beck, M. L., Hopkins, W. A., & Hawley, D. M. (2015). Relationships among plumage  
347 coloration, blood selenium concentrations and immune responses of adult and nestling  
348 tree swallows. *Journal of Experimental Biology*, 218(21), 3415–3424.  
349 <https://doi.org/10.1242/jeb.123794>
- 350 Birkhead, T. R., Pellatt, E. J., Matthews, I. M., Roddis, N. J., Hunter, F. M., McPhie, F., &  
351 Castillo-Juarez, H. (2006). Genic capture and the genetic basis of sexually selected traits  
352 in the zebra finch. *Evolution*, 60(11), 2389–2398.
- 353 Bonneaud, C., Sinsheimer, J. S., Richard, M., Chastel, O., & Sorci, G. (2009). Mhc  
354 polymorphisms fail to explain the heritability of phytohaemagglutinin- induced skin  
355 swelling in a wild passerine. *Biology Letters*, 5(6), 784–787.  
356 <https://doi.org/10.1098/rsbl.2009.0435>
- 357 Brommer, J. E., Alho, J. S., Biard, C., Chapman, J. R., Charmantier, A., Dreiss, A., ...  
358 Winkel, W. (2010). Passerine extrapair mating dynamics: a bayesian modeling approach  
359 comparing four species. *The American Naturalist*, 176(2), 178–187.  
360 <https://doi.org/10.1086/653660>
- 361 Brommer, J. E., & Klueen, E. (2012). Exploring the genetics of nestling personality traits in a  
362 wild passerine bird: testing the phenotypic gambit. *Ecology and Evolution*, 2(12), 3032–  
363 3044. <https://doi.org/10.1002/ece3.412>
- 364 Butler, D. G., Cullis, B. R., Gilmour, A. R., & Gogel, B. J. (2009). ASReml-R reference  
365 manual. Department of Primary Industries and Fisheries, State of Queensland, Australia.
- 366 Charmantier, A., Wolak, M. E., Grégoire, A., Fargevieille, A., & Doutrelant, C. (2017).  
367 Colour ornamentation in the blue tit : quantitative genetic (co)variances across sexes.  
368 *Heredity*, 118, 125–134. <https://doi.org/10.1038/hdy.2016.70>

- 369 Charmantier, A., & Réale, D. (2005). How do misassigned paternities affect the estimation of  
370 heritability in the wild? *Molecular Ecology*, *14*, 2839–2850.  
371 <https://doi.org/10.1111/j.1365-294X.2005.02619.x>
- 372 Cichoń, M., & Dubiec, A. (2005). Cell-mediated immunity predicts the probability of local  
373 recruitment in nestling blue tits. *Journal of Evolutionary Biology*, *18*(4), 962–966.  
374 <https://doi.org/10.1111/j.1420-9101.2005.00910.x>
- 375 Coster, A.(2012). pedigree: Pedigree functions. R package version 1.4.[https://CRAN.R-](https://CRAN.R-project.org/package=pedigree)  
376 [project.org/package=pedigree](https://CRAN.R-project.org/package=pedigree)
- 377 Doutrelant, C., Gregoire, A., Grnac, N., Gomez, D., Lambrechts, M., & Perret, P. (2008).  
378 Female coloration indicates female reproductive capacity in blue tits. *Journal of*  
379 *Evolutionary Biology*, *21*, 226–233. <https://doi.org/10.1111/j.1420-9101.2007.01451.x>
- 380 Doutrelant, C., Grégoire, A., Midamegbe, A., Lambrechts, M., & Perret, P. (2012). Female  
381 plumage coloration is sensitive to the cost of reproduction. An experiment in blue tits.  
382 *Journal of Animal Ecology*, *81*(1), 87–96. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2656.2011.01889.x)  
383 [2656.2011.01889.x](https://doi.org/10.1111/j.1365-2656.2011.01889.x)
- 384 Drobniak, S. M., Wiejaczka, D., Arct, A., Dubiec, A., & Gustafsson, L. (2013). Low cross-  
385 sex genetic correlation in carotenoid-based plumage traits in the blue tit nestlings  
386 (*Cyanistes caeruleus*). *PLoS ONE*, *8*(7), e69786.  
387 <https://doi.org/10.1371/journal.pone.0069786>
- 388 Evans, S. R., & Sheldon, B. C. (2012). Quantitative genetics of a carotenoid-based color :  
389 heritability and persistent natal environmental effects in the great tit. *The American*  
390 *Naturalist*, *1*, 79–94. <https://doi.org/10.1086/663198>
- 391 Figuerola, J., Senar, J. C., & Pascual, J. (1999). The use of a colorimeter in field studies of  
392 blue tit *Parus caeruleus* coloration. *Ardea*, *87*(2), 269–275.
- 393 Fischer, T. M., Gilmour, A. R., & Werf, J. H. Van Der. (2004). Computing approximate  
394 standard errors for genetic parameters derived from random regression models fitted by  
395 average information REML. *Genetics, Selection, Evolution : GSE*, *36*(3), 363–369.  
396 <https://doi.org/10.1186/1297-9686-36-3-363>
- 397 Galván, I. (2011). Ultraviolet-blue plumage colouration can be perceived as an indicator of  
398 fluctuating asymmetry by blue tits (*Cyanistes caeruleus*). *Journal of Ornithology*, *152*(2),  
399 223–230. <https://doi.org/10.1007/s10336-010-0568-y>
- 400 Hadfield, J. D., Nutall, A., Osorio, D., & Owens, I. P. F. (2007). Testing the phenotypic  
401 gambit: phenotypic, genetic and environmental correlations of colour. *Journal of*  
402 *Evolutionary Biology*, *20*(2), 549–557. [https://doi.org/10.1111/j.1420-](https://doi.org/10.1111/j.1420-9101.2006.01262.x)  
403 [9101.2006.01262.x](https://doi.org/10.1111/j.1420-9101.2006.01262.x)
- 404 Hadfield, J. D., Burgess, M. D., Lord, A., Phillimore, A. B., Clegg, S. M., & Owens, I. P. F.  
405 (2006). Direct versus indirect sexual selection : genetic basis of colour, size and  
406 recruitment in a wild bird. *Proceedings of the Royal Society of London B: Biological*  
407 *Sciences*, *273*, 1347–1353. <https://doi.org/10.1098/rspb.2005.3459>

- 408 Hamilton, W. D., & Zuk, M. (1982). Heritable true fitness and bright birds: a role for  
409 parasites? *Science*, *218*, 384–386.
- 410 Hõrak, P., Tegelmann, L., Ots, I., & Møller, A. P. (1999). Immune function and survival of  
411 great tit nestlings in relation to growth conditions. *Oecologia*, *121*, 316–322.  
412 [https://doi.org/10.1007/978-3-642-38865-1\\_45](https://doi.org/10.1007/978-3-642-38865-1_45)
- 413 Hunt, S., Bennett, A. T. D., Cuthill, I. C., & Griffiths, R. (1998). Blue tits are ultraviolet tits.  
414 *Proceedings of the Royal Society of London B: Biological Sciences*, *265*, 451–455.
- 415 Hunt, S., Cuthill, I. C., Bennett, A. T. D., & Griffiths, R. (1999). Preferences for ultraviolet  
416 partners in the blue tit. *Animal Behaviour*, *58*(4), 809–815.  
417 <https://doi.org/10.1006/anbe.1999.1214>
- 418 Jacot, A., & Kempenaers, B. (2007). Effects of nestling condition on UV plumage traits in  
419 blue tits: An experimental approach. *Behavioral Ecology*, *18*(1), 34–40.  
420 <https://doi.org/10.1093/beheco/arl054>
- 421 Johnsen, A., Delhey, K., Andersson, S., & Kempenaers, B. (2003). Plumage colour in nestling  
422 blue tits : sexual dichromatism, condition dependence and genetic effects. *Proceedings of*  
423 *the Royal Society of London B: Biological Sciences*, *270*, 1263–1270.  
424 <https://doi.org/10.1098/rspb.2003.2375>
- 425 Jones, A. G., & Ratterman, N. L. (2009). Mate choice and sexual selection : What have we  
426 learned since Darwin ? *Proceedings in the National Academy of Sciences*, *106*(S1),  
427 10001–10008.
- 428 Kim, S.-Y., Fargallo, J. A., Vergara, P., & Martínez-Padilla, J. (2013). Multivariate heredity  
429 of melanin-based coloration, body mass and immunity. *Heredity*, *111*, 139–146.  
430 <https://doi.org/10.1038/hdy.2013.29>
- 431 Kinnard, T. B., & Westneat, D. F. (2009). Phenotypic and genetic variance of house sparrows  
432 (*Passer domesticus*) early in development. *The Auk*, *126*(4), 884–895.  
433 <https://doi.org/10.1525/auk.2009.08119>
- 434 Kirkpatrick, M., & Barton, N. H. (1997). The strength of indirect selection on female mating  
435 preferences. *Proceedings of the National Academy of Sciences of the United States of*  
436 *America*, *94*(4), 1282–1286. <https://doi.org/VL-94>
- 437 Kruuk, L. E. B., & Hadfield, J. D. (2007). How to separate genetic and environmental causes  
438 of similarity between relatives. *Journal of Evolutionary Biology*, *20*(5), 1890–1903.  
439 <https://doi.org/10.1111/j.1420-9101.2007.01377.x>
- 440 Linden, M., Gustafsson, L., & Part, T. (1992). Selection of fledging mass in the collared  
441 flycatcher and the great tit. *Ecology*. <https://doi.org/10.2307/1938745>
- 442 López-Rull, I., Celis, P., Salaberria, C., Puerta, M., & Gil, D. (2011). Post-fledging  
443 recruitment in relation to nestling plasma testosterone and immunocompetence in the  
444 spotless starling. *Functional Ecology*, *25*(3), 500–508. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2435.2010.01783.x)  
445 [2435.2010.01783.x](https://doi.org/10.1111/j.1365-2435.2010.01783.x)

- 446 Merilä, J., Kruuk, L. E. B., & Sheldon, B. C. (2001). Natural selection on the genetical  
447 component of variance in body condition in a wild bird population. *Journal of*  
448 *Evolutionary Biology*, *14*, 918–929.
- 449 Merilä, J., & Sheldon, B. C. (2000). Lifetime reproductive success and heritability in nature.  
450 *The American Naturalist*, *155*(3), 301–310. <https://doi.org/10.1086/303330>
- 451 Monrós, J. S., Belda, E. J., & Barba, E. (2002). Post-fledging survival of individual great tits :  
452 the effect of hatching date and fledging mass. *Oikos*, *99*, 481–488.
- 453 Moreno, J., Merino, S., Sanz, J. J., Arriero, E., Morales, J., & Tomás, G. (2005). Nestling  
454 cell-mediated immune response, body mass and hatching date as predictors of local  
455 recruitment in the pied flycatcher *Ficedula hypoleuca*. *Journal of Avian Biology*, *36*(3),  
456 251–260. <https://doi.org/10.1111/j.0908-8857.2005.03413.x>
- 457 Morrissey, M. (2018). pedantics: Functions to Facilitate Power and Sensitivity Analyses for  
458 Genetic Studies of Natural Populations. R package version 1.7. [https://CRAN.R-](https://CRAN.R-project.org/package=pedantics)  
459 [project.org/package=pedantics](https://CRAN.R-project.org/package=pedantics)
- 460 Norris, K. (1993). Heritable variation in a plumage indicator of viability in male great tits  
461 (*Parus major*). *Nature*, *362*, 537–539.
- 462 Perrins, C. M. (1965). Population Fluctuations and Clutch-Size in the Great Tit, *Parus major*  
463 *L.* *Journal of Animal Ecology*, *34*(3), 601–647.
- 464 Peters, A., Delhey, K., Johnsen, A., & Kempenaers, B. (2007). The condition-dependent  
465 development of carotenoid-based and structural plumage in nestling blue tits : males and  
466 females differ. *The American Naturalist*, *169*, S122–S136.
- 467 Pickett, S. R. A., Weber, S. B., McGraw, K. J., Norris, K. J., & Evans, M. R. (2013).  
468 Environmental and parental influences on offspring health and growth in great tits  
469 (*Parus major*). *PLoS ONE*, *8*(7), e69695. <https://doi.org/10.1371/journal.pone.0069695>
- 470 Pitala, N., Gustafsson, L., Sendecka, J., & Brommer, J. E. (2007). Nestling immune response  
471 to phytohaemagglutinin is not heritable in collared flycatchers. *Biology Letters*, *3*, 418–  
472 421. <https://doi.org/10.1098/rsbl.2007.0135>
- 473 Qvarnström, A., Brommer, J. E., & Gustafsson, L. (2006). Evolution of mate choice in the  
474 wild. *Nature*, *441*, 84–86. <https://doi.org/10.1038/nature05501>
- 475 Radersma, R., Komdeur, J., & Tinbergen, J. M. (2015). Early morning fledging improves  
476 recruitment in Great Tits *Parus major*. *Ibis*, *157*(2), 351–355.  
477 <https://doi.org/10.1111/ibi.12230>
- 478 Radersma, R., Tinbergen, J. M., & Komdeur, J. (2011). Do brood sex ratio, nestling  
479 development and sex affect fledging timing and order? An experimental study on great  
480 tits. *Animal Behaviour*, *81*(1), 69–75. <https://doi.org/10.1016/j.anbehav.2010.09.007>

- 481 Reid, J. M. (2014). Quantitative genetic approaches to understanding sexual selection and  
482 mating system evolution in the wild. In A. Charmantier, D. Garant, & L. E. B. Kruuk  
483 (Eds.), *Quantitative Genetics in the Wild* (pp. 34–53). Oxford: Oxford University Press.
- 484 Remeš, V., & Matysioková, B. (2013). More ornamented females produce higher-quality  
485 offspring in a socially monogamous bird : an experimental study in the great tit (*Parus*  
486 *major*). *Frontiers in Zoology*, *10*(14), 1–10.
- 487 Rémy, A., Grégoire, A., Perret, P., & Doutrelant, C. (2010). Mediating male-male  
488 interactions: The role of the UV blue crest coloration in blue tits. *Behavioral Ecology*  
489 *and Sociobiology*, *64*(11), 1839–1847. <https://doi.org/10.1007/s00265-010-0995-z>
- 490 Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of*  
491 *Statistical Software*, *48*(2), 1-36. <http://www.jstatsoft.org/v48/i02/>.
- 492 Sakaluk, S. K., Wilson, A. J., Bowers, E. K., Johnson, L. S., Masters, B. S., Johnson, B. G.,  
493 ... Thompson, C. F. (2014). Genetic and environmental variation in condition, cutaneous  
494 immunity, and haematocrit in house wrens. *BMC Evolutionary Biology*, *14*, 242.  
495 <https://doi.org/10.1186/s12862-014-0242-8>
- 496 Senar, J. C., Figuerola, J., & Pascual, J. (2002). Brighter yellow blue tits make better parents.  
497 *Proceedings of the Royal Society of London B: Biological Sciences*, *269*, 257–261.  
498 <https://doi.org/10.1098/rspb.2001.1882>
- 499 Siefferman, L., & Hill, G. E. (2007). The effect of rearing environment on blue structural  
500 coloration of eastern bluebirds (*Sialia sialis*). *Behavioral Ecology and Sociobiology*,  
501 *61*(12), 1839–1846. <https://doi.org/10.1097/IOP.0b013e3181d3da43>.Risk
- 502 Siefferman, L., Shawkey, M. D., Bowman, R., & Woolfenden, G. E. (2008). Juvenile  
503 coloration of Florida scrub-jays (*Aphelocoma coerulescens*) is sexually dichromatic and  
504 correlated with condition. *Journal of Ornithology*, *149*(3), 357–363.  
505 <https://doi.org/10.1007/s10336-008-0289-7>
- 506 Smits, J. E., Bortolotti, G. R., & Tella, J. L. (1999). Simplifying the phytohaemagglutinin  
507 skin-testing technique in studies of avian immunocompetence. *Functional Ecology*,  
508 *13*(4), 567–572. <https://doi.org/10.1046/j.1365-2435.1999.00338.x>
- 509 Svensson, L. (1992). *Identification guide to European passerines*. Stockholm.
- 510 Thompson, C. F., Sakaluk, S. K., Masters, B. S., Johnson, B. G. P., Vogel, L. A., Forsman, A.  
511 M., & Johnson, L. S. (2013). Condition-dependent sex difference in nestling house wren  
512 (*Troglodytes aedon*) response to phytohaemagglutinin injection. *Canadian Journal of*  
513 *Zoology*, *92*, 1–7. <https://doi.org/10.1139/cjz-2013-0140>
- 514 Tinbergen, J., & Boerlijst, M. (1990). Nestling weight and survival in individual great tits  
515 (*Parus major*). *Journal of Animal Ecology*, *59*(3), 1113–1127.
- 516 Velando, A., Torres, R., & Espinosa, I. (2005). Male coloration and chick condition in blue-  
517 footed booby : a cross-fostering experiment. *Behavioral Ecology and Sociobiology*, *58*,  
518 175–180. <https://doi.org/10.1007/s00265-005-0911-0>

- 519 Verboven, N., & Visser, M. E. (1998). Seasonal variation in local recruitment of great tits:  
520 The Importance of Being Early. *Oikos*, *81*, 511–524. <https://doi.org/10.2307/3546771>
- 521 Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., ...  
522 Nussey, D. H. (2009). An ecologist's guide to the animal model. *The Journal of Animal*  
523 *Ecology*, *79*(1), 13–26. <https://doi.org/10.1111/j.1365-2656.2009.01639.x>

524 **Supporting information**

525

526 **Table S1:** Repeatability of each color trait measured by each spectrometer

527 **Figure S1:** Average reflectance spectrum measured by each spectrometer

528 **Figure S2:** Distribution of the six traits measured in nestlings

529 **Figure S3:** Distribution of the univariate animal model residuals for all six traits

530 **Tables S2-S7:** Sex-specific variances and cross-sex covariances estimated by bivariate animal  
531 models for each trait separately

532 **Figure S4:** Additive genetic variances of all six traits and their covariances in males and females

533 **Tables S8-S9:** Genetic correlation matrix estimated by a multivariate mixed model in females  
534 and males respectively.

535 **Table S10:** Cross-sex genetic correlation and SE for each trait

536 **Tables S11-S16:** Fixed and random effects estimated by the animal model for each trait

537 **Tables S17:** Variances of wing web thickness measured on day 13 and day 14 and their  
538 covariances estimated on different levels by a bivariate animal model

539 **Tables S18-S22:** Covariance matrices derived from the multivariate animal model

540 **Figure S5:** Correlations and loadings estimated in the structural equation models

541 **Tables S23:** Phenotypic correlation matrix derived from a multivariate mixed model

542 **Text S1:** R code for performing quantitative genetic analyses

543 **Text S2-S3:** R code for performing structural equation modelling and simulations

544 **Table 1:** Number of individuals, rearing broods, mean and standard deviation (SD) for each  
 545 studied trait.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
n.individuals	2851	2851	2851	1516	3210	3200
n.broods	364	364	364	173	390	389
mean (SD)	9.46 (2.74)	340.78 (10.61)	0.33 (0.02)	0.54 (0.17)	11.40 (1.08)	46.34 (3.39)

546

547 **Table 2:** Animal model estimates (and standard errors) of variance components of all six traits.  
 548 Heritability ( $h^2$ ) and ratio  $V_{CE}/V_P$  (and their standard error) were calculated for each trait, where  
 549  $V_P$  is the sum of all estimated variances, except for color traits where it does not include  $V_{ME}$ .  
 550 For tail color traits,  $V_I$ ,  $V_{CE}$  and  $V_{ME}$  were estimated separately for each spectrophotometer used  
 551 (1 and 2 respectively) and thus two  $h^2$  and ratios  $V_{CE}/V_P$  were calculated.

		Brightness	Hue	UV chroma	PHA	Body mass	Wing
$V_A$		0.26 (0.13)	1.2 (2.02)	6.24E-06 (3.93E-06)	5.17E-06 (1.36E-03)	0.25 (0.04)	3.26 (0.46)
$V_{CE}$	1	0.94 (0.19)	11.39 (2.52)	1.63E-05 (3.54E-06)	7.83E-06 (1.21E-06)	0.49 (0.04)	4.87 (0.57)
	2	1.68 (0.21)	24.11 (3.06)	1.35E-04 (1.78E-05)			
$V_I$	1	1.43 (0.16)	19.66 (2.76)	2.84E-05 (3.79E-06)			
	2	2.89 (0.17)	40.32 (2.75)	3.32E-04 (1.58E-05)			
$V_{ME}$	1	2.32 (0.06)	82.47 (2.12)	4.83E-05 (3.24E-06)			
	2	6.55 (0.11)	122.6 (2.06)	5.51E-04 (9.25E-06)			
$V_R$					1.30E-02 (1.02E-03)	0.19 (0.02)	3.46 (0.35)
$h^2$	1	0.10 (0.05)	0.04 (0.06)	0.12 (0.08)	0.19 (0.06)	0.27 (0.04)	0.28 (0.05)
	2	0.05 (0.03)	0.02 (0.03)	0.01 (0.008)			
$V_{CE}/V_P$	1	0.36 (0.05)	0.35 (0.06)	0.32 (0.05)	0.30 (0.04)	0.52 (0.02)	0.42 (0.03)
	2	0.35 (0.03)	0.37 (0.03)	0.29 (0.03)			

552

553 **Table 3:** Additive genetic correlation (and standard error) matrix estimated by the multivariate  
 554 animal model. Correlations between color measures and performance traits are highlighted in  
 555 grey.

Brightness	Hue	UV chroma	PHA	Body mass	Wing
-0.72 (0.40)					
-0.13 (0.41)	-0.34 (0.54)				
-0.05 (0.25)	-0.04 (0.30)	-0.50 (0.33)			
0.21 (0.20)	0.13 (0.23)	-0.17 (0.28)	0.23 (0.16)		
0.50 (0.24)	-0.51 (0.28)	-0.35 (0.35)	0.32 (0.19)	0.26 (0.13)	


556


## Supplementary material belonging to the article


## “Tail color signals performance in blue tit nestlings”

**Table S1:** Repeatability of each color measured by each spectrometer

Trait	Repeatability 1 (95% CI)	Repeatability 2 (95% CI)
brightness	0.51 (0.48-0.55)	0.40 (0.38-0.43)
hue	0.28 (0.32-0.25)	0.33 (0.31-0.36)
UV chroma	0.50 (0.47-0.54)	0.43 (0.41-0.45)

**Figure S1:** Average reflectance spectra from nestlings' tail feathers in 2005-2006 (first spectrometer, red line), and in 2003+2007-2009 (second spectrometer, blue line). The dotted vertical line represents the peak reflectance for both spectrophotometers.

**Figure S2:** Distribution of the six traits (averages per individual) measured in nestlings.

**Figure S3:** Distribution of the univariate animal model residuals for all six traits

**Table S2:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with tail brightness in males and in females as two response variables. Common environment and residual error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Component	Parameter	Estimate	SE	z.ratio
Common environment 1	$V_m$	1.12	0.26	4.40
	$COV_{mf}$	0.90	0.20	4.49
	$V_f$	0.78	0.23	3.35
Common environment 2	$V_m$	1.50	0.27	5.60
	$COV_{mf}$	1.68	0.24	6.99
	$V_f$	2.00	0.34	5.85
Additive genetic	$V_m$	0.22	0.19	1.21
	$COV_{mf}$	0.31	0.17	1.85
	$V_f$	0.45	0.29	1.56
Residual 1	$V_m$	1.65	0.21	7.76
	$V_f$	2.08	0.31	6.72
Residual 2	$V_m$	4.36	0.26	16.57
	$V_f$	4.89	0.33	14.94

**Table S3:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with tail hue in males and in females as two response variables. Common environment and residual error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Component	Parameter	Estimate	SE	z.ratio
Common environment 1	$V_m$	11.81	4.17	2.83
	$COV_{mf}$	12.38	3.38	3.67
	$V_f$	17.87	4.69	3.81
Common environment 2	$V_m$	29.72	4.96	5.99
	$COV_{mf}$	24.48	3.78	6.48
	$V_f$	20.29	4.68	4.34
Additive genetic	$V_m$	4.96	4.96	1.00
	$COV_{mf}$	4.94	4.00	1.24
	$V_f$	14.25	5.58	2.56
Residual 1	$V_m$	49.27	6.09	8.09
	$V_f$	25.05	5.32	4.71
Residual 2	$V_m$	71.25	4.96	14.36
	$V_f$	80.45	5.79	13.90

**Table S4:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with tail UV chroma in males and in females as two response variables. Common environment and residual error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Component	Parameter	Estimate	SE	z.ratio
Common environment 1	$V_f$	1.26E-05	3.95E-06	3.20
	$COV_{mf}$	1.67E-05	3.83E-06	4.35
	$V_m$	2.20E-05	5.62E-06	3.91
Common environment 2	$V_f$	1.34E-04	2.64E-05	5.07
	$COV_{mf}$	1.43E-04	2.09E-05	6.84
	$V_m$	1.58E-04	2.65E-05	5.96
Additive genetic	$V_f$	1.21E-05	7.45E-06	1.63
	$COV_{mf}$	1.21E-05	5.75E-06	2.11
	$V_m$	1.22E-05	8.14E-06	1.50
Residual 1	$V_f$	2.75E-05	6.23E-06	4.42
	$V_m$	3.63E-05	6.98E-06	5.20
Residual 2	$V_f$	5.02E-04	2.70E-05	18.58
	$V_m$	4.29E-04	2.32E-05	18.53

**Table S5:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with PHA in males and in females as two response variables.

Component	Parameter	Estimate	SE	z.ratio
Common environment	$V_f$	6.16E-03	1.38E-03	4.45
	$COV_{mf}$	7.54E-03	1.29E-03	5.85
	$V_m$	9.88E-03	1.71E-03	5.77
Additive genetic	$V_f$	7.47E-03	2.15E-03	3.47
	$COV_{mf}$	5.71E-03	1.57E-03	3.64
	$V_m$	4.49E-03	1.90E-03	2.37
Residual	$V_f$	1.15E-02	1.60E-03	7.22
	$V_m$	1.30E-02	1.56E-03	8.31

**Table S6:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with body mass in males and in females as two response variables.


Component	Parameter	Estimate	SE	z.ratio
Common environment	$V_f$	0.44	0.04	10.25
	$COV_{mf}$	0.43	0.04	11.08
	$V_m$	0.43	0.04	10.02
Additive genetic	$V_f$	0.24	0.04	5.69
	$COV_{mf}$	0.24	0.04	6.44
	$V_m$	0.28	0.05	6.09
Residual	$V_f$	0.15	0.03	5.65
	$V_m$	0.17	0.03	5.83


**Table S7:** Variances and covariances, their standard error (SE), and z ratio, estimated by a bivariate animal model with wing length in males and in females as two response variables.

Component	Parameter	Estimate	SE	z.ratio
Common environment	$V_f$	4.94	0.50	9.81
	$COV_{mf}$	4.79	0.46	10.47
	$V_m$	5.08	0.53	9.53
Additive genetic	$V_f$	1.86	0.47	4.00
	$COV_{mf}$	2.02	0.44	4.60
	$V_m$	2.46	0.61	4.03
Residual	$V_f$	3.31	0.33	9.94
	$V_m$	3.89	0.43	8.97


**Figure S4: A)** Additive genetic variances and their 95%CI in both sexes for all six traits and **B)** additive genetic covariances between these traits. These (co)variances and their uncertainties were estimated by a multivariate animal model for each sex separately.

### A) Variances



**B) Covariances (part 1)**

## B) Covariances (part 2)



**Table S8:** Genetic correlation matrix estimated by a multivariate mixed model **in females**. In this model, fixed effects were similar to fixed effects fitted in the multivariate animal model and only the residual component was estimated. Correlations (and their standard error) are in the lower triangle and the diagonal (underlined numbers) contains phenotypic variances (and their standard error) estimated by the model. Correlations between color measures and performance traits are highlighted in grey.

Brightness	Hue	UV chroma	PHA	Body mass	Wing
-0.59 (0.61)					
-0.30 (0.71)	-0.33 (0.73)				
0.04 (0.35)	-0.09 (0.36)	-0.55 (0.38)			
0.28 (0.30)	0.30 (0.30)	-0.50 (0.37)	0.33 (0.18)		
0.61 (0.38)	-0.52 (0.39)	-0.40 (0.48)	0.38 (0.24)	0.25 (0.18)	

**Table S9:** Genetic correlation matrix estimated by a multivariate mixed model **in males**. In this model, fixed effects were similar to fixed effects fitted in the multivariate animal model and only the residual component was estimated. Correlations (and their standard error) are in the lower triangle and the diagonal (underlined numbers) contains phenotypic variances (and their standard error) estimated by the model. Correlations between color measures and performance traits are highlighted in grey.

Brightness	Hue	UV chroma	PHA	Body mass	Wing
-0.79 (0.67)					
-0.24 (0.73)	-0.22 (1.09)				
-0.10 (0.41)	-0.02 (0.55)	-0.28 (0.62)			
0.15 (0.27)	-0.10 (0.35)	0.05 (0.44)	0.45 (0.23)		
0.61 (0.34)	-0.56 (0.42)	-0.38 (0.54)	0.21 (0.27)	0.4 (0.17)	

**Table S10:** Cross-sex genetic correlation and SE for each trait

Trait	$R_{g_{mf}}$	SE
Brightness	0.99	0.53
Hue	0.59	0.48
UV chroma	0.95	0.52
PHA	0.98	0.17
Body mass	0.92	0.05
Wing length	0.94	0.08

**Table S11:** Fixed and random effects (and their standard error SE) estimated by the animal model for tail brightness. The statistical significance of additive genetic variance was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter. Common environment, individual, and measurement error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment 1	0.94	0.19		
Common environment 2	1.68	0.21		
Additive genetic	0.26	0.13	$\chi^2=6.19$	0.01
Individual 1	1.43	0.16		
Individual 2	2.89	0.17		
Measurement error 1	2.32	0.06		
Measurement error 2	6.55	0.11		
<b>Fixed effects</b>				
Intercept	6.85	0.44	$F_{1,287.2}=842.20$	<0.001
Vane	0.28	0.02	$F_{1,1894.0}=324.60$	<0.001
Year			$F_{5,287.63} = 36.18$	<0.001
	2005	1.15		
	2006	-0.65		
	2007	-1.91		
	2008	-0.60		
	2009	-1.70		
Sex			$F_{2,2308.1} = 9.58$	<0.001
	female	0.16		
	male	-0.18		

**Table S12:** Fixed and random effects (and their standard error SE) estimated by the animal model for tail hue. The statistical significance of additive genetic variance was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter. Common environment, individual, and measurement error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment 1	11.39	2.52		
Common environment 2	24.11	3.06		
Additive genetic	1.20	2.02	$\chi^2=0.20$	0.65
Individual 1	19.66	2.76		
Individual 2	40.32	2.75		
Measurement error 1	82.47	2.12		
Measurement error 2	122.60	2.06		
<b>Fixed effects</b>				
Intercept	354.16	1.68	$F_{1,219.7}=1.89E+05$	<0.001
Vane	-0.44	0.06	$F_{1,1629.8}=52.1$	<0.001
Year			$F_{5,283.7}=24.09$	<0.001
	2005	-4.93	1.82	
	2006	-10.89	1.79	
	2007	-10.28	1.85	
	2008	-9.79	1.79	
	2009	-4.65	1.79	
Sex			$F_{2,2387.1}=5.98$	0.002
	female	-0.42	0.79	
	male	-1.43	0.78	

**Table S13:** Fixed and random effects (and their standard error SE) estimated by the animal model for tail UV chroma. The statistical significance of additive genetic variance was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter. Common environment, individual, and measurement error variances were estimated separately for each spectrophotometer (1: feathers from 2005-2006, 2: feathers from 2003, 2007-2009).

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment 1	1.63E-05	3.54E-06		
Common environment 2	1.35E-04	1.78E-05		
Additive genetic	6.24E-06	3.93E-06	$\chi^2=4.31$	0.04
Individual 1	2.84E-05	3.79E-06		
Individual 2	3.32E-04	1.58E-05		
Measurement error 1	4.83E-05	1.24E-06		
Measurement error 2	5.51E-04	9.25E-06		
<b>Fixed effects</b>				
Intercept	3.32E-01	3.85E-03	$F_{1,149.6}=72020.00$	<0.001
Vane	-3.58E-04	8.97E-05	$F_{1,779.1}=15.92$	<0.001
Year			$F_{5,295.7}=66.67$	<0.001
	2005	-1.74E-02	3.95E-03	
	2006	-1.30E-02	3.93E-03	
	2007	9.45E-03	4.29E-03	
	2008	1.79E-03	4.14E-03	
	2009	6.58E-05	4.10E-03	
Sex			$F_{2,1083.4}=157.10$	<0.001
	female	-1.93E-03	1.20E-03	
	male	6.10E-03	1.19E-03	

**Table S14:** Fixed and random effects (and their standard error SE) estimated by the animal model for PHA response. The statistical significance of random effects was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter.

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment	7.83E-03	1.21E-03	$\chi^2=153.09$	<0.001
Additive genetic	5.17E-03	1.36E-03	$\chi^2=46.48$	<0.001
Residual	1.30E-02	1.02E-03		
<b>Fixed effects</b>				
Intercept	0.49	0.03	$F_{1,196.7} = 3920.0$	<0.001
Year			$F_{3,207.9} = 6.52$	<0.001
	2005	0.04		
	2006	0.06		
	2007	-0.02		
Sex			$F_{1,1422.8} = 1.69$	0.85
	female	0.035		
	male	0.037		

**Table S15:** Fixed and random effects (and their standard error SE) estimated by the animal model for body mass. The statistical significance of random effects was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter.

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment	0.49	0.04	$\chi^2=1093.62$	<0.001
Additive genetic	0.25	0.04	$\chi^2=185.67$	<0.001
Residual	0.19	0.02		
<b>Fixed effects</b>				
Intercept	8.51	0.27	$F_{1,494.3} = 2401.00$	<0.001
Tarsus	0.13	0.01	$F_{1,3107.0} = 184.50$	<0.001
Year			$F_{5,404.4} = 12.95$	<0.001
	2005	0.42		
	2006	0.68		
	2007	0.17		
	2008	0.85		
	2009	0.92		
Sex			$F_{1,2928.2} = 213.00$	<0.001
	female	-0.27		
	male	0.21		

**Table S16:** Fixed and random effects (and their standard error SE) estimated by the animal model for wing length. The statistical significance of random effects was tested using LRT with 1df. The statistical significance of fixed effects was tested using conditional Wald-F tests. Coefficients of years and sex effects are reported as contrasts to year 2003 for the former and to unsexed individuals for the latter.

Effect	Estimate	SE	Test statistic	p.value
<b>Random effects</b>				
Common environment	4.87	0.46	$\chi^2=616.89$	<0.001
Additive genetic	3.26	0.57	$\chi^2=64.74$	<0.001
Residual	3.46	0.35		
<b>Fixed effects</b>				
Intercept	44.56	0.69	$F_{1,435.6} = 97880.00$	<0.001
Year			$F_{5,404.0} = 7.34$	<0.001
	2005	1.31		
	2006	2.85		
	2007	0.44		
	2008	1.43		
	2009	0.84		
Sex			$F_{1,2953.4} = 25.18$	<0.001
	female	0.09		
	male	0.72		

**Table S17:** Variances of wing web thickness measured on day 13 and day 14 and their covariances estimated on different levels by a bivariate animal model (and their standard errors (SE)). In this model, year and sex were fitted as fixed effects for each response separately and random effects included common environment effects, individual identity, additive genetic effects and measurement error. Measurement error covariance was not fitted as both measures were not taken at the same time. Variance of the PHA response on each level can then be estimated as the sum of both variances minus twice their covariance and heritability is calculated as  $VA/VP$  where  $VP$  does not include measurement error.

<b>(Co)Variance component</b>	<b>Estimate</b>	<b>SE</b>
Common environment: WingWeb_d13	3.89	0.51
Common environment:WingWeb_d14:WingWeb_d13	7.28	1.25
Common environment: WingWeb_d14	33.77	4.87
Additive genetic: WingWeb_d13	0.52	0.23
Additive genetic: WingWeb_d14:WingWeb_d13	-1.16	0.64
Additive genetic: WingWeb_d14	11.97	3.62
Individual: WingWeb_d13	3.89	0.23
Individual: WingWeb_d14:WingWeb_d13	2.29	0.61
Individual: WingWeb_d14	55.62	3.28
Measurement error: WingWeb_d13	0.71	0.03
Measurement error : WingWeb_d14	3.24	0.08
Common environment: PHA	23.10	3.77
Additive genetic: PHA	14.82	3.98
Individual: PHA	54.94	3.41

**Table S18:** Additive genetic covariance matrix derived from the multivariate animal model. The diagonal contains additive genetic variances and standard errors are printed below each estimate in grey. Covariances between color measures and performance traits are highlighted in yellow.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
Brightness	0.28 0.12					
Hue	-0.80 0.44	4.44 2.90				
UV chroma	-1.60E-04 4.93E-04	-1.63E-03 2.59E-03	5.19E-06 3.58E-06			
PHA	-1.87E-03 9.72E-03	-0.01 0.05	-8.30E-05 5.47E-05	5.32E-03 1.31E-03		
Body mass	0.05 0.05	0.13 0.23	-1.86E-04 3.04E-04	8.15E-03 5.46E-03	0.23 0.03	
Wing	0.34 0.17	-1.42 0.78	-1.01E-03 1.04E-03	3.05E-02 1.81E-02	0.16 0.08	1.73 0.37

**Table S19:** First common environment (CE1) covariance matrix derived from the multivariate animal model. The diagonal contains CE1 variances and standard errors are printed below each estimate in grey. Covariances between color measures and performance traits are highlighted in yellow.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
Brightness	1.04 0.21					
Hue	-0.42 0.57	12.93 3.00				
UV chroma	-1.50E-03 6.52E-04	-4.17E-03 2.43E-03	1.74E-05 3.69E-06			
PHA	0.04 0.02	-0.01 0.06	7.64E-05 6.53E-05	1.12E-02 2.18E-03		
Body mass	0.34 0.11	0.31 0.41	9.01E-04 4.58E-04	5.25E-02 0.01	0.60 0.10	
Wing	1.50 0.43	1.21 1.53	1.80E-04 1.70E-03	1.41E-01 0.04	1.18 0.30	7.92 1.40

**Table S20:** Second common environment (CE2) covariance matrix derived from the multivariate animal model. The diagonal contains CE1 variances and standard errors are printed below each estimate in grey. Covariances between color measures and performance traits are highlighted in yellow.

	Brightness	Hue	UV chroma	PHA	Mass	Wing
Brightness	1.95 0.24					
Hue	-1.52 0.65	25.09 3.43				
UV chroma	-1.16E-02 1.77E-03	-1.74E-02 5.91E-03	1.44E-04 1.84E-05			
PHA	0.01 0.02	3.43E-03 5.84E-02	-1.01E-04 1.30E-04	3.45E-03 9.62E-04		
Mass	0.32 0.08	-0.09 0.29	-2.20E-03 6.70E-04	1.89E-02 6.11E-03	0.46 0.05	
Wing	1.51 0.25	-2.00 0.90	-7.99E-03 2.09E-03	3.65E-03 2.00E-02	0.67 0.11	4.18 0.44

**Table S21:** First residual (RES1) covariance matrix derived from the multivariate animal model. The diagonal contains RES1 variances and standard errors are printed below each estimate in grey. Covariances between color measures and performance traits are highlighted in yellow.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
Brightness	2.02 0.16					
Hue	-0.02 0.54	45.29 3.61				
UV chroma	-1.75E-03 5.50E-04	-1.01E-02 2.81E-03	4.08E-05 3.73E-06			
PHA	-3.96E-03 1.07E-02	0.03 0.05	1.02E-04 5.30E-05	1.44E-02 1.28E-03		
Body mass	0.01 0.05	-0.19 0.26	6.53E-04 2.75E-04	1.51E-02 4.84E-03	0.27 0.03	
Wing	0.41 0.22	1.53 1.06	3.81E-03 1.14E-03	0.05 0.02	0.72 0.10	7.78 0.54


**Table S22:** Second residual (RES2) covariance matrix derived from the multivariate animal model. The diagonal contains RES2 variances and standard errors are printed below each estimate in grey. Covariances between color measures and performance traits are highlighted in yellow.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
Brightness	4.68 0.18					
Hue	0.82 0.55	84.91 3.39				
UV chroma	-2.98E-02 1.34E-03	-0.08 0.01	4.61E-04 1.55E-05			
PHA	8.84E-03 1.51E-02	-0.05 0.06	7.23E-05 1.26E-04	1.14E-02 1.06E-03		
Body mass	0.06 0.04	-0.50 0.18	-3.15E-04 3.30E-04	5.99E-03 3.77E-03	0.19 0.02	
Wing	0.95 0.15	-1.54 0.64	-7.07E-03 1.22E-03	7.42E-03 1.39E-02	0.33 0.05	3.35 0.24

**Table S23:** Phenotypic correlation matrix estimated by a multivariate mixed model. In this model, fixed effects were similar to fixed effects fitted in the multivariate animal model and only the residual component was estimated. Correlations (and their standard error) are in the lower triangle and the diagonal (underlined numbers) contains phenotypic variances (and their standard error) estimated by the model. Correlations between color measures and performance traits are highlighted in grey.

	Brightness	Hue	UV chroma	PHA	Body mass	Wing
Brightness	<u>5.85 (0.15)</u>					
Hue	-0.06 (0.02)	<u>99.5 (2.64)</u>				
UV chroma	-0.59 (0.01)	-0.36 (0.02)	<u>4.5E-04 (1.20E-05)</u>			
PHA	0.07 (0.03)	-0.05 (0.03)	-0.01 (0.03)	<u>2.55E-02 (9.18E-04)</u>		
Body mass	0.19 (0.02)	-0.06 (0.02)	-0.07 (0.02)	0.33 (0.02)	<u>0.90 (0.02)</u>	
Wing	0.32 (0.02)	-0.13 (0.02)	-0.12 (0.02)	0.21 (0.02)	0.41 (0.01)	<u>10.71 (0.27)</u>

**Figure S5:** Correlations (solid arrows) and loadings (dashed arrows) estimated in the structural equation models.



*Text S1: R code for performing the quantitative genetic analyses*

```

#Open data files
pedigree<-read.table("Pedigree_BT.txt",header=T)
Data.means<-read.table("Data_nestlings.txt",header=T)
Data.repeats<-read.table("Data_nestlings_repeats.txt",header=T)

#create the inverse of the relationship matrix from the pedigree file
library(asreml)
ainv<-asreml.Ainverse(pedigree)$ginv

#Pin function to estimate genetic correlations and their SEs based on animal model
estimates
pin<-function (object, transform)
{
  pframe <- as.list(object$gammas)
  names(pframe) <- paste("V", seq(1, length(pframe)), sep = "")
  tvalue <- eval(deriv(transform[[length(transform)]], names(pframe)), pframe)
  X <- as.vector(attr(tvalue, "gradient"))
  X[object$gammas.type == 1] <- 0
  tname <- if (length(transform) == 3)
    transform[[2]]
  else ""
  n <- length(pframe)
  i <- rep(1:n, 1:n)
  j <- sequence(1:n)
  k <- 1 + (i > j)
  Vmat <- object$ai
  se <- sqrt(sum(Vmat * X[i] * X[j] * k))
  data.frame(row.names = tname, Estimate = tvalue, SE = se)
}

#Pin function to estimate heritability and its SE based on animal model estimates
in model with heterogeneous residuals
pin2<-function (object, transform)
{
  pframe <- as.list(object$gammas)
  names(pframe) <- paste("V", seq(1, length(pframe)), sep = "")
  tvalue <- eval(deriv(transform[[length(transform)]], names(pframe)), pframe)
  X <- as.vector(attr(tvalue, "gradient"))
  # X[object$gammas.type == 1] <- 0#Do not run this line
  tname <- if (length(transform) == 3)
    transform[[2]]
  else ""
  n <- length(pframe)
  i <- rep(1:n, 1:n)
  j <- sequence(1:n)
  k <- 1 + (i > j)
  Vmat <- object$ai
  se <- sqrt(sum(Vmat * X[i] * X[j] * k))
  data.frame(row.names = tname, Estimate = tvalue, SE = se)
}

#####
#Estimate heritability of nestling color traits (using repeated measures)
#####

#Prepare data for the models
Data.repeats$Ring<-as.factor(Data.repeats$Ring)
Data.repeats$NestID<-as.factor(Data.repeats$NestID)
Data.repeats$GeneticID<-as.factor(Data.repeats$GeneticID)
Data.repeats$Year<-as.factor(Data.repeats$Year)
Data.repeats$Sex<-as.factor(Data.repeats$Sex)
Data.repeats$machine<-as.factor(Data.repeats$machine)
Data.repeats<-droplevels(Data.repeats)

```

```

#I) Brightness
modelbria<-asreml(fixed=brightness_n~ 1 + Sex +Vane + Year
, random= ~ped(Ring) + at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelbria)$varcomp
wald.asreml(modelbria,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelbria$coefficients$fixed#Coefficients of fixed effects

pin2(modelbria, h2~V3/(V1+V3+V4))#First h2
pin2(modelbria, h2~V3/(V2+V3+V5))#Second h2

pin2(modelbria, CE2~V1/(V1+V3+V4))#First VCE/VP
pin2(modelbria, CE2~V2/(V2+V3+V5))#Second VCE/VP

modelbrib<-asreml(fixed=brightness_n~ 1 + Sex +Vane + Year
, random= ~at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

1-pchisq(2*(modelbria$loglik-modelbrib$loglik),1)#Test if VA in brightness is
significantly different from zero

#II) Hue
modelhuea<-asreml(fixed=hue_n~ 1 + Sex +Vane + Year
, random= ~ped(Ring) + at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit",maxiter=100)

summary(modelhuea)$varcomp
wald.asreml(modelhuea,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelhuea$coefficients$fixed#Coefficients of fixed effects

pin2(modelhuea, h2~V3/(V1+V3+V4))#First h2
pin2(modelhuea, h2~V3/(V2+V3+V5))#Second h2

pin2(modelhuea, CE2~V1/(V1+V3+V4))#First VCE/VP
pin2(modelhuea, CE2~V2/(V2+V3+V5))#Second VCE/VP

modelhueb<-asreml(fixed=hue_n~ 1 + Sex +Vane + Year
, random= ~ at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelhueb)
1-pchisq(2*(modelhuea$loglik-modelhueb$loglik),1)#Test if VA in hue is
significantly different from zero

#III) UV chroma
modelUVa<-asreml(fixed=UV.chrome_n~ 1 + Sex +Vane + Year
, random= ~ped(Ring) + at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelUVa)$varcomp

```

```

wald.asreml(modelUVa,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelUVa$coefficients$fixed#Coefficients of fixed effects

pin2(modelUVa, h2~V3/(V1+V3+V4))#First h2
pin2(modelUVa, h2~V3/(V2+V3+V5))#Second h2

pin2(modelUVa, CE2~V1/(V1+V3+V4))#First VCE/VP
pin2(modelUVa, CE2~V2/(V2+V3+V5))#Second VCE/VP

modelUVb<-asreml(fixed=UV.chrome_n~ 1 + Sex +Vane + Year
, random= ~ at(machine):ide(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.repeats
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

1-pchisq(2*(modelUVa$loglik-modelUVb$loglik),1)#Test if VA in UV chroma is
significantly different from zero

#####
#Estimate heritability of nestling traits (using individual averages)
#####

#Prepare data for the models
Data.means$NestID<-as.factor(Data.means$NestID)
Data.means$GeneticID<-as.factor(Data.means$GeneticID)
Data.means$Year<-as.factor(Data.means$Year)
Data.means$machine<-as.factor(Data.means$machine)
Data.means$Ring<-as.factor(Data.means$Ring)
Data.means$Sex<-as.factor(Data.means$Sex)

# I) Brightness
modelbri<-asreml(fixed=brightness_n~ 1 + Sex +Vane + Year
, random= ~ped(Ring) +at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.means
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelbri)$varcomp
wald.asreml(modelbri,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelbri$coefficients$fixed #Coefficients of fixed effects

pin2(modelbri, h2~V3/(V1+V3+V4))#First h2
pin2(modelbri, h2~V3/(V2+V3+V5))#Second h2

pin2(modelbri, CE~V1/(V1+V3+V4))#First VCE/VP
pin2(modelbri, CE~V2/(V2+V3+V5))#Second VCE/VP

modelbri2<-asreml(fixed=brightness_n~ 1 + Sex +Vane + Year
, random= ~at(machine):NestID
, rcov= ~ at(machine):units
, data=Data.means
, na.method.X="include", na.method.Y="omit")

summary(modelbri2)$varcomp
1-pchisq(2*(modelbri$loglik-modelbri2$loglik),1) #Test if VA in brightness is
significantly different from zero

# II) Hue
modelhue<-asreml(fixed=hue_n~ 1 +Vane+Sex+ Year
, random= ~ped(Ring) +at(machine):NestID
, rcov= ~ at(machine):units

```

```

      , data=Data.means
      ,ginverse=list(Ring=ainv)
      , na.method.X="include", na.method.Y="omit")

summary(modelhue)$varcomp
wald.asreml(modelhue,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelhue$coefficients$fixed#Coefficients of fixed effects

pin2(modelhue, h2~V3/(V1+V3+V4))#First h2
pin2(modelhue, h2~V3/(V2+V3+V5))#Second h2

pin2(modelhue, CE~V1/(V1+V3+V4))#First VCE/VP
pin2(modelhue, CE~V2/(V2+V3+V5))#Second VCE/VP

modelhue2<-asreml(fixed=hue_n~ 1 +Vane+Sex+ Year
      , random= ~at(machine):NestID
      , rcov= ~ at(machine):units
      , data=Data.means
      , na.method.X="include", na.method.Y="omit")

summary(modelhue2)
1-pchisq(2*(modelhue$loglik-modelhue2$loglik),1)#Test if VA in hue is significantly
different from zero

# III) UV chroma

modelUV<-asreml(fixed=UV.chrome_n~ 1 +Vane +Sex + Year
      , random= ~ped(Ring) +at(machine):NestID
      , rcov= ~ at(machine):units
      , data=Data.means
      ,ginverse=list(Ring=ainv)
      , na.method.X="include", na.method.Y="omit")

summary(modelUV)$varcomp

wald.asreml(modelUV,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelUV$coefficients$fixed#Coefficients of fixed effects

pin2(modelUV, h2~V3/(V1+V3+V4))#First h2
pin2(modelUV, h2~V3/(V2+V3+V5))#Second h2

pin2(modelUV, CE~V1/(V1+V3+V4))#First VCE/VP
pin2(modelUV, CE~V2/(V2+V3+V5))#Second VCE/VP

modelUV2<-asreml(fixed=UV.chrome_n~ 1 +Vane+Sex+ Year
      , random= ~at(machine):NestID
      , rcov= ~ at(machine):units
      , data=Data.means
      , na.method.X="include", na.method.Y="omit")

summary(modelUV2)
1-pchisq(2*(modelUV$loglik-modelUV2$loglik),1)#Test if VA in UV chroma is
significantly different from zero

# IV) PHA

modelPHA<-asreml(fixed=PHA~ 1+ Year+Sex
      , random= ~ped(Ring) +NestID
      , data=Data.means
      ,ginverse=list(Ring=ainv)
      , na.method.X="include", na.method.Y="omit")

summary(modelPHA)$varcomp
wald.asreml(modelPHA,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelPHA$coefficients$fixed#Coefficients of fixed effects

```

```

pin(modelPHA, h2~V2/(V1+V2+V3))# h2
pin(modelPHA, CE~V1/(V1+V2+V3))# VCE/VP

modelPHA2<-asreml(fixed=PHA~ 1+ Year+Sex
, random= ~NestID
, data=Data.means
, na.method.X="include", na.method.Y="omit")
summary(modelPHA2)$varcomp

1-pchisq(2*(modelPHA$loglik-modelPHA2$loglik),1)#Test if VA in PHA is significantly
different from zero

# V) Size-corrected weight

modelW<-asreml(fixed=Weight_d16~ 1+ Year+Sex + Tarsus
, random= ~ped(Ring) +NestID
, data=Data.means
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelW)$varcomp
wald.asreml(modelW,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelW$coefficients$fixed#Coefficients of fixed effects

pin(modelW, h2~V2/(V1+V2+V3))# h2
pin(modelW, CE~V1/(V1+V2+V3))# VCE/VP

modelW2<-asreml(fixed=Weight_d16~ 1+ Year+Sex + Tarsus
, random= ~NestID
, data=Data.means
, na.method.X="include", na.method.Y="omit")

summary(modelW2)$varcomp
1-pchisq(2*(modelW$loglik-modelW2$loglik),1)#Test if VA in weight is significantly
different from zero

# VI) Wing length
modelWing<-asreml(fixed=Wing~ 1+ Year+Sex
, random= ~ped(Ring) +NestID
, data=Data.means
,ginverse=list(Ring=ainv)
, na.method.X="include", na.method.Y="omit")

summary(modelWing)$varcomp
wald.asreml(modelWing,ssType = "conditional", denDF="numeric")#Test significance of
fixed effects
modelWing$coefficients$fixed#Coefficients of fixed effects

pin(modelWing, h2~V2/(V1+V2+V3))# h2
pin(modelWing, CE~V1/(V1+V2+V3))# VCE/VP

modelWing2<-asreml(fixed=Wing~ 1+ Year+Sex
, random= ~NestID
, data=Data.means
, na.method.X="include", na.method.Y="omit")

summary(modelWing2)$varcomp
1-pchisq(2*(modelWing$loglik-modelWing2$loglik),1)#Test if VA in Wing length is
significantly different from zero

#####
#Multivariate models
#####

###Multivariate model for all condition traits

```

```

multi.nestling.w<-asreml(cbind(PHA,Weight_d16,Wing)~ trait + trait:Year + trait:Sex
+ at(trait,2):Tarsus,

random=~us(trait,init=c(0.005,0.009,0.3,0.03,0.3,9)):ped(Ring)+
us(trait,init=c(0.007,0.3,0.47,0.08,0.9,3.6)):NestID,
      rcov=~units:diag(trait,init=c(0.01, 0.16, 0.7)),
      ginverse=list(Ring=ainv),
      data = Data.means,na.method.X="include",
      maxiter=100)
summary(multi.nestling.w)
pin(multi.nestling.w, cor~V8/(sqrt(V7*V9))#Weight-PHA
pin(multi.nestling.w, cor~V10/(sqrt(V7*V12))#Wing-PHA
pin(multi.nestling.w, cor~V11/(sqrt(V12*V9))#Weight-Wing

#Check if the correlations between performance traits are influenced by wing
thickness before injection
multi.nestling.w2<-asreml(cbind(PHA,Weight_d16,Wing,WingWeb_d13)~ trait +
trait:Year + trait:Sex + at(trait,2):Tarsus,
      random=~us(trait,init=c(0.006,0.02,0.37,0.06,0.86,5.27,-
0.03,0.1,0.1,0.57)):ped(Ring)+
us(trait,init=c(0.007,0.03,0.48,0.06,0.73,4.64,0.03,0.1,0.1,3.97)):NestID,
      rcov=~units:diag(trait,init=c(0.01,0.12,2.28,4.19)),
      ginverse=list(Ring=ainv),
      data = Data.means,na.method.X="include",
      maxiter=300)
summary(multi.nestling.w2)
pin(multi.nestling.w2, cor~V12/(sqrt(V11*V13))
pin(multi.nestling.w2, cor~V14/(sqrt(V11*V16))
pin(multi.nestling.w2, cor~V15/(sqrt(V13*V16))

#Test these correlations 1 by 1
sv<-asreml(cbind(PHA,Weight_d16,Wing)~ trait + trait:Year + trait:Sex +
at(trait,2):Tarsus,
      random=~us(trait):ped(Ring)+
us(trait,init=c(0.007,0.3,0.47,0.08,0.9,3.6)):NestID,
      rcov=~units:diag(trait,init=c(0.01, 0.16, 0.7)),
      ginverse=list(Ring=ainv),
      data = Data.means,na.method.X="include",
      maxiter=100,start.values = T)
gam<-sv$gammas.table
gam$Value[5]<-0#Values 2,4,5
gam$Constraint[5]<-"F"

multi.nestling.wb<-asreml(cbind(PHA,Weight_d16,Wing)~ trait + trait:Year +
trait:Sex + at(trait,2):Tarsus,
      random=~us(trait):ped(Ring)+
us(trait,init=c(0.007,0.3,0.47,0.08,0.9,3.6)):NestID,
      rcov=~units:diag(trait,init=c(0.01, 0.16, 0.7)),
      ginverse=list(Ring=ainv),
      data = Data.means,na.method.X="include",
      maxiter=100,G.param = gam)
summary(multi.nestling.wb)

1-pchisq(2*(multi.nestling.w$loglik-multi.nestling.wb$loglik),1)#Test for the
statistical significance of the genetic correlation

###Multivariate model for all coloration traits

multi.nestling.c<-asreml(cbind(brightness_n,UV.chrome_n,hue_n)~ trait + trait:Year
+ trait:Sex + trait:Vane,
      random=~ us(trait,init=c(0.90,-1.22e-3,1.62e-5,-0.25,-6e-
3,13.11)):at(machine,init=c(1,2)):NestID + us(trait,init=c(0.24,-5e-4,8e-6,-0.4,-
4e-3,8)):ped(Ring),
      rcov=~ units:us(trait,init=c(2,-1.44e-3,3.7e-5,-0.56,-5.8e-
3,40.5)):at(machine,init=c(1,2)),
      ginverse=list(Ring=ainv),
      data = Data.means,na.method.X="include",

```

```

maxiter=300)
summary(multi.nestling.c)$varcomp

#Test these correlations 1 by 1
sv<-asreml(cbind(brightness_n,UV.chrome_n,hue_n)~ trait + trait:Year + trait:Sex +
trait:Vane,
  random=~ us(trait,init=c(0.90,-1.22e-3,1.62e-5,-0.25,-6e-
3,13.11)):at(machine,init=c(1,2)):NestID + us(trait,init=c(0.24,-5e-4,8e-6,-0.4,-
4e-3,8)):ped(Ring) ,
  rcov=~ units:us(trait,init=c(2,-1.44e-3,3.7e-5,-0.56,-5.8e-
3,40.5)):at(machine,init=c(1,2)),
  ginverse=list(Ring=ainv),
  data = Data.means,na.method.X="include",
  maxiter=300,start.values = T)

gam<-sv$gammas.table
gam$Value[17]<-0#values 14,16,17
gam$Constraint[17]<-"F"

multi.nestling.cb<-asreml(cbind(brightness_n,UV.chrome_n,hue_n)~ trait + trait:Year
+ trait:Sex + trait:Vane,
  random=~ us(trait,init=c(0.90,-1.22e-3,1.62e-5,-0.25,-6e-
3,13.11)):at(machine,init=c(1,2)):NestID + us(trait):ped(Ring) ,
  rcov=~ units:us(trait,init=c(2,-1.44e-3,3.7e-5,-0.56,-5.8e-
3,40.5)):at(machine,init=c(1,2)),
  ginverse=list(Ring=ainv),
  data = Data.means,na.method.X="include",
  maxiter=300,G.param=gam)

summary(multi.nestling.cb)
1-pchisq(2*(multi.nestling.c$loglik-multi.nestling.cb$loglik),1)#Test for the
statistical significance of the genetic correlation

###Multivariate model for all six traits

#Need to specify starting values
sv<-asreml(cbind(brightness_n,hue_n,UV.chrome_n,PHA, Weight_d16,Wing)~ trait +
trait:Year + trait:Sex + at(trait,1):Vane+ at(trait,2):Vane+ at(trait,3):Vane+
at(trait,5):Tarsus,
  random=~ us(trait):ped(Ring) + us(trait):at(machine):NestID ,
  rcov=~ units:us(trait):at(machine),
  ginverse=list(Ring=ainv),
  data = Data.means,na.method.X="include",
  maxiter=300,start.values = TRUE)
gam<-sv$gammas.table

gam[1:21,1:2]#pedigree
gam$Value[1:21]<-c(2.77E-01, -7.98E-01, 4.45E+00, -1.60E-04, -1.63E-03,
5.19E-06, -1.85E-03,
-5.69E-03, -8.29E-05, 5.32E-03, 5.31E-02, 1.28E-01,
-1.86E-04, 8.15E-03,
2.30E-01, 3.44E-01, -1.42E+00, -1.01E-03, 3.04E-02,
1.65E-01, 1.73E+00)

gam[22:42,1:2]#NestID1
gam$Value[c(22:42)]<-c(1.04E+00, -4.21E-01, 1.29E+01, -1.50E-03, -4.17E-03,
1.74E-05,4.19E-02,
-9.47E-03, 7.64E-05, 1.12E-02, 3.40E-01, 3.06E-01,
9.01E-04,5.25E-02,
5.96E-01, 1.50E+00, 1.21E+00, 1.80E-04, 1.41E-01,
1.18E+00, 7.92E+00)

gam[43:63,1:2]#NestID2
gam$Value[c(43:63)]<-c(1.95E+00, -1.52E+00, 2.51E+01, -1.16E-02, -1.74E-02,
1.44E-04, 1.35E-02,
3.47E-03, -1.01E-04, 3.45E-03, 3.22E-01, -8.74E-02,
-2.20E-03, 1.89E-02,

```

```

        4.61E-01, 1.51E+00, -2.00E+00, -7.99E-03, 3.69E-03,
        6.67E-01, 4.18E+00)
gam[65:85,1:2]#residual1
gam$Value[c(65:85)]<-c(2.02E+00, -1.70E-02, 4.53E+01, -1.75E-03, -1.01E-02,
4.08E-05, -3.98E-03,
2.56E-02, 1.02E-04, 1.44E-02, 1.11E-02, -1.93E-01,
6.53E-04, 1.51E-02,
2.73E-01, 4.08E-01, 1.53E+00, 3.81E-03, 4.60E-02,
7.21E-01, 7.77E+00)
gam[87:107,1:2]#residual2
gam$Value[c(87:107)]<-c(4.68E+00, 8.19E-01, 8.49E+01, -2.98E-02, -8.10E-02
,4.61E-04 ,8.83E-03,
-5.42E-02, 7.22E-05, 1.14E-02, 5.60E-02, -
5.04E-01, -3.15E-04, 5.99E-03,
1.88E-01, 9.52E-01, -1.54E+00, -7.07E-03, 7.48E-03,
3.31E-01, 3.34E+00)

#Run the model
multi.total.nestling<-asreml(cbind(brightness_n,hue_n,UV.chrome_n,PHA,
Weight_d16,Wing)~ trait + trait:Year + trait:Sex + at(trait,1):Vane+
at(trait,2):Vane+ at(trait,3):Vane+ at(trait,5):Tarsus,
random=~ us(trait):ped(Ring) +
us(trait):at(machine):NestID ,
rcov=~ units:us(trait):at(machine),
ginverse=list(Ring=ainv),
data = Data.means,na.method.X="include",G.param=gam,
R.param=gam,
maxiter=100)

summary(multi.total.nestling)$varcomp

#Calculate genetic correlations and their SE using the pin function

pin(multi.total.nestling,rg~ V44/sqrt(V43*V45))#hue-bri
pin(multi.total.nestling,rg~ V46/sqrt(V43*V48))#UV-bri
pin(multi.total.nestling,rg~ V47/sqrt(V45*V48))#UV-hue

pin(multi.total.nestling,rg~ V49/sqrt(V43*V52))#PHA-bri
pin(multi.total.nestling,rg~ V50/sqrt(V45*V52))#PHA-hue
pin(multi.total.nestling,rg~ V51/sqrt(V48*V52))#PHA-UV

pin(multi.total.nestling,rg~ V53/sqrt(V43*V57))#Weight- Bri
pin(multi.total.nestling,rg~ V54/sqrt(V45*V57))#Weight- Hue
pin(multi.total.nestling,rg~ V55/sqrt(V48*V57))#Weight- UV
pin(multi.total.nestling,rg~ V56/sqrt(V57*V52))#Weight- PHA

pin(multi.total.nestling,rg~ V58/sqrt(V43*V63))#Wing -Bri
pin(multi.total.nestling,rg~ V59/sqrt(V45*V63))#Wing -Hue
pin(multi.total.nestling,rg~ V60/sqrt(V48*V63))#Wing- UV
pin(multi.total.nestling,rg~ V61/sqrt(V52*V63))#Wing -PHA
pin(multi.total.nestling,rg~ V62/sqrt(V57*V63))#Wing -Weight

###Multivariate model on the phenotypic level

multi.pheno.nestling<-asreml(cbind(brightness_n,hue_n,UV.chrome_n,PHA,
Weight_d16,Wing)~ trait + trait:Year + trait:Sex + at(trait,1):Vane+
at(trait,2):Vane+ at(trait,3):Vane+ at(trait,5):Tarsus,
rcov=~ units:us(trait),
data = Data.means,na.method.X="include",maxiter=100)
summary(multi.pheno.nestling)$varcomp

#Calculate phenotypic correlations and their SE using the pin function

pin(multi.pheno.nestling,rg~ V3/sqrt(V2*V4))#hue-bri
pin(multi.pheno.nestling,rg~ V5/sqrt(V2*V7))#UV-bri
pin(multi.pheno.nestling,rg~ V6/sqrt(V4*V7))#UV-hue

```

```

pin(multi.pheno.nestling,rg~ V8/sqrt(V2*V11))#PHA-bri
pin(multi.pheno.nestling,rg~ V9/sqrt(V4*V11))#PHA-hue
pin(multi.pheno.nestling,rg~ V10/sqrt(V7*V11))#PHA-UV

pin(multi.pheno.nestling,rg~ V12/sqrt(V2*V16))#Weight- Bri
pin(multi.pheno.nestling,rg~ V13/sqrt(V4*V16))#Weight- Hue
pin(multi.pheno.nestling,rg~ V14/sqrt(V7*V16))#Weight- UV
pin(multi.pheno.nestling,rg~ V15/sqrt(V11*V16))#Weight- PHA

pin(multi.pheno.nestling,rg~ V17/sqrt(V2*V22))#Wing -Bri
pin(multi.pheno.nestling,rg~ V18/sqrt(V4*V22))#Wing -Hue
pin(multi.pheno.nestling,rg~ V19/sqrt(V7*V22))#Wing- UV
pin(multi.pheno.nestling,rg~ V20/sqrt(V11*V22))#Wing -PHA
pin(multi.pheno.nestling,rg~ V21/sqrt(V16*V22))#Wing -Weight

#####
#A different approach to calculate heritability of PHA
#Excludes measurement error
#####
WingWebs<-read.table("WingWebs.txt",header=T)
WingWebs$Ring<-as.factor(WingWebs$Ring)
WingWebs$NestID<-as.factor(WingWebs$NestID)
WingWebs$Year<-as.factor(WingWebs$Year)
WingWebs$Sex<-as.factor(WingWebs$Sex)

#Run a multivariate model with repeated measures of wing web before and after
injection
multi<-asreml(cbind(WingWeb_d13,WingWeb_d14)~ trait + trait:Year + trait:Sex ,
              random=~us(trait):ped(Ring)+ + us(trait):ide(Ring)+us(trait):NestID,
              rcov=~units:diag(trait),
              ginverse=list(Ring=ainv),
              data = WingWebs,na.method.X="include",
              maxiter=1000)
summary(multi)$varcomp

#Calculate heritability of the difference=PHA
pin(multi, h2~(V4+V6-2*V5) / ((V4+V6-2*V5)+(V1+V3-2*V2)+(V7+V9-2*V8)))#heritability
pin(multi, CE2~(V1+V3-2*V2) / ((V4+V6-2*V5)+(V1+V3-2*V2)+(V7+V9-2*V8)))#VCE/VP

#####
#Estimate cross-sex genetic correlations
#####

#Create the table for analyses
Data.means.fem<-Data.means[Data.means$Sex=="f",]
Data.means.mal<-Data.means[Data.means$Sex=="m",]

names(Data.means.fem)[5:10]<-paste(names(Data.means.fem)[5:10],"_f",sep="")
names(Data.means.mal)[5:10]<-paste(names(Data.means.mal)[5:10],"_m",sep="")

Data.means.mal$hue_n_f<-NA
Data.means.mal$brightness_n_f<-NA
Data.means.mal$UV.chrome_n_f<-NA
Data.means.mal$PHA_f<-NA
Data.means.mal$Weight_d16_f<-NA
Data.means.mal$Wing_f<-NA

Data.means.fem$hue_n_m<-NA
Data.means.fem$brightness_n_m<-NA
Data.means.fem$UV.chrome_n_m<-NA
Data.means.fem$PHA_m<-NA
Data.means.fem$Weight_d16_m<-NA
Data.means.fem$Wing_m<-NA

Data.means.fem<-Data.means.fem[,names(Data.means.mal)]
names(Data.means.fem)

```

```

Data.means.sexes<-rbind(Data.means.mal,Data.means.fem)
Data.means.sexes<-Data.means.sexes[order(Data.means.sexes$machine),]

#Run the bivariate model for each trait
biv.sexes.bri<-asreml(cbind(brightness_n_m,brightness_n_f)~ trait + trait:Year +
trait:Vane,
                    random=~us(trait):ped(Ring)+ us(trait):NestID:at(machine)
                    ,
                    rcov=~units:diag(trait):at(machine),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=500)
summary(biv.sexes.bri)
pin(biv.sexes.bri, cor~V8/(sqrt(V7*V9)))#r=0.99 se=0.53

biv.sexes.hue<-asreml(cbind(hue_n_m,hue_n_f)~ trait + trait:Year + trait:Vane,
                    random=~us(trait):ped(Ring)+ us(trait):NestID:at(machine) ,
                    rcov=~units:diag(trait):at(machine),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=500)
summary(biv.sexes.hue)
pin(biv.sexes.hue, cor~V8/(sqrt(V7*V9)))#r=0.59 se=0.48

biv.sexes.UV<-asreml(cbind(UV.chrome_n_f,UV.chrome_n_m)~ trait + trait:Year +
trait:Vane,
                    random=~us(trait):ped(Ring)+ us(trait):NestID:at(machine) ,
                    rcov=~units:diag(trait):at(machine),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=500)
biv.sexes.UV<-update(biv.sexes.UV)
summary(biv.sexes.UV)

pin(biv.sexes.UV, cor~V8/(sqrt(V7*V9)))#r=0.95 se=0.52

biv.sexes.PHA<-asreml(cbind(PHA_f,PHA_m)~ trait + trait:Year,
                    random=~us(trait):ped(Ring)+ us(trait):NestID ,
                    rcov=~units:diag(trait),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=400)
summary(biv.sexes.PHA)

pin(biv.sexes.PHA, cor~V5/(sqrt(V4*V6)))#r=0.98 se=0.17

biv.sexes.W<-asreml(cbind(Weight_d16_f,Weight_d16_m)~ trait + trait:Year +
trait:Tarsus,
                    random=~us(trait):ped(Ring)+ us(trait):NestID ,
                    rcov=~units:diag(trait),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=300)
summary(biv.sexes.W)

pin(biv.sexes.W, cor~V5/(sqrt(V4*V6)))#r=0.92 se=0.05

biv.sexes.Wing<-asreml(cbind(Wing_f,Wing_m)~ trait + trait:Year,
                    random=~us(trait):ped(Ring)+ us(trait):NestID ,
                    rcov=~units:diag(trait),
                    ginverse=list(Ring=ainv),
                    data = Data.means.sexes,na.method.X="include",
                    maxiter=300)
summary(biv.sexes.Wing)

pin(biv.sexes.Wing, cor~V5/(sqrt(V4*V6)))#r=0.94 se=0.08

```

*Text S2: R code for performing structural equation modelling on the G matrix*

```

library(lavaan)

#Extract additive genetic (co)variances estimates from the full model
Gmat<-summary(multi.total.nestling)$varcomp[43:63,]

#Create covariance matrix based on these estimates
cov.color_n_gen <- matrix(Gmat[c(1,2,4,7,11,16,
                                2,3,5,8,12,17,
                                4,5,6,9,13,18,
                                7,8,9,10,14,19,
                                11,12,13,14,15,20,
                                16,17,18,19,20,21),1],6,6,byrow=TRUE)

rownames(cov.color_n_gen) <- colnames(cov.color_n_gen) <- c("BriN","Hue","UV",
"PHA","Weight","Wing")

#Transform it into a correlation martix
cor.color_n_gen <- cov2cor(cov.color_n_gen)

#Model where the three performance traits load on a single "performance" factor
HS.modelW <-
  'L1 =~lam1*Weight+ lam2*Wing + lam3*PHA
Weight~~b1*Weight
Wing~~b2*Wing
PHA~~b3*PHA
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2)
'

fitW <- cfa(HS.modelW, sample.cov =cor.color_n_gen,sample.nobs=306,std.lv=TRUE)
standardizedsolution(fitW)

#First model: estimate the correlation between brightness and performance
HS.modelA <-
  'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*BriN
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
BriN~~0*BriN
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2) '
fitA <- cfa(HS.modelA, sample.cov =cor.color_n_gen,sample.nobs=306,std.lv=TRUE)
standardizedsolution(fitA)

#Second model: estimate the correlation between hue and performance

HS.modelB <-
  'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*Hue
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
Hue~~0*Hue
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2) '
fitB <- sem(HS.modelB, sample.cov =cor.color_n_gen,sample.nobs=306,std.lv=TRUE)
standardizedsolution(fitB)

#Third model: estimate the correlation between UV chroma and performance
HS.modelC <-

```

```
'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*UV
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
UV~~0*UV
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2) '
fitC <- cfa(HS.modelC, sample.cov =cor.color_n_gen, sample.nobs=306, std.lv=TRUE)
standardizedsolution(fitC)
```

*Text S3: R code for performing simulations*

```

library(lavaan)
library(asreml)
library(pedantics)
#Open data files
pedigree<-read.table("Pedigree_BT.txt",header=T)
ainv<-asreml.Ainverse(pedigree)$ginv

data.chicks<-read.table("Data_nestlings.txt",header=T)
data.chicks<-unique(data.chicks[,c(1,3)])
NestIDS<-unique(data.chicks$NestID)
names(data.chicks)[1]<-"id"
#Extract additive genetic (co)variances estimates from the full model
Gmat<-summary(multi.total.nestling)$varcomp[43:63,1]
Emat1<-summary(multi.total.nestling)$varcomp[65:85,1]
CEmat1<-summary(multi.total.nestling)$varcomp[1:21,1]

#Create covariance matrices based on these estimates
SigmaG <- matrix(Gmat[c(1,2,4,7,11,16,
                        2,3,5,8,12,17,
                        4,5,6,9,13,18,
                        7,8,9,10,14,19,
                        11,12,13,14,15,20,
                        16,17,18,19,20,21)],6,6,byrow=TRUE)

SigmaE <- matrix(Emat1[c(1,2,4,7,11,16,
                        2,3,5,8,12,17,
                        4,5,6,9,13,18,
                        7,8,9,10,14,19,
                        11,12,13,14,15,20,
                        16,17,18,19,20,21)],6,6,byrow=TRUE)

SigmaCE <- matrix(CEmat1[c(1,2,4,7,11,16,
                        2,3,5,8,12,17,
                        4,5,6,9,13,18,
                        7,8,9,10,14,19,
                        11,12,13,14,15,20,
                        16,17,18,19,20,21)],6,6,byrow=TRUE)

#Run simulations
Results.sim<-data.frame("iteration"=NA,
"Conv1"=NA,"Conv2"=NA,"Conv3"=NA,"Bri.Perf"=NA, "Hue.Perf"=NA, "UV.Perf"=NA,
                        "lam1a"=NA, "lam2a"=NA,"lam3a"=NA,"lam1b"=NA,
"lam2b"=NA,"lam3b"=NA,"lam1c"=NA, "lam2c"=NA,"lam3c"=NA)

for ( i in 1:1000){

  Results.sim[i,1]<-i
#Simulate data using pedantics
simPhen<-phensim(pedigree,traits=6, randomA=SigmaG,randomE=SigmaE)$phenotypes

#Add CE effects
simPhen$id<-as.factor(simPhen$id)
CEeffects<-as.data.frame(rmvnorm(length(NestIDS),sigma=SigmaCE))
CEeffects$NestID<-as.factor(NestIDS)
simPhen<-merge(simPhen,data.chicks)
simPhen<-merge(simPhen,CEeffects)
Datasim<-simPhen[,c(1,2)]
Datasim$trait_1<-simPhen$trait_1+simPhen$V1
Datasim$trait_2<-simPhen$trait_2+simPhen$V2
Datasim$trait_3<-simPhen$trait_3+simPhen$V3
Datasim$trait_4<-simPhen$trait_4+simPhen$V4
Datasim$trait_5<-simPhen$trait_5+simPhen$V5
Datasim$trait_6<-simPhen$trait_6+simPhen$V6
Datasim$NestID<-as.factor(Datasim$NestID)
###Analyse using asreml

```

```

#Run the models: need to run 3 different models because 6-trait model never
converges despite starting values
sv1<-
c(SigmaG[1,1],SigmaG[1,4],SigmaG[4,4],SigmaG[1,5],SigmaG[4,5],SigmaG[5,5],SigmaG[1,
6],SigmaG[4,6],SigmaG[5,6],SigmaG[6,6])
sv2<-
c(SigmaG[2,2],SigmaG[2,4],SigmaG[4,4],SigmaG[2,5],SigmaG[4,5],SigmaG[5,5],SigmaG[2,
6],SigmaG[4,6],SigmaG[5,6],SigmaG[6,6])
sv3<-
c(SigmaG[3,3],SigmaG[3,4],SigmaG[4,4],SigmaG[3,5],SigmaG[4,5],SigmaG[5,5],SigmaG[3,
6],SigmaG[4,6],SigmaG[5,6],SigmaG[6,6])

sv1b<-
c(SigmaE[1,1],SigmaE[1,4],SigmaE[4,4],SigmaE[1,5],SigmaE[4,5],SigmaE[5,5],SigmaE[1,
6],SigmaE[4,6],SigmaE[5,6],SigmaE[6,6])
sv2b<-
c(SigmaE[2,2],SigmaE[2,4],SigmaE[4,4],SigmaE[2,5],SigmaE[4,5],SigmaE[5,5],SigmaE[2,
6],SigmaE[4,6],SigmaE[5,6],SigmaE[6,6])
sv3b<-
c(SigmaE[3,3],SigmaE[3,4],SigmaE[4,4],SigmaE[3,5],SigmaE[4,5],SigmaE[5,5],SigmaE[3,
6],SigmaE[4,6],SigmaE[5,6],SigmaE[6,6])

sv1c<-
c(SigmaCE[1,1],SigmaCE[1,4],SigmaCE[4,4],SigmaCE[1,5],SigmaCE[4,5],SigmaCE[5,5],Sig
maCE[1,6],SigmaCE[4,6],SigmaCE[5,6],SigmaCE[6,6])
sv2c<-
c(SigmaCE[2,2],SigmaCE[2,4],SigmaCE[4,4],SigmaCE[2,5],SigmaCE[4,5],SigmaCE[5,5],Sig
maCE[2,6],SigmaCE[4,6],SigmaCE[5,6],SigmaCE[6,6])
sv3c<-
c(SigmaCE[3,3],SigmaCE[3,4],SigmaCE[4,4],SigmaCE[3,5],SigmaCE[4,5],SigmaCE[5,5],Sig
maCE[3,6],SigmaCE[4,6],SigmaCE[5,6],SigmaCE[6,6])

Test.multi1<-asreml(cbind(trait_1,trait_4, trait_5,trait_6)~ trait,
                    random=~ us(trait,init=sv1):ped(id)+ us(trait,init=sv1c):NestID
                    ,
                    rcov=~ units:us(trait,init=sv1b),
                    ginverse=list(id=ainv),
                    data = Datasim,na.method.X="include",
                    maxiter=300)

Test.multi2<-asreml(cbind(trait_2,trait_4, trait_5,trait_6)~ trait,
                    random=~ us(trait,init=sv2):ped(id)+ us(trait,init=sv2c):NestID
                    ,
                    rcov=~ units:us(trait,init=sv2b),
                    ginverse=list(id=ainv),
                    data = Datasim,na.method.X="include",
                    maxiter=300)

Test.multi3<-asreml(cbind(trait_3,trait_4, trait_5,trait_6)~ trait,
                    random=~ us(trait,init=sv3):ped(id)+ us(trait,init=sv3c):NestID
                    ,
                    rcov=~ units:us(trait,init=sv3b),
                    ginverse=list(id=ainv),
                    data = Datasim,na.method.X="include",
                    maxiter=300)

Results.sim[i,2]<-Test.multi1$converge
Results.sim[i,3]<-Test.multi2$converge
Results.sim[i,4]<-Test.multi3$converge

Gmatsim1<-summary(Test.multi1)$varcomp[11:20,1]
Gmatsim2<-summary(Test.multi2)$varcomp[11:20,1]
Gmatsim3<-summary(Test.multi3)$varcomp[11:20,1]

#Extract covariance matrix
cov1<- matrix(Gmatsim1[c(1,2,4,7,

```

```

                2,3,5,8,
                4,5,6,9,
                7,8,9,10)],4,4,byrow=TRUE)

rownames(cov1) <- colnames(cov1) <- c("BriN","PHA","Weight","Wing")

cov2<- matrix(Gmatsim2[c(1,2,4,7,
                        2,3,5,8,
                        4,5,6,9,
                        7,8,9,10)],4,4,byrow=TRUE)

rownames(cov2) <- colnames(cov2) <- c("Hue","PHA","Weight","Wing")

cov3<- matrix(Gmatsim3[c(1,2,4,7,
                        2,3,5,8,
                        4,5,6,9,
                        7,8,9,10)],4,4,byrow=TRUE)

rownames(cov3) <- colnames(cov3) <- c("UV","PHA","Weight","Wing")

#Transform it into a correlation matrix
cor1 <- cov2cor(cov1)
cor2 <- cov2cor(cov2)
cor3 <- cov2cor(cov3)

###Run SEMS on this correlation matrix
#Test Brightness and performance
HS.modelA <-
  'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*BriN
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
BriN~~0*BriN
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2) '

fitA <- cfa(HS.modelA, sample.cov =cor1,sample.nobs=306,std.lv=TRUE)
solA<-data.frame(standardizedsolution(fitA))
Results.sim[i,"Bri.Perf"]<-solA[11,4]

#Test Hue and performance
HS.modelB <-
  'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*Hue
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
Hue~~0*Hue
b1==1-(lam1^2)
b2==1-(lam2^2)
b3==1-(lam3^2) '

fitB <- sem(HS.modelB, sample.cov =cor2,sample.nobs=306,std.lv=TRUE)
solB<-data.frame(standardizedsolution(fitB))
Results.sim[i,"Hue.Perf"]<-solB[11,4]

#Test UV and performance
HS.modelC <-
  'L1 =~lam1*PHA +lam2*Weight+ lam3*Wing
L2=~1*UV
PHA~~b1*PHA
Weight~~b2*Weight
Wing~~b3*Wing
UV~~0*UV
b1==1-(lam1^2)

```

```

b2==1-(lam2^2)
b3==1-(lam3^2) '

fitC <- cfa(HS.modelC, sample.cov =cor3,sample.nobs=306,std.lv=TRUE)
solC<-data.frame(standardizedsolution(fitC))
Results.sim[i,"UV.Perf"]<-solC[11,4]

Results.sim[i,"lam1a"]<-solA[1,4]
  Results.sim[i,"lam2a"]<-solA[2,4]
  Results.sim[i,"lam3a"]<-solA[3,4]

  Results.sim[i,"lam1b"]<-solB[1,4]
  Results.sim[i,"lam2b"]<-solB[2,4]
  Results.sim[i,"lam3b"]<-solB[3,4]

  Results.sim[i,"lam1c"]<-solC[1,4]
  Results.sim[i,"lam2c"]<-solC[2,4]
  Results.sim[i,"lam3c"]<-solC[3,4]

}###End of simulations

#Save results
write.table(Results.sim,"Result.sim.txt",row.names = F)

#Open results
Results.sim<-read.table("Result.sim.txt",header=T)

#Keep only the models where the 3 performance traits correlate positively
reduced1<-Results.sim[Results.sim$lam1a>0.05
&Results.sim$lam2a>0.05&Results.sim$lam3a>0.05&Results.sim$Conv1==TRUE,c(1,5,8:10)
]
reduced2<-Results.sim[Results.sim$lam1b>0.05
&Results.sim$lam2b>0.05&Results.sim$lam3b>0.05&Results.sim$Conv2==TRUE,c(1,6,11:13)
]
reduced3<-Results.sim[Results.sim$lam1c>0.05
&Results.sim$lam2c>0.05&Results.sim$lam3c>0.05&Results.sim$Conv3==TRUE,
c(1,7,14:16)]

hist(reduced1$Bri.Perf,xlim=c(-1,1),breaks=20)
hist(reduced2$Hue.Perf,xlim=c(-2,1),breaks=20)
hist(reduced3$UV.Perf,xlim=c(-2,1),breaks=20)

#Median and 95% CI of correlation estimates
median(reduced1$Bri.Perf)
median(reduced2$Hue.Perf)
median(reduced3$UV.Perf)

quantile(reduced1$Bri.Perf,probs=c(0.025,0.975))
quantile(reduced2$Hue.Perf,probs=c(0.025,0.975))
quantile(reduced3$UV.Perf,probs=c(0.025,0.975))

```

Dear Editor of Journal of Evolutionary Biology,

Thank you for your quick reply. We made the corrections that you requested in our manuscript "Tail color signals performance in blue tit nestlings" (JEB-2018-00555.R3).

Sincerely,

The author

-----

JEB ms # JEB-2018-00555.R2

Dear Dr. Class,

I am willing to consider acceptance of your paper for publication in JEB, provided you revise it along the lines recommended.

- Remove double brackets  $)), ), (($  throughout (e.g. l 244)
- state what you're reporting (line 250 - what's in the brackets? Report consistently - if that's se, then it's inconsistently reported - see line 256).
- report 95%CI as range (with a dash between, not a comma), throughout
- use space consistently in reporting results throughout (before and after numbers, =, 95%CI, 95 % CI ect.)

Please pay careful attention to the formatting of tables, figures and references, as well as the style used for reporting the results of statistical tests (see Instructions for Authors, <http://onlinelibrary.wiley.com/journal/10.1111/%28ISSN%291420-9101/homepage/ForAuthors.html>)

Please make sure any in-line statistics conform to the Instructions for authors ("In-line statistical results should be presented as Test-statistics: degrees of freedom as subscript(s) to test-statistics (e.g.  $F_{1,12} = \dots$  or  $t_8 = \dots$ ), followed by P-value., e.g. ( $F_{1,12} = 4.931$ ,  $P = 0.0464$ ). Statistical results in tables should be comprehensive, allowing future meta-analyses. Depending on the details of the analyses, results reported may include parameter estimates, test-statistics, degrees of freedom, significance levels and err/residual model information (e.g. error MS's and df's in ANOVA or regression models). Since exact P-values can be useful for meta-analyses, we recommend that these are quoted even when non-significant, e.g.  $t_{23}=0.25$ ,  $P=0.34$ , or  $F_{2,32}=1.12$ ,  $P=0.55$ . However, non significant tests (i.e.  $P > 0.05$ ) should always be interpreted as such.")

Please submit your revised paper in an editable format, within 30 days from the date of this letter, since otherwise it will be considered as a newly submitted manuscript. You can access the revision submission by clicking on the link Manuscripts with Decisions in your Author Center.

Sincerely,

Julia Schroeder