

<https://helda.helsinki.fi>

Quantification of folate in the main steps of traditional processing of tef injera, a cereal based fermented staple food

Tamene, Aynadis

2019-05

Tamene , A , Kariluoto , S , Baye , K & Humblot , C 2019 , ' Quantification of folate in the main steps of traditional processing of tef injera, a cereal based fermented staple food ' , Journal of Cereal Science , vol. 87 , pp. 225-230 . <https://doi.org/10.1016/j.jcs.2019.04.005>

<http://hdl.handle.net/10138/327305>

<https://doi.org/10.1016/j.jcs.2019.04.005>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Quantification of folate in the main steps of traditional processing of tef *injera*, a cereal based**
2 **fermented staple food**

3 Aynadis Tamene^a, Susanna Kariluoto^b, Kaleab Baye^{a*} and Christèle Humblot^c

4 ^aCenter for Food Science and Nutrition, College of Natural and Computational Sciences; Addis Ababa
5 University, P.O. Box 150201, Addis Ababa, Ethiopia. aynadis.tamene@aau.edu.et and
6 kaleabbaye@gmail.com

7 ^bDepartment of Food and Environmental Sciences, University of Helsinki, P.O. Box 66, FIN-00014,
8 Finland. susanna.kariluoto@helsinki.fi

9 ^cUMR Nutripass, IRD, University of Montpellier / Montpellier SupAgro, Montpellier, France.

10 christele.humblot@ird.fr

11 *correspondence should be addressed to: kaleabbaye@gmail.com

12

13

14

15 **Highlights**

16 Tef flour had an average folate content of 59 µg/100 g of dry matter content.

17 Fermentation increased batter folate content up to 148% compared with that of flour.

18 Thermal treatment (baking) always reduced folate content.

19

20

21 **Abstract**

22 *Injera is an Ethiopian fermented flatbread preferably made from whole grain cereal (teff). Tef it is*
23 *increasingly used to produce gluten-free pasta and bread, but the folate content of teff and*
24 *products made from it remains unknown. Given that folate deficiencies lead to several health*
25 *disorders, the aim of this study was to quantify folate in each of the three main steps of traditional*
26 *processing of teff injera. Total folate contents of teff flour, fermented batter and injera were*
27 *determined through microbiological assays using Lactobacillus rhamnosus (ATCC 7469). Folate*
28 *content of teff flour was 8.7 µg/100 g of dry matter content, which is in the same range as the richest*
29 *cereals like oats. The increase in folate content due to fermentation was highly variable (60-*
30 *148%). Cooking always led to folate losses, with a maximum of 52.8%. Altogether, injera*
31 *processing increased folate retention between 38.0 and 121.8%. Folate content of injera was 14.3*
32 *µg/100 g on fresh weight-basis. Tef Injera can contribute up to 10% of the recommended nutrient*
33 *intake of folate for children aged 1–3 and women of reproductive age. Although the folate content*
34 *of teff is already high, future studies should focus on optimizing the folate content of injera.*

35 **Key words:** Fermentation, Folate, *Injera*, Tef.

36

37

38

39

40 **1. Introduction**

41 Every year, inadequate folate intake predisposes women to birth complications like neural tube
42 defects (Moore *et al.*, 2003). Inadequate maternal folate status has also been associated with low
43 infant birth weight, preterm delivery and fetal growth retardation (Scholl and Johnson, 2000).
44 Megaloblastic anemia and elevated blood concentrations of homocysteine have also been linked
45 to folate deficiencies (Bailey and Gregory, 2006) and (Ho *et al.*, 2011). Although animal source
46 foods (liver, kidney, chicken giblets, egg yolk), and vegetable source foods (pulses, and dark-green
47 leafy vegetables) are rich sources of folate, regular consumption of some of these foods is limited
48 in low and middle income countries (LMIC). Instead, diets are predominantly based on cereal and
49 pulses (Lee *et al.*, 2013). Despite cereal and pulses non-negligible folate content, combined with
50 the limited availability and access to folic acid fortified foods, and the low compliance/adherence
51 to folic acid supplementation during pregnancy, a significant proportion of the population in LMIC
52 is at risk of folate deficiency and its adverse effects (McLean *et al.*, 2008; Haidar, 2010).

53

54 In Africa, the preparation of many cereal-based staple foods includes a fermentation step (Humblot
55 and Guyot, 2008). *Injera* is a staple food that is widely consumed in Ethiopia (Baye *et al.*, 2013),
56 and is often prepared from tef (*Eragrostis tef*), an ancient cereal crop indigenous to Ethiopia
57 (Yetneberk *et al.*, 2004). Tef is becoming popular worldwide thanks to its nutritional profile
58 (gluten-free, high dietary fibre content, high iron content etc.). It is increasingly used in health
59 food to produce gluten-free pasta and bread (Zhu, 2018). But to the best of our knowledge, its
60 folate content has never been estimated. Fermentation and thermal treatment - baking - the two
61 main processes used to prepare injera may have an effect on the folate content of *injera*. For
62 example, baking has been found to cause up to 25% folate losses in wheat and rye sour batter

63 breads (Kariluoto *et al.*, 2004 and Gujska and Majewska, 2005). Unlike heat treatment, household
64 fermentation can either increase or decrease the initial folate content of the flour (Saubade *et al.*,
65 2017a). During fermentation of food products, yeasts and some bacteria have been found to
66 increase the folate content of the original raw material. But the folate originally present in the
67 cereal or that resulting from microbial synthesis can also be consumed by other bacteria (LeBlanc
68 *et al.*, 2007; Moslehi-Jenabian *et al.*, 2010; Holzapfel, 2002 Keagy *et al.*, 1975 and Kariluoto *et*
69 *al.*, 2004). Thus, the final folate content of the fermented food is a balance between production
70 and consumption by microorganisms.

71
72 Information on folates in tef *injera* is rare, and the effect of fermentation and baking in households
73 is largely unknown. This is unfortunate because daily intake and hence the risk of folate deficiency
74 cannot be estimated without precise knowledge of the folate content of widely consumed foods
75 like *injera*. A better understanding of the dynamics
76 of folate retention after tef fermentation and of the extent of folate losses caused by thermal
77 treatment when *injera* is prepared in the household should help optimize the preparation of *injera*
78 by increasing the production and retention of folate.

79
80 The aim of this study was to quantify folate in the main steps of traditional processing of tef *injera*.
81 The folate content of tef flour was quantified to complete the food composition table in Ethiopia,
82 which currently does not include folate. The traditional preparation of tef *injera* in urban
83 households was characterized in detail and the effect of fermentation and thermal treatment on
84 folate content evaluated. Finally, the contribution of tef *injera* to meeting folate requirements was
85 estimated.

86 **2. Material and methods**

87 **2.1. Chemicals**

88 All the chemicals used in this study were purchased from Sigma-Aldrich Chemie GmbH,
89 Switzerland.

90

91 **2.2. Sampling of tef flour, batter and injera**

92 Detailed observations of the traditional *injera* making process were made in 20 selected
93 households (where *injera* is traditionally prepared) in Addis Ababa. Since there are 10 sub-cities
94 in Addis Ababa, two households were selected for observation in each sub-city, which resulted the
95 flow diagram presented in Fig. 1. Briefly, the process begins by milling whole tef grain into flour,
96 mixing 4–5 kg of tef flour and with 5–6 L of tap water. Fermentation is started by inoculation by
97 backslopping using 1 L of leftover (called *ersho*) from a previous successful spontaneous
98 traditional fermentation. The mixture is then allowed to ferment for an average of 3–4 days at room
99 temperature (called 1st stage fermentation). After fermentation, the liquid present on top of the
100 batter (the supernatant) is discarded and replaced with the same volume of fresh tap water. Then,
101 a portion equivalent to 1/11th of the fermented batter (1 L) is mixed with 3 L of tap water, boiled
102 for 10 min and allowed to cool to ~45 °C. The resulting product is called *absit*, it serves as a batter
103 binder and is added back to the fermented batter, which is allowed to ferment for an additional 2-
104 3 h (2nd stage fermentation). Finally, 450 mL of the fermented rather liquid batter is poured onto
105 a hot clay griddle, covered, and baked for 1–2 min. The resulting flat bread is called *injera*.

106 Samples of tef flour (n=60), batter (n=60), and *injera* (n=60) were collected from the 20
107 households on three separate occasions (referred to as sampling occasion 1, 2 and 3) at intervals
108 of approximately one month, giving a total of 180 samples for the experiment. The samples were
109 collected from each household using a simple random sampling technique. The samples were
110 collected aseptically and placed in sterile plastic bottles covered with aluminum foil to protect
111 them from direct light and transported back to the laboratory in an ice box. The dry matter (DM)
112 content of all three types of samples (flour, batter and *injera*) and the pH of the batter samples
113 were determined immediately. DM content was determined by drying the samples at 105 °C in
114 open dishes to constant weight. The remaining samples were stored at – 20 °C for further folate
115 analysis. All the samples of tef flour collected from the 20 households were mixtures of
116 red and white teff varieties.

117 **2.3. Determination of pH**

118 pH was measured using a fresh aliquot of the batter immediately after diluting with deionized
119 water (1:1, v/v) and compared with data in the literature.

120 **2.4. Effect of processing**

121 The effect of traditional household, i.e. fermentation, thermal treatment (baking), and of *injera*
122 processing as a whole on the total folate content of tef *injera* were evaluated and are expressed as
123 percentage retention.

124 **2.5. Contribution of tef *injera* to the recommended nutrient intake (RNI) of folate**

125 Based on the data gathered in the Ethiopian National Food Consumption survey conducted in 2013
126 (EPHI, 2013), we estimated the contribution of tef *injera* to the recommended nutrient intake
127 (RNI) of folate for children aged 1-3, and women of reproductive age. These population groups
128 were selected because they are at an increased risk for folate deficiency.

129

130 **2.6. Folate analysis**

131 The total folate contents of tef flour, batter and *injera* were determined using the reference
132 microbiological method, after tri-enzyme extraction (Kariluoto, 2004). All analytical procedures
133 were carried out under yellow or subdued light. Alternatively, aluminum foil was used to cover
134 the samples and calibrants. Sample extracts were kept under nitrogen atmosphere.

135 **2.6.1. Extraction and tri-enzyme treatment**

136 For the analysis of total folate using the microbiological assay, samples weighing 1 to 1.5 g,
137 depending on the estimated folate content in each sample, were extracted in triplicate (Kariluoto
138 and Piironen, 2009). Extraction was followed by tri-enzyme treatment (α -amylase, hog kidney
139 conjugase and protease) with some modifications: 200 μ L of α -amylase was added in the extracted
140 samples and allowed to settle for 30 min before the pH was adjusted to 4.9 using HCl. This
141 pretreatment facilitated sample homogenization and pH adjustment. Hog kidney conjugase (1 mL)
142 and 800 μ L of α -amylase were then added to the samples. Hog kidney conjugase was prepared
143 from fresh hog kidneys according to Gregory *et al.* (1984). Its activity was tested according to
144 Vahteristo *et al.* (1996). After the enzymes were inactivated in a boiling water bath and cooled on
145 ice, the samples were brought to exactly 25 mL with 0.5% sodium ascorbate and directly analyzed
146 with the microbiological assay.

147

148 **2.6.2. Microbiological assay**

149 Ninety-six-well microtiter plates were used for the assay and the total folate content was
150 determined based on the growth of folate-dependent strain *Lactobacillus rhamnosus* ATCC 7469
151 as the test organism and 5-CHO-H₄ folate as the calibrant. Two dilutions were made from each
152 sample extract using 0.5% sodium ascorbate solution and eight levels of calibrant (0–80 pg/well)
153 in each plate. The plates were incubated for 18 h at 35 °C and turbidity was measured with a
154 microplate reader (Multiskan EX; Labsystems, Helsinki, Finland) at 595 nm. The performance of
155 the method was confirmed by analyzing a blank sample. Certified CRM 121 reference material
156 was analyzed as a quality control in each set of samples. A control chart previously constructed by
157 Kariluoto *et al.* (2004) was used for the folate content of the reference material (certified value
158 500-700 ng/g DM), and a coefficient of variation (CV) < 10% among analytical replicates was
159 considered acceptable.

160 **2.7. Statistical analysis**

161 Statistical analysis of folate was computed using SPSS version 20. The folate analyses were carried
162 out in triplicate and the average values and standard deviations were calculated. Differences
163 between means of folate values in tef flour, batter and *injera* were evaluated using one way-
164 analysis of variance (ANOVA) and Tukey's post hoc test. Differences in means were considered
165 statistically significant with a p-value ≤ 0.05 .

166

167 **3. Results**

168 **3.1. pH of the batter**

169 Measured at 25 °C, the pH of the batter samples ranged from 2.8 to 5.6 with an average of $3.54 \pm$
170 0.4.

171 **3.2. Folate content of tef flour, batter and injera**

172 Microbiologically determined folate contents ranged from 31 to 89 $\mu\text{g}/100$ g DM in tef flour
173 (n=60), 26 to 82 $\mu\text{g}/100$ g DM in batter (n=60) and 21 to 61 $\mu\text{g}/100$ g DM in *injera* (n=60) (Figure
174 2).

175 The average total folate content of *injera* (39 ± 8 $\mu\text{g}/100$ g DM) was significantly lower than that
176 of the batter (52 ± 12 $\mu\text{g}/100$ g DM) and of the flour (59 ± 11 $\mu\text{g}/100$ g DM; $P < 0.05$). High
177 variability was observed among the samples produced by the same household (data not shown).
178 To see if the folate content of samples produced by different households differed, we also
179 compared the households by pairs. After fermentation, only the samples taken from one household
180 had higher folate contents than the other 19 households, and it was the same household where the
181 folate content of the original tef flour was also higher (data not shown).

182 **3.3 Effect of fermentation**

183 Percentage retention of folate after fermentation was calculated by comparing the amount of folate
184 content in tef flour and in the fermented batter on dry matter basis (Figure 3A). A folate retention
185 value $> 100\%$ showed that fermentation increased folate content whereas retention values $< 100\%$
186 indicated folate consumption/losses. Folate retention after fermentation of tef batter ranged from
187 59 to 148% (Figure 3A).

188 Folate retention of more than 100 % was observed in 12/60 cases. In 9/60 cases, retention of about
189 100% was recorded, while in the remaining cases retention was less than 100%.

190 **3.4 Effect of thermal treatment**

191 In all the households, on all three sampling occasions, the folate retention value due to thermal
192 treatment was less than 100%, it ranged from 47 to 96% with average folate retention of 67.7%
193 (Figure 3B).

194 **3.5 Effect of injera processing**

195 The folate retention value due to *injera* processing as a whole ranged from 38 to 97%, showing
196 that the folate content in the final product (*injera*) was almost always lower than the folate content
197 of the original ingredient (tef flour) (Figure 3C). One sample (out of the 60) showed folate retention
198 > 100% (122%, household 5, sampling occasion 2).

199 **3.6 Contribution of tef injera to folate requirements (RNI).**

200 Microbiologically determined folate contents of tef *injera* per fresh weight ranged from 7.1 to 20.1
201 µg/100 g, with an average folate content of 14.3 µg/100 g. The contribution of *injera* to the RNI
202 of folate in the two population groups (children aged from one to three and women of reproductive
203 age) was estimated and the results are listed in Table 1. *Injera* consumption ranged between 23
204 and 66 g/day for children and between 131-202 g/day for women (EPHI, 2013). Using this
205 consumption data, we calculated that folate intake from tef *injera* contributes a maximum of 10%
206 of the RNI for both the children and the women of reproductive age.

207

208 **4. Discussion**

209 To our knowledge, this is the first study to evaluate the folate content of tef flour, fermented tef
210 batter, and tef *injera* to determine the fate of folate in the preparation of this cereal-based fermented

211 Ethiopian food. Our study shows that the whole grain cereal (tef) flour is a relatively good source
212 of folate. The effect of the fermentation step in *injera* processing was highly variable but increased
213 folate content in some cases, whereas baking invariably led to losses. Using available levels of
214 *injera* consumption (EPHI, 2013), we calculated that tef *Injera* contributes up to 10% of the daily
215 folate requirements of vulnerable groups like young children and women of reproductive age.

216

217 The average total folate content of tef flour (52.1 µg/100 g DM) was higher than that reported for
218 other cereals including oats, rice, whole wheat flour and maize (30–40 µg/100 g) but was slightly
219 lower than values reported for sorghum flour (77.0 µg/100 g) (Hager *et al.*, 2012). It is worth
220 noting that our observations of high intra- and interhousehold variability in tef folate content were
221 made possible by repeated and representative sampling. Several factors like the mixture of tef
222 varieties, the milling conditions and the duration and conditions of storage could partly explain the
223 observed variability (Monks *et al.*, 2013; Czarnowska and Gujska, 2012), and such high variability
224 in folate content is common. For example, highly variable folate contents have been reported for
225 oats (49.5–60.4 µg/100 g DM), wheat (36.4–77.4 µg/100 g DM) and rye (64–93 µg/100 g DM
226 (Shewry *et al.*, 2008; Piironen *et al.*, 2008; Kariluoto *et al.*, 2001).

227

228 Recent studies suggest that fermenting cereals has the potential to increase folate content, and that
229 its effectiveness as a strategy to combat folate deficiencies should be further explored (Saubade *et*
230 *al.*, 2017b). The possibility of increasing folate by fermentation was indeed confirmed in our study,
231 as cases in folate retentions of more than 100% were observed. Our results suggest that in these

232 cases of fermentation, folate producing microorganisms dominated those that do not produce or
233 that consume folate. Which microorganisms and which conditions led to folate production in these
234 cases of fermentation warrants further detailed investigations, but this observation already
235 confirms the potential of tef fermentation to produce folate.

236

237 The pH of the fermented batter samples was 3.5 ± 0.4 , which is consistent with data in the literature
238 on tef *injera* fermentation (Baye *et al.*, 2013; Fischer *et al.*, 2014; Yigzaw *et al.*, 2004).

239 For any produced folate to be of use, it will need to survive the baking temperature. Folate is heat
240 labile and can also interact with oxygen in the presence of light (Sotiriadis and Hoskins, 1982).

241 Folate retention relative to the fermented batter was highly variable (47–96%), possibly explained
242 by the difference in *injera* baking conditions (time-temperature). Folate losses due to baking are
243 widely reported (Hefni and Witthöft, 2011; Kariluoto *et al.*, 2004), but the fact that more than 90%
244 of the folates were retained in some examples of baking *injera* suggests that this process could be
245 optimized to minimize folate losses.

246 Some of the folate present in the original flour and synthesized by microorganisms responsible for
247 fermentation can be lost during heat treatment, as reported in previous studies (e.g. Saubade *et al.*,
248 2017a). This means the final folate content of cooked products can be lower than the folate content
249 of the original raw material. In this study, we also showed that the folate content in the final product
250 (*injera*) was almost always lower than the folate content of the original tef flour (Fig. 3C).
251 However, in one case, the final cooked products had higher folate content than the original flour.
252 It has been shown that fermentation can lead to such high folate production that even after cooking,

253 the final products have higher folate contents than the original flour (Kariluoto *et al.*, 2004). This
254 suggests that if appropriate selection is used and microorganisms responsible for folate production
255 are used, it will be possible to increase the amount of folate in the final fermented and baked
256 products. This has been accomplished in other traditional fermented foods made from other cereals
257 such as pearl millet or rye (Greppi *et al.*, 2017; Kariluoto *et al.*, 2006).

258 According to the U.S. Food and Drug administration (FDA, 2016), foods that contribute 10% or
259 more of the daily folate requirements can be considered as good sources of folate. Based on portion
260 size estimates obtained from the national food consumption survey, the maximum folate content
261 in our tef *injera* (20.1 µg/100 g per fresh weight) samples could contribute up to 10% of the RNI
262 of children (1–3 years) and woman of reproductive age. Considering that the national food
263 consumption survey was conducted during the lean season (June–September) when food
264 consumption can be lower (EPHI, 2013), the folate contribution to the RNI is likely
265 underestimated. Assuming two servings (~350 g/serving) of *injera* are consumed per day by
266 women of reproductive age, tef *injera* could contribute up to 25% of the RNI. It is worth noting
267 that *injera* is always consumed with a stew, and that depending on the nature of the stew (e.g.
268 legume based), additional folate intakes may be expected.

269 Several limitations need to be taken into consideration when interpreting our findings. First,
270 although different types of cereals are used to make *injera* (Baye *et al.*, 2015), we focused only on
271 tef *injera*. Second, the extent to which discarding the supernatants contributed to folate loss was
272 not quantified making it difficult to ascribe losses to material loss or possible consumption by
273 microorganisms during fermentation. Third, although the present study revealed that fermentation
274 can increase folate content, evaluating the responsible microorganisms and the conditions that

275 enable folate production were beyond the scope of the present study. However, studies addressing
276 these issues are now underway in our laboratory.

277 Notwithstanding the above limitations, our study quantified the folate content of a widely
278 consumed staple in Ethiopia for the first time. In addition, the study has advanced our
279 understanding of the possible effect of *injera* processing (fermentation and thermal treatment) on
280 folate content.

281 **Conclusion**

282 Tef is a relatively good source of folate and fermenting it to prepare *injera* can increase or decrease
283 its folate content, while baking invariably leads to folate losses. The increase in folate content we
284 observed in some cases, could be attributed to production by the microorganisms involved in
285 fermentation. Reduced folate content could be the result of folate consumption by other
286 microorganisms, or losses due to discarding the supernatant. As fermentation was not controlled,
287 we recorded the net folate content resulting from a balance between the consumption and
288 production of folate by microorganisms. This finding also points to the need for further
289 investigation of the conditions that favor folate production through fermentation while minimizing
290 losses due to thermal treatment (baking). Further studies should also identify the microorganisms
291 (yeast and bacteria) responsible for the folate production during fermentation and their use as a
292 starter culture for the preparation of folate-enriched *injera* should be evaluated. Studies addressing
293 these issues are underway in our laboratory.

294

295

296

297 **Acknowledgements**

298 This study was conducted in the framework of the Folea project (www.folea.eu), which is part
299 of the ERA- Net « Developing African-European joint collaboration for Science and Technology
300 » (ERAfrica), financially support provided by the European Commission under the 7th Framework
301 Programme (ERAfrica ERAFRICA IC-027, FP-226154). Additional support was obtained from
302 the graduate program of Addis Ababa University. The authors would like to thank the families in
303 Addis Ababa who allowed us to collect samples of their private tef fermentations. We are also
304 deeply grateful for Ethiopian Public Health Institute for allowing us access to the National Food
305 Consumption data.

306

307 **Disclosure of interest**

308 The authors report no conflict of interest.

309 **References**

- 310 Bailey L.B., Gregory J.Fr. (2006). Folate, [in:] Present knowledge in nutrition external link icon.
311 B. Bowman and R. Russell (ed.). DC, International Life Sciences Institute, Washington,
312 pp. 278 301.
- 313 Baye, K., Mouquet-Rivier, C., Icard-Vernière, C., Rochette, I., & Guyot, J. P. (2013). Influence of
314 flour blend composition on fermentation kinetics and phytate hydrolysis of sourdough used
315 to make *injera*. *Food Chemistry*, 138(1), 430-436.
- 316 Baye, K., Guyot, J. P., Icard-Verniere, C., Rochette, I., & Mouquet-Rivier, C. (2015). Enzymatic
317 degradation of phytate, polyphenols and dietary fibers in Ethiopian *injera* flours: Effect on
318 iron bioaccessibility. *Food Chemistry*, 174, 60-67
- 319 Czarnowska, M., & Gujska, E. (2012). Effect of freezing technology and storage conditions on
320 folate content in selected vegetables. *Plant foods for human nutrition*, 67(4), 401-406.
- 321 EPHI (2013). Ethiopian national food consumption survey.
322 [https://www.ephi.gov.et/images/pictures/National%20Food%20Consumption%20Survey%20Re](https://www.ephi.gov.et/images/pictures/National%20Food%20Consumption%20Survey%20Report_Ethiopia.pdf)
323 [port_Ethiopia.pdf](https://www.ephi.gov.et/images/pictures/National%20Food%20Consumption%20Survey%20Report_Ethiopia.pdf)
- 324 FDA (2016). Labeling & Nutrition - Guidance for Industry: A Food Labeling Guide (10. Appendix
325 B: Additional Requirements for Nutrient Content Claims).
- 326 Fischer, M. M., Egli, I. M., Aeberli, I., Hurrell, R. F., & Meile, L. (2014). Phytic acid degrading
327 lactic acid bacteria in tef-injera fermentation. *International journal of food*
328 *microbiology*, 190, 54-60.

329 Gregory, J.F., Sartain, D.B., & Day, B.P.F. (1984). Fluorometric determination of folacin in
330 biological materials using high performance liquid chromatography. *Journal of Nutrition*.
331 114: 341-353.

332 Greppi, A., Hemery, Y., Berrazaga, I., Almaksour, Z., & Humblot, C. (2017). Ability of
333 lactobacilli isolated from traditional cereal-based fermented food to produce folate in
334 culture media under different growth conditions. *LWT*, 86, 277-284.

335 Gujska, E., & Majewska, K. (2005). Effect of baking process on added folic acid and endogenous
336 folates stability in wheat and rye breads. *Plant Foods for Human Nutrition*, 60(2), 37-42.

337 Hager, A. S., Wolter, A., Jacob, F., Zannini, E., & Arendt, E. K. (2012). Nutritional properties and
338 ultra-structure of commercial gluten free flours from different botanical sources compared
339 to wheat flours. *Journal of Cereal Science*, 56(2), 239-247.

340 Haidar, J. (2010). Prevalence of anaemia, deficiencies of iron and folic acid and their determinants
341 in Ethiopian women. *Journal of health, population, and nutrition*, 28(4), 359.

342 Hefni, M., & Witthöft, C. M. (2011). Increasing the folate content in Egyptian baladi bread using
343 germinated wheat flour. *LWT-Food Science and Technology*, 44(3), 706-712.

344 Ho, R. C., Cheung, M. W., Fu, E., Win, H. H., Zaw, M. H., Ng, A., & Mak, A. (2011). Is high
345 homocysteine level a risk factor for cognitive decline in elderly? A systematic review,
346 meta-analysis, and meta-regression. *The American Journal of Geriatric Psychiatry*, 19(7),
347 607-617.

348 Holzapfel, W. H. (2002). Appropriate starter culture technologies for small-scale fermentation in
349 developing countries. *International Journal of Food Microbiology*, 75(3), 197-212.

350 Humblot, C., & Guyot, J. P. (2008). Other fermentations. In *Molecular Techniques in the*
351 *Microbial Ecology of Fermented Foods* (pp. 208-224). Springer New York.

352 Kariluoto, M. S., Vahteristo, L. T., & Piironen, V. I. (2001). Applicability of microbiological assay
353 and affinity chromatography purification followed by high-performance liquid
354 chromatography (HPLC) in studying folate contents in rye. *Journal of the Science of Food
355 and Agriculture*, 81(9), 938-942.

356 Kariluoto S, Vahteristo L, Salovaara H, Katina K, Liukkonen K, Piironen V. (2004). Effect of
357 baking method and fermentation on folate content of rye and wheat breads. *Cereal
358 Chemistry* 81(1):134-139.

359 Kariluoto, S., Aittamaa, M., Korhola, M., Salovaara, H., Vahteristo, L., & Piironen, V. (2006).
360 Effects of yeasts and bacteria on the levels of folates in rye sourdoughs. *International
361 journal of food microbiology*, 106(2), 137-143.

362 Kariluoto, S., & Piironen, V. (2009). Total folate. In P. Shewry, & J. Ward (Eds.).
363 HEALTHGRAIN methods, analysis of bioactive components in small grain cereals. St
364 Paul, Minnesota, USA : AACC International.

365 Keagy, P. M., Stokstad, E. L. R., & Fellers, D. A. (1975). Folacin stability during bread processing
366 and family flour storage. *Cereal Chemistry*.

367 LeBlanc, J. G., de Giori, G. S., Smid, E. J., Hugenholtz, J., & Sesma, F. (2007). Folate production
368 by lactic acid bacteria and other food-grade microorganisms. *Communicating Current
369 Research and Educational Topics and trends in Applied Microbiology*, 1, 329-339.

370 Lee, S. E., Talegawkar, S. A., Merialdi, M., & Caulfield, L. E. (2013). Dietary intakes of women
371 during pregnancy in low-and middle-income countries. *Public health nutrition*, 16(8),
372 1340-1353.

373 McLean, E., de Benoist, B., & Allen, L. H. (2008). Review of the magnitude of folate and vitamin
374 B12 deficiencies worldwide. *Food and Nutrition Bulletin*, 29(2_suppl1), S38-S51.

375 Monks, J. L. F., Vanier, N. L., Casaril, J., Berto, R. M., de Oliveira, M., Gomes, C. B., Carvalho,
376 M. P., Dias, A. R. G., & Elias, M. C. (2013). Effects of milling on proximate composition,
377 folic acid, fatty acids and technological properties of rice. *Journal of Food Composition
378 and Analysis*, 30(2), 73-79.

379 Moore, L. L., Bradlee, M. L., Singer, M. R., Rothman, K. J., & Milunsky, A. (2003). Folate intake
380 and the risk of neural tube defects: an estimation of dose-response. *Epidemiology*, 200-205.

381 Moslehi-Jenabian, S., Lindegaard, L., & Jespersen, L. (2010). Beneficial effects of probiotic and
382 food borne yeasts on human health. *Nutrients*, 2(4), 449-473.

383 Piironen, V., Edelmann, M., Kariluoto, S., & Bedó, Z. (2008). Folate in wheat genotypes in the
384 HEALTHGRAIN diversity screen. *Journal of Agricultural and Food Chemistry*, 56(21),
385 9726-9731.

386 Saubade, F., Hemery, Y. M., Guyot, J. P., & Humblot, C. (2017a). Lactic acid fermentation as a
387 tool for increasing the folate content of foods. *Critical Reviews in Food Science and
388 Nutrition*, 57(18), 3894-3910.

389 Saubade, F., Humblot, C., Hemery, Y. M., & Guyot, J. P. (2017b). PCR screening of an African
390 fermented pearl-millet porridge metagenome to investigate the nutritional potential of its
391 microbiota. *International Journal of Food Microbiology*, 244, 103-110.

392 Scholl, T. O., & Johnson, W. G. (2000). Folic acid: influence on the outcome of pregnancy. *The
393 American Journal of Clinical Nutrition*, 71(5), 1295s-1303s.

394 Shewry, P. R., Piironen, V., Lampi, A. M., Nyström, L., Li, L., Rakszegi, M., Fras, A., Boros, D.,
395 Gebruers, K., Courtin, C.M., & Delcour, J. A. (2008). Phytochemical and fiber components
396 in oat varieties in the HEALTHGRAIN diversity screen. *Journal of Agricultural and Food*
397 *Chemistry*, 56(21), 9777-9784.

398 Sotiriadis, P. K., & Hoskins, F. H. (1982). Vitamin retention during storage of processed foods. I.
399 Effect of ascorbic acid on folates in cowpeas, okra and tomatoes. *Scientia Horticulturae*,
400 16(2), 125-130.

401 Vahteristo, L., Ollilainen, V., Koivistoinen, P., and Varo, P. (1996). Improvements in the analysis
402 of reduced folate monoglutamates and folic acid in food by high-performance liquid
403 chromatography. *Journal of Agricultural and Food Chem.* 44: 477-482.

404 World Health Organization. (2004). *Vitamin and mineral requirements in human nutrition: report*
405 *of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21-30 September 1998* (No.
406 Ed. 2). World Health Organization.

407 Yetneberk, S., de Kock, K.L., Rooney, L.W., Taylor, J.R.N. (2004). Effects of sorghum cultivar
408 on injera quality. *Cereal Chemistry*, 81, 314-321

409 Yigzaw, Y., Gorton, L., Solomon, T., & Akalu, G. (2004). Fermentation of seeds of Teff
410 (*Eragrostis teff*), grass-pea (*Lathyrus sativus*), and their mixtures: aspects of nutrition and
411 food safety. *Journal of agricultural and food chemistry*, 52(5), 1163-1169.

412 Zhu, F. (2018). Chemical composition and food uses of teff (*Eragrostis tef*). *Food chemistry*, 239,
413 402-415.

414

415

416 **Table 1** Calculated contribution of tef *injera* to folate requirements

Population group	Age (years)	RNI ($\mu\text{g/day}$ in DFEs) ^a	<i>Injera</i> intake (g/day) ^b	Contribution to RNI (%)	
Children	1-3 years	150	Minimum	23.0	2.2
			Maximum	108.1	10.3
			Average	66.1	6.3
Women	19-45	400	Minimum	130.7	3.7
			Maximum	291.2	10.4
			Average	202.4	7.2


417 ^a Recommended nutrient intakes (RNIs) are from WHO, 2004; DFEs, Dietary folate equivalents

418 ^b Estimated using consumption values obtained from the Ethiopian National Food Consumption Survey (EPHI, 2013)

419

420

421



422

423 **Figure 1.** Flow diagram of the traditional *injera* making process observed in Ethiopian urban
 424 households.

425 *samples were taken for folate analysis

426 “*Ersho*”: inoculum used for *injera* making, it is a leftover from a previous successful
 427 spontaneous traditional fermentation.


428

429

430

431

432



433

434 **Figure 2.** Box plot showing the distribution of total folate content of tef flour (n=60), fermented
 435 batter (n=60) and *injera* (n=60) in $\mu\text{g}/100\text{ g DM}$

436 The lower edge of the box corresponds to the 25th percentile, the upper edge to the 75th percentile,
 437 and the line across the middle corresponds to the median (50th percentile). The vertical lines
 438 extending outside the box represent the full range of observations.

439 DM, Dry matter basis; different superscript letters indicate a statistically significant difference (p
 440 < 0.05).

441

442


443

444

445

446

447



448

449

450

451 **Figure 3.** Folate retention (%) due to fermentation, thermal treatment and *injera* processing

452 A: Folate retention (%) after 1st stage fermentation; $\text{Folate retention (\%)} = \text{Folate}_{\text{flour}} / \text{Folate}_{\text{batter}} \times 100$.

453 B: Folate retention (%) during cooking; $\text{Folate retention (\%)} = \text{Folate}_{\text{batter}} / \text{Folate}_{\text{injera}} \times 100$.

454 C: Folate retention (%) due to *injera* processing; $\text{Folate retention (\%)} = \text{Folate}_{\text{flour}} / \text{Folate}_{\text{injera}} \times 100$.