

<https://helda.helsinki.fi>

---

## Iron-Phosphorus Feedbacks Drive Multidecadal Oscillations in Baltic Sea Hypoxia

Jilbert, Tom

2021-12-28

---

Jilbert , T , Gustafsson , B G , Veldhuijzen , S , Reed , D C , Helmond , N A G M , Hermans , M & Slomp , C P 2021 , ' Iron-Phosphorus Feedbacks Drive Multidecadal Oscillations in Baltic Sea Hypoxia ' , Geophysical Research Letters , vol. 48 , no. 24 , ARTN e2021GL095908 . <https://doi.org/10.1029/2021GL095908>

---

<http://hdl.handle.net/10138/338546>

<https://doi.org/10.1029/2021GL095908>

---

cc\_by\_nc

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Geophysical Research Letters®



## RESEARCH LETTER

10.1029/2021GL095908

### Key Points:

- Past hypoxic intervals in the Baltic Sea were characterized by multidecadal oscillations in oxygen stress
- Regularly paced internal oscillations caused by feedbacks in coupled iron-phosphorus dynamics
- External loading of phosphorus and climate forcing dictate the amplitude of internal oscillatory behavior

### Supporting Information:

Supporting Information may be found in the online version of this article.

### Correspondence to:

T. Jilbert,  
[tom.jilbert@helsinki.fi](mailto:tom.jilbert@helsinki.fi)

### Citation:

Jilbert, T., Gustafsson, B. G., Veldhuijzen, S., Reed, D. C., van Helmond, N. A. G. M., Hermans, M., & Slomp, C. P. (2021). Iron-phosphorus feedbacks drive multidecadal oscillations in Baltic Sea hypoxia. *Geophysical Research Letters*, 48, e2021GL095908. <https://doi.org/10.1029/2021GL095908>

Received 31 AUG 2021

Accepted 29 NOV 2021

### Author Contributions:

**Conceptualization:** Tom Jilbert, Bo G. Gustafsson, Caroline P. Slomp

**Data curation:** Tom Jilbert

**Formal analysis:** Tom Jilbert, Simon Veldhuijzen, Daniel C. Reed, Niels A. G. M. Helmond, Martijn Hermans, Caroline P. Slomp

**Funding acquisition:** Tom Jilbert, Caroline P. Slomp

**Investigation:** Tom Jilbert, Bo G. Gustafsson, Simon Veldhuijzen, Daniel C. Reed, Niels A. G. M. Helmond, Martijn Hermans, Caroline P. Slomp

**Methodology:** Tom Jilbert, Bo G. Gustafsson, Simon Veldhuijzen, Daniel C.

## Iron-Phosphorus Feedbacks Drive Multidecadal Oscillations in Baltic Sea Hypoxia

Tom Jilbert<sup>1,2,3,4</sup> , Bo G. Gustafsson<sup>2,5</sup> , Simon Veldhuijzen<sup>4</sup>, Daniel C. Reed<sup>4,6</sup>, Niels A. G. M. van Helmond<sup>4</sup> , Martijn Hermans<sup>1,3,4</sup> , and Caroline P. Slomp<sup>4</sup> 

<sup>1</sup>Aquatic Biogeochemistry Research Unit (ABRU), Ecosystems and Environment Research Program, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland, <sup>2</sup>Tvärminne Zoological Station, University of Helsinki, Hanko, Finland, <sup>3</sup>Department of Geosciences and Geography, Environmental Geochemistry Group, Faculty of Science, University of Helsinki, Helsinki, Finland, <sup>4</sup>Department of Earth Sciences (Geochemistry), Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands, <sup>5</sup>Baltic Nest Institute, Baltic Sea Centre, Stockholm University, Stockholm, Sweden, <sup>6</sup>Fisheries & Oceans Canada, Bedford Institute of Oceanography, Dartmouth, NS, Canada

**Abstract** Hypoxia has occurred intermittently in the Baltic Sea since the establishment of brackish-water conditions at ~8,000 years B.P., principally as recurrent hypoxic events during the Holocene Thermal Maximum (HTM) and the Medieval Climate Anomaly (MCA). Sedimentary phosphorus release has been implicated as a key driver of these events, but previous paleoenvironmental reconstructions have lacked the sampling resolution to investigate feedbacks in past iron-phosphorus cycling on short timescales. Here we employ Laser Ablation (LA)-ICP-MS scanning of sediment cores to generate ultra-high resolution geochemical records of past hypoxic events. We show that in-phase multidecadal oscillations in hypoxia intensity and iron-phosphorus cycling occurred throughout these events. Using a box model, we demonstrate that such oscillations were likely driven by instabilities in the dynamics of iron-phosphorus cycling under preindustrial phosphorus loads, and modulated by external climate forcing. Oscillatory behavior could complicate the recovery from hypoxia during future trajectories of external loading reductions.

**Plain Language Summary** Hypoxia in the coastal ocean is expanding worldwide, and inputs of nutrients from waste water and agriculture are mainly to blame. Nutrients feed plankton blooms, which then consume oxygen as they decay. Because much of this decay takes place at the seafloor, sediments play an important role in deoxygenation, and in the recycling of nutrients in coastal regions. It is known that the amount of iron oxides in sediments has a strong effect on nutrient recycling during algal decay. Iron oxide-rich sediments in healthy oxygen-rich areas can trap phosphorus, a key nutrient element from decaying algae. In contrast, iron oxide-poor sediments in deoxygenated areas release phosphorus back to the water to fuel more algal growth. Our study shows that changes in the distribution of iron oxides between deep and shallow areas of the Baltic Sea led to self-sustaining variability (oscillations) in oxygen stress on decadal timescales during past intervals in the Sea's 8000-year history. We use a model to demonstrate that under certain conditions of climate and nutrient pressure, such variability may occur naturally, and therefore may influence the future recovery of the Baltic Sea from its present nutrient-rich state.

## 1. Introduction

Marginal marine basins such as the Baltic Sea, Black Sea and many estuaries are naturally susceptible to hypoxia due to stratification and isolation of deep water masses (Conley et al., 2009; Eckert et al., 2013; Lefort et al., 2012). Hypoxia has been further exacerbated in modern coastal systems by anthropogenic nutrient inputs, leading to an expansion of marine 'dead zones' worldwide (Rabalais et al., 2010). Positive feedbacks in the phosphorus (P) cycle play a key role in intensifying hypoxic conditions in aquatic systems. Phosphorus release from sediments accelerates under low-oxygen conditions, due to the dissolution of iron oxide-bound P (Fe-P; Mortimer, 1941) and the preferential regeneration of P from organic matter (Ingall et al., 1993). Enhanced sedimentary P release, in turn, sustains high productivity and oxygen demand (Vahtera et al., 2007; Van Cappellen & Ingall, 1994).

An important caveat to the positive feedback between P cycling and hypoxia is that in spatially finite systems, the release of Fe-P from sediments cannot proceed indefinitely. Low-oxygen conditions at any given location may rapidly exhaust the local sedimentary inventory of Fe-P (Jilbert et al., 2011; Reed et al., 2011), while the

© 2021 The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](https://creativecommons.org/licenses/by-nc/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Reed, Niels A. G. M. Helmond, Caroline P. Slomp

**Project Administration:** Tom Jilbert,

Caroline P. Slomp

**Resources:** Caroline P. Slomp

**Software:** Bo G. Gustafsson

**Supervision:** Caroline P. Slomp

**Validation:** Bo G. Gustafsson, Daniel C.

Reed, Niels A. G. M. Helmond, Martijn

Hermans, Caroline P. Slomp

**Visualization:** Tom Jilbert, Bo G.

Gustafsson, Daniel C. Reed, Martijn

Hermans

**Writing – original draft:** Tom Jilbert,

Bo G. Gustafsson, Caroline P. Slomp

**Writing – review & editing:** Tom Jilbert,

Bo G. Gustafsson, Simon Veldhuijzen,

Daniel C. Reed, Niels A. G. M. Helmond,

Martijn Hermans, Caroline P. Slomp

maximum area of potential hypoxia may be limited by thermohaline stratification (Carstensen et al., 2014). Theoretically, if external P loads are insufficient to maintain high productivity after the exhaustion of the Fe-P pool, a reverse of the positive feedback may be induced, leading to sequestration of P and abatement of hypoxia. This dual directionality of the Fe-P feedback has been proposed to explain the rapid transitions at the onset and termination of centennial-timescale hypoxic events in the Baltic Sea (Jilbert & Slomp, 2013).

The hypoxic events defined in Jilbert and Slomp (2013) are subdivisions of longer hypoxic intervals (HI) in the sedimentary record of the Baltic Sea (Zillén et al., 2008), which occurred during the Holocene Thermal Maximum (HTM<sub>HI</sub>, ~8,000–4,000 cal. yr. B.P.), the Medieval Climate Anomaly (MCA<sub>HI</sub>, 1,200–750 cal. yr. B.P.) and the modern period (Mod<sub>HI</sub>, ~1980–present). Hypoxic conditions during these intervals are thought to have been induced by a combination of physical oceanographic forcing and external nutrient inputs, and sustained by P cycle feedbacks. Due to constraints on the sampling resolution of sediment cores, investigation of multiannual to multidecadal variability in past hypoxia during these intervals is challenging. However, such variability is essential to study because of the relevance for managing the recovery from modern hypoxia, one major goal of the Baltic Sea Action Plan (HELCOM, 2007, 2021). Here, we present new, ultra-high resolution records of past Baltic Sea hypoxia and Fe-P dynamics using LA-ICP-MS line scanning of resin-embedded sediments. These records cover the entire HTM<sub>HI</sub> and MCA<sub>HI</sub> intervals and thus significantly extend the initial LA-ICP-MS data presented in Jilbert and Slomp (2013). Using time-dependent box-model, we show that multidecadal oscillations observed in the sediment proxies could plausibly have been sustained by instabilities in the feedbacks between Fe-P dynamics and hypoxia on these timescales.

## 2. Materials and Methods

### 2.1. Sediment Sampling

A gravity core (0–406 cm) was collected from site F80 (58°00.00′N, 19°53.81′E, water depth 191 m) in the Fårö Deep of the Baltic Sea during the HYPER/COMBINE cruise of R/V Aranda (May/June 2009), and sectioned in a nitrogen-filled glovebox at 0.5–2.0 cm intervals. Undisturbed U-channel subsamples were taken from laminated intervals and embedded with Spurr's epoxy resin by the fluid displacive method under nitrogen (Jilbert et al., 2008).


### 2.2. Sample Preparation and Geochemical Analysis

Discrete sediment samples were freeze-dried and ground under nitrogen. Aliquots of 0.125 g dry sediment were digested in 2.5 ml HF (40%) and 2.5 ml of HClO<sub>4</sub> (70%)/HNO<sub>3</sub> (65%) mixture at 90°C overnight, before evaporation at 160°C and redissolution in 1M HNO<sub>3</sub>. Solutions were analyzed by ICP-OES for total Al, Fe and Mo and P (precision and accuracy <10%).

Resin-embedded sediment trays were cut to expose the internal surface of the sediment, and sectioned into shorter blocks to fit inside the Laser Ablation (LA) ICP-MS sampling chamber. LA-ICP-MS line scanning was performed at 0.04 mm s<sup>-1</sup> using an Excimer laser (193 nm, spot size 120 μm, repetition rate 10 Hz, fluence 8 J cm<sup>-2</sup>) coupled to an Element 2 ICP-MS (Hennekam et al., 2015). Count intensities of the following isotopes were measured for use in this study: <sup>27</sup>Al, <sup>31</sup>P, <sup>57</sup>Fe, <sup>79</sup>Br, and <sup>98</sup>Mo. Calibration of the LA-ICP-MS data was performed according to a modified version of the two-step procedure described in the supplement of Jilbert and Slomp (2013). Full details of the calibration procedure are given in the Supporting Information S1.

### 2.3. Rationale of the Geochemical Proxies

Sedimentary Mo/Al and Fe/Al are established proxies for hypoxia intensity and Fe shuttling, respectively (Jilbert & Slomp, 2013; Lenz et al., 2015). The ratio of bromine to phosphorus (Br/P) is here used as a surrogate for C/P, an indicator of the intensity of preferential P regeneration (Jilbert & Slomp, 2013). The rationale for using Br/P is that C cannot be measured in the epoxy-embedded samples by LA-ICP-MS, but Br and C are strongly correlated in sediments due to the bromination of organic matter by seawater bromide during degradation (Ziegler et al., 2008), hence LA-ICP-MS-derived Br provides a surrogate for C. Absolute Br contents were also determined on selected samples by repeated digestion in H<sub>2</sub>O<sub>2</sub> followed by ICP-MS analysis (Ziegler et al., 2008).



**Figure 1.** Key features of the box model used in this study. (a): Schematic of the model design. Surface water and deep water boxes (blue) are each underlain by a reactive sediment layer (gray). Six time-dependent state variables are included:  $P$  = water column phosphorus concentration,  $O_2$  = water column oxygen concentration (deep water only),  $C_{org}$  = sediment organic carbon inventory,  $P_{org}$  = sediment organic phosphorus inventory, Fe-P = sediment iron-bound phosphorus inventory. Equations governing exchange fluxes and their relationship to state variables are given in the Supporting Information (Text S3, Figure S4). (b): Principle of the sigmoid function used in the model. The attainable Fe-P inventory of the sediments is controlled by the dimensionless ratio between oxygen flux to sediments and sedimentation rate of organic matter ( $O_2$  supply demand ratio).

These values were used to generate discrete sample Br/Al ratios for the final step of the calibration of the LA-ICP-MS data.


#### 2.4. Sediment Dating and Time Series Analysis

The age-depth model used for the sediments in this study is described in the supplement of Jilbert and Slomp (2013). Briefly, the discrete-sample  $C_{org}$  profile was tuned to the Loss on Ignition (LOI) profile of reference core 372740-3 from the Gotland Deep (Lougheed et al., 2012). 372740-3 is dated using known paleomagnetic secular variation (PSV) features and lead (Pb) isochrons (12 features total in the 406 cm core covering ~7,000 years, giving an average spacing of ~600 years). Linear sedimentation rate is assumed between dating points. In the early part of the record, this age-depth model differs by up to 500 years from a recently published model for Baltic Sea sediments based on benthic foraminiferal  $^{14}C$  dating (Warden et al., 2017). However, this offset is not expected to impact on the conclusions of the present study, which focuses on frequencies of variability rather than absolute ages of events.

Time series analysis of the LA-ICP-MS Mo/Al, Fe/Al, and Br/P profiles was performed in Analyseries 1.1.1 (Paillard et al., 1996). Raw elemental ratios were detrended and normalized to unit variance, resampled to 1 year resolution and processed by Fast Fourier Transform (FFT, Blackman-Tukey method with Bartlett window) to yield a power spectrum of cyclic components in each profile. Cross-spectral analysis (same settings as above) was also applied to investigate coherence and phase relations between the elemental ratio data. A 20–100-year band-pass filter was subsequently applied to the detrended, normalized data for visual comparison of the time series.

#### 2.5. Box Modeling

A box model was created to represent the coupled cycling of P and Fe in the Baltic Sea. The basic formulation is the same for the  $HTM_{HI}$ ,  $MCA_{HI}$ , and  $Mod_{HI}$  (Figure 1a), with six state variables related by a system of equations. A full model description, including calibration and validation against present day data, is provided in the Supporting Information S1. Production of organic matter in surface waters is limited by the phosphorus concentration  $P$ . Organic matter sinks vertically and accumulates in the sediments, where remineralization occurs. Phosphorus



**Figure 2.** LA-ICP-MS line scan data of Baltic Sea sediments from (a)  $HI_{HTM}$  and (b)  $HI_{MCA}$ . (left) High-resolution geochemical profiles of sediments from site F80. Horizontal gray bars indicate the subdivision of  $HTM_{HI}$  and  $MCA_{HI}$  into numbered hypoxic events, as given in Jilbert and Slomp (2013). Upper panels = calibrated LA-ICP-MS profiles of Mo/Al. Lower panels = 20–100 years bandpass-filtered profiles of Mo/Al, Br/P and Fe/Al (detrended and normalized to unit variance prior to filtering). (center) Blackman-Tukey spectral analysis of Mo/Al, Br/P, and Fe/Al data for the entire time intervals shown on the left. Gray field indicates the period 60–100 years. (right) Coherence and phase analysis of Mo/Al and Br/P (solid lines) and Mo/Al and Fe/Al (dashed lines) for the entire time intervals shown on the left. Phase in radians (0 = in phase,  $\Pi$  = antiphase).

may be regenerated from sediments due to remineralization or release from sedimentary Fe-P, or buried. Oxygen is constant in the surface layer but allowed to vary in the deep-water box, simulating the observed oxygen depletion in the sub-halocline water column of the Baltic Sea (Carstensen et al., 2014). A key feature of the model is a sigmoid function describing the relationship between the sedimentary Fe-P inventory and the oxygen *supply-demand ratio*, the dimensionless ratio between oxygen flux to sediments and oxygen demand by organic matter mineralization (Figure 1b). The oxygen flux to sediments is directly proportional to oxygen concentration. Advective exchange of dissolved constituents is permitted, both vertically between the two layers and laterally between these layers and a hypothetical adjacent sea (simulating the real exchange between the Baltic and the North Sea). External loading of P is prescribed for each time interval based on literature estimates. The sensitivity and stability of steady state solutions to changes in external loads and vertical exchange were assessed. Simulations with variable frequency and amplitude of vertical mixing were conducted to demonstrate the potential of variable climate forcing to trigger and control oscillatory behavior. Spectral analysis was performed on 10,000-year long annual oxygen time-series using FFT with Hamming window of the autocovariance limited to 512 time steps.

### 3. Results and Discussion

#### 3.1. Evidence for Oscillations in Past Hypoxia and Phosphorus Regeneration

During  $HTM_{HI}$  and  $MCA_{HI}$ , multidecadal oscillations are observed in the LA-ICP-MS line scan profiles of Mo/Al and Br/P (Figure 2, left). Spectral analysis of the full data series shows multiple peaks across the decadal to multidecadal bands. A clear peak is observed in the 60–100-year band in both parameters (Figure 2, center). This band is also characterized by high coherence and close-to-zero phase relation between Mo/Al and Br/P (Figure 2, right), indicating in-phase oscillatory signals. Especially in the intervals of lower overall Mo/Al (e.g.,

1.6–1.2 ky BP; 5.4–5.0 ky BP), the oscillation is clearly visible in the raw and bandpass-filtered data (Figure 2, left and Figures S2 and S3). This result implies that a close coupling existed between hypoxia and phosphorus regeneration on multidecadal timescales during past hypoxic events, and that both were highly variable on these timescales.

### 3.2. Evidence for Oscillations in Past Fe Shuttling

In the intervals where the oscillations in Mo/Al and Br/P are most pronounced (e.g., 1.6–1.2 ky BP; 5.3–4.9 ky BP), the LA-ICP-MS data also show similarly paced variability in Fe/Al (Figure 2, left and Figures S2 and S3). Accordingly, a similar peak in the 60–100-year band is observed in the power spectrum of the Fe/Al data (Figure 2, center), as well as high-coherence and close-to-zero phase relation between Mo/Al and Fe/Al (Figure 2, right). This indicates that shelf-to-basin shuttling of Fe in the Baltic Sea was also sensitive to multidecadal variability in hypoxia. Namely, during periods of low deep-water oxygen, more Fe was transported laterally downslope into the deep basins via cycles of dissolution and reprecipitation (Lyons & Severmann, 2006).

### 3.3. Oscillations in Box Model Simulations

Model simulations suggest that multidecadal oscillations in hypoxia and phosphorus regeneration may have been an intrinsic feature of biogeochemical cycles in the Baltic Sea under the forcing conditions of the  $HTM_{HI}$  and  $MCA_{HI}$  (Figure 3). Steady-state solutions of the model are unstable when parameters including basin geometry and external loading of P are set to constant realistic values for these intervals, indicating the presence of unforced oscillations. The periodicity of these unforced oscillations in the model is typically 130–170 years, slightly longer than observed in the sediment records (Figure 3). Within each oscillation, during the period of low deep water oxygen, the sediment Fe-P inventory is at a minimum, whereas deep water phosphorus ( $P$ ), and sediment organic carbon ( $C_{org}$ ) and phosphorus ( $P_{org}$ ), show maximum values (Figure 3). Conversely, when deep water oxygen is high, the opposite trends are observed.


### 3.4. Mechanism and Frequency of the Oscillations

The presence of relatively large amounts of Fe-P in Baltic Sea sediments under low P loading conditions is a prerequisite for the observed instability. The quantitative representation of the sigmoid function used in the model (Figure S5) shows that the sensitivity of the Fe-P inventory to the oxygen supply-demand ratio is high at intermediate values ( $Fe-P = 50-150 \text{ mmol m}^{-2}$ ). During the *recharge* phase (Figure 1b), storage of Fe-P in sediments provides a reinforcing feedback toward a higher supply-demand ratio. Conversely, the *discharge* phase is characterized by release of Fe-P and a reinforcing feedback toward lower supply-demand ratio. However when oxygen is plentiful and productivity low (high values of the supply-demand ratio), the sediment Fe-P pool becomes increasingly saturated (close to  $200 \text{ mmol m}^{-2}$  Fe-P), leading to a leveling-off in the Fe-P inventory. Similarly, when oxygen is scarce and productivity high, the Fe-P concentration levels off due to the approaching exhaustion of the sedimentary Fe-P pool. This insensitivity at high and low values makes the system vulnerable to a switch in directionality.


Even in our relatively simple model formulation, the frequency of the intrinsic oscillations is a complex function of the prescribed parameters. Frequency is influenced partly by the geometry of the Baltic Sea basins, which is a major factor controlling the residence time of P in both the shallow and deep layers in the model. Indeed, the periodicity of simulated oscillations during the  $HTM_{HI}$ , when the subhalocline hypoxic area of the Baltic was expanded due to glacio-isostatic effects (Jilbert et al., 2015) is slightly longer than that during the  $MCA_{HI}$  (Figures 3a–3d). However, the frequency of real-world oscillations may also be externally driven. Indeed, the inherent instability of the model can be triggered by variability in external driving forces as discussed below.

### 3.5. Influence of Climatic Drivers on Oscillations

The observed oscillations in the sediment records indicate dominant periods of 60–100 years, which is consistent with periods of the Atlantic Multidecadal Oscillation (AMO), or Atlantic Multidecadal Variability (AMV), that may have been present throughout the Holocene (Knudsen et al., 2011). The AMO in turn modulates the influence of the North Atlantic Oscillation (NAO) on the conditions in the Baltic Sea region (Börgel et al., 2020).



**Figure 3.** State variables in box model simulations of biogeochemical cycles during past hypoxic intervals in the Baltic Sea. (a and b): sedimentary P inventories, (c and d): oxygen concentration in deep water box, (e): sensitivity analysis showing mean values, at steady state, of state variables in response to changes in external P loading. (f): sensitivity analysis showing mean values, at steady state, of state variables in response to changes in vertical exchange between shallow and deep-water boxes. Blue shaded areas in (e–f) indicate the range in P input and vertical exchange, where the steady state solutions are unstable, according to linear stability analysis, causing oscillations in time-dependent solutions.



**Figure 4.** Power spectra of deep-water oxygen concentrations for different cases of imposed cyclic variation of the vertical exchange rate (amplitude in  $\text{m y}^{-1}$  as deviation from  $11 \text{ m y}^{-1}$ , the  $\text{MCA}_{\text{HI}}$  default value), representing variability of climatic forcing. Each row represents a given period ( $T$ ) and the columns different amplitude of the oscillation. The dashed line graphs show the power spectra of the inherent oscillation with constant vertical exchange rate (cf. Figure 3c).

Thus, it is reasonable to assume that the observed oscillations in Baltic Sea sediment records may to some extent have been externally driven by climatic variability as well as the internal instability of biogeochemical cycles.

To explore the impact of climatic variability, we imposed a sinusoidal perturbation to the vertical exchange relative to the parameter settings for the model simulations of the  $\text{MCA}_{\text{HI}}$ . Vertical exchange is chosen because it has a direct and influential impact on oxygen conditions and subsequent effects on the Fe-P inventory. In Figure 4, oxygen power spectra are shown for a number of simulations with different amplitudes and periods of the vertical exchange perturbation. It is evident that for certain periods of climatic oscillations, the inherent model instability is triggered and the oxygen oscillations are strongly amplified compared to the unforced case (e.g., for amplitude =  $3 \text{ m y}^{-1}$  and  $T = 80$  and  $100$  years). In such cases, the period of oscillations is largely determined by the climate. This external influence could partly explain the generally shorter periods observed in the sediment records (60–100 years, similar to AMO/AMV forcing) compared to the inherent oscillations of the model (130–170 years). In terms of actual historical variability in AMO/AMV forcing, we note that Wang et al. (2017) showed the magnitude of AMV in the 60–100-year band to be higher in the centuries following the  $\text{MCA}_{\text{HI}}$  than during it. However, since the Baltic Sea was in a stable oxic state after 0.8 ky BP (Jilbert & Slomp, 2013) there was no amplification of the climatic signal to be observed in the sediment records.

### 3.6. Sensitivity to External Phosphorus Loading and Vertical Exchange

The presence of multidecadal oscillations in the model simulations is highly sensitive to external loading of P. Using a parameterization for the modern Baltic Sea, external loading of <8,000 tonnes P/year yields solutions in which no oscillations are observed because the system remains in a quasipermanent oxic state (Figure 3e). As external loading increases, oscillations are observed within a range of ~8,000–12,000 tonnes P/year. However at high external loading (>12,000 tonnes P/year), oscillations disappear again because the system remains in a quasipermanent hypoxic state.

The modern P loading to the Baltic Proper is ~25,000 tonnes P/year, and loading during the late 20th century was significantly in excess of this value (Gustafsson et al., 2012). Such high loads are sufficient to suppress oscillatory behavior in the model, by sustaining high productivity and holding the system in a quasipermanent hypoxic state. A simulation of the modern development of hypoxia since 1900 supports this observation (Figures S6 and S7). Therefore, under constant future loading scenarios, we would not expect to observe multidecadal oscillations in Baltic Sea hypoxia during the coming centuries.

Although enhanced nutrient loading to the Baltic during the MCA has been suggested (Zillén & Conley, 2010), little evidence for eutrophication has been found in coastal paleoenvironmental data from this period (Jokinen et al., 2018; Ning et al., 2016; Norbäck Ivarsson et al., 2019). Therefore, we imposed an external loading of 8,000 tonnes P/year for both the MCA<sub>HI</sub> and HTM<sub>HI</sub> simulations, as derived from estimates of the preindustrial loading to the Baltic Sea (Gustafsson et al., 2012). Because this value is at the lower limit of the range yielding oscillations in most parameterizations (Figures 3c and 3d), we adjusted the vertical exchange parameter *D* in MCA<sub>HI</sub> and HTM<sub>HI</sub> simulations to ~20% lower than the modern value (Table S1). Lower values of *D* favor stagnation and the development of deep water hypoxia, and tip the system into an oscillatory state at external loading of <8,000 tonnes P/year (Figures S8 and S9). Our simulations therefore suggest that climatic factors controlling vertical mixing may be capable of triggering hypoxia under preanthropogenic P loading, despite a recent study indicating otherwise (Meier et al., 2019).

### 3.7. Implications for Future Management of the Baltic Sea

Under the BSAP (HELCOM, 2007, 2021), riparian countries agreed to limit external P loading to the Baltic in order to reduce the occurrence of hypoxia and cyanobacterial blooms in the future. As part of the plan, maximum allowable inputs of P to the whole Baltic Sea were set at 21,716 tonnes P/year (about 18,500 tonnes P/year in the region covered by the model). Our findings suggest that such targets, if met, will improve water quality but are unlikely to force the system to a fully oxic state (Figure 3e). Rather, deep water oxygen concentrations are likely to remain close to the hypoxia threshold of 63  $\mu\text{mol/L}$ . This finding is consistent with existing simulations from more complex coupled physical-biogeochemical models for the Baltic Sea (e.g., Meier et al., 2018). To achieve a transition beyond hypoxia to quasipermanent oxic conditions would require a further load reduction and our model shows that this will likely proceed via an interval during which the system is vulnerable to multidecadal oscillations in hypoxia, creating a challenge for assessments of the success of management efforts. These results highlight the need to include more details of Fe-P interactions in models used for management decisions in the Baltic Sea (e.g., BALTSEM; Savchuk et al., 2012). We suggest that the extent to which oscillations may impact upon the actual recovery trajectory in the coming centuries will depend on the speed with which loading reduction targets are achieved. More rapid reductions are more likely to starve the system of P and therefore to promote a more linear recovery trajectory.

## 4. Conclusions

Our study shows that feedbacks in the coupled cycling of Fe and P drove multidecadal oscillations in the intensity of oxygen depletion during past hypoxic intervals in the 8,000 years history of the Baltic Sea. In-phase oscillatory profiles of Mo/Al, Br/P and Fe/Al in sediments suggest large-scale synchronicity in deep water hypoxia, P regeneration and Fe shuttling, respectively. A box model of coupled Fe and P cycling suggests that internal multidecadal instability in hypoxia intensity is controlled by nonlinear recharging and discharging of the sedimentary Fe-P reservoir in deeper areas. Crucially, our model simulations show that oscillatory behavior is sensitive to external P loading. While low external loads favor stable oxic conditions, and high loads sustain stable hypoxia, intermediate loads lead to instability causing oscillatory behavior on multidecadal timescales. This observation

has implications for predicting the trajectory of recovery from modern hypoxia in the Baltic Sea, since external P loading is expected to decline in accordance with the BSAP. Variable external climate forcing may influence the frequency and amplitude of oscillations through its impact on vertical water mass exchange.

### Data Availability Statement

Open Research: The raw LA-ICP-MS data and time series analysis outputs are available free of charge (CC-BY 4.0) on Zenodo at <https://doi.org/10.5281/zenodo.5222925> (<https://doi.org/10.5281/zenodo.5222925>). The box model (v1.0.2) used for investigating the mechanisms of oscillations is available (MIT Licence: Copyright Bo Gustafsson) at <https://doi.org/10.5281/zenodo.5235401> (<https://doi.org/10.5281/zenodo.5235401>). The model code was developed using Xcode IDE (<https://developer.apple.com/xcode/>) in Swift language.

### Acknowledgments

This work was supported by European Research Council (ERC) Starting Grant (Consolidator Level) 278364 and Netherlands Organisation for Scientific Research (NWO) Vici grant 865.13.005 to Caroline Slomp, Academy of Finland Research Fellowship 317684 to Tom Jilbert, the EU BONUS-HYPER project, Baltic Bridge collaboration between University of Helsinki and Stockholm University and the program of the Netherlands Earth System Science Center (NESSC), financially supported by the Ministry of Education, Culture and Science (OCW).

### References

- Börgel, F., Frauen, C., Neumann, T., & Meier, H. E. M. (2020). The Atlantic Multidecadal Oscillation controls the impact of the North Atlantic Oscillation on North European climate. *Environmental Research Letters*, 15(10), 104025. <https://doi.org/10.1088/1748-9326/aba925>
- Carstensen, J., Andersen, J. H., Gustafsson, B. G., & Conley, D. J. (2014). Deoxygenation of the Baltic Sea during the last century. *Proceedings of the National Academy of Sciences*, 111(15), 5628–5633. <https://doi.org/10.1073/pnas.1323156111>
- Conley, D. J., Björck, S., Bonsdorff, E., Carstensen, J., Destouni, G., Gustafsson, B. G., et al. (2009). Hypoxia-related processes in the Baltic Sea. *Environmental Science & Technology*, 43(10), 3412–3420. <https://doi.org/10.1021/es802762a>
- Eckert, S., Brumsack, H.-J., Severmann, S., Schnetger, B., März, C., & Fröllje, H. (2013). Establishment of euxinic conditions in the Holocene Black Sea. *Geology*, 41(4), 431–434. <https://doi.org/10.1130/G33826.1>
- Gustafsson, B. G., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B., et al. (2012). Reconstructing the development of Baltic Sea eutrophication 1850–2006. *Ambio*, 41(6), 534–548. <https://doi.org/10.1007/s13280-012-0318-x>
- HELCOM. (2007). *HELCOM Baltic Sea action plan. Adopted in Krakow*.
- HELCOM. (2021). *Baltic Sea Action Plan, 2021 update. Adopted in Lübeck*.
- Hennekam, R., Jilbert, T., Mason, P. R. D., de Lange, G. J., & Reichert, G.-J. (2015). High-resolution line-scan analysis of resin-embedded sediments using laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS). *Chemical Geology*, 403, 42–51. <https://doi.org/10.1016/j.chemgeo.2015.03.004>
- Ingall, E. D., Bustin, R., & Van Cappellen, P. (1993). Influence of water column anoxia on the burial and preservation of carbon and phosphorus in marine shales. *Geochimica et Cosmochimica Acta*, 57(2), 303–316. [https://doi.org/10.1016/0016-7037\(93\)90433-W](https://doi.org/10.1016/0016-7037(93)90433-W)
- Jilbert, T., Conley, D. J., Gustafsson, B. G., Funkey, C. P., & Slomp, C. P. (2015). Glacio-isostatic control on hypoxia in a high-latitude shelf basin. *Geology*, 43(5), 427–430. <https://doi.org/10.1130/G36454.1>
- Jilbert, T., de Lange, G. J., & Reichert, G.-J. (2008). Fluid displacive resin embedding of laminated sediments: Preserving trace metals for high-resolution paleoclimate investigations. *Limnology and Oceanography: Methods*, 6(1), 16–22. <https://doi.org/10.4319/lom.2008.6.16>
- Jilbert, T., Slomp, C. P., Gustafsson, B. G., & Boer, W. (2011). Beyond the Fe-P-redox connection: Preferential regeneration of phosphorus from organic matter as a key control on Baltic Sea nutrient cycles. *Biogeosciences*, 8(6), 1699–1720. <https://doi.org/10.5194/bg-8-1699-2011>
- Jilbert, T., & Slomp, C. P. (2013). Rapid high-amplitude variability in Baltic Sea hypoxia during the Holocene. *Geology*, 41(11), 1183–1186. <https://doi.org/10.1130/G34804.1>
- Jokinen, S. A., Virtasalo, J. J., Jilbert, T., Kaiser, J., Dellwig, O., Arz, H. W., et al. (2018). A 1500-year multiproxy record of coastal hypoxia from the northern Baltic Sea indicates unprecedented deoxygenation over the 20th century. *Biogeosciences*, 15(13), 3975–4001. <https://doi.org/10.5194/bg-15-3975-2018>
- Knudsen, M. F., Seidenkrantz, M.-S., Jacobsen, B. H., & Kuijpers, A. (2011). Tracking the Atlantic Multidecadal Oscillation through the last 8,000 years. *Nature Communications*, 2(1), 1–8. <https://doi.org/10.1038/ncomms1186>
- Lefort, S., Gratton, Y., Mucci, A., Dadou, I., & Gilbert, D. (2012). Hypoxia in the Lower St. Lawrence Estuary: How physics controls spatial patterns. *Journal of Geophysical Research: Oceans*, 117(C7). <https://doi.org/10.1029/2011JC007751>
- Lenz, C., Jilbert, T., Conley, D. J., & Slomp, C. P. (2015). Hypoxia-driven variations in iron and manganese shuttling in the Baltic Sea over the past 8 kyr. *Geochemistry, Geophysics, Geosystems*, 16(10), 3754–3766. <https://doi.org/10.1002/2015GC005960>
- Lougheed, B. C., Snowball, I., Moros, M., Kabel, K., Muscheler, R., Virtasalo, J. J., & Wacker, L. (2012). Using an independent geochronology based on palaeomagnetic secular variation (PSV) and atmospheric Pb deposition to date Baltic Sea sediments and infer 14C reservoir age. *Quaternary Science Reviews*, 42, 43–58. <https://doi.org/10.1016/j.quascirev.2012.03.013>
- Lyons, T. W., & Severmann, S. (2006). A critical look at iron paleoredox proxies: New insights from modern euxinic marine basins. *Geochimica et Cosmochimica Acta*, 70(23), 5698–5722. <https://doi.org/10.1016/j.gca.2006.08.021>
- Meier, H. E. M., Edman, M. K., Eilola, K. J., Placke, M., Neumann, T., Andersson, H. C., et al. (2018). Assessment of eutrophication abatement scenarios for the Baltic Sea by multi-model ensemble simulations. *Frontiers in Marine Science*, 5, 440. <https://doi.org/10.3389/fmars.2018.00440>
- Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., et al. (2019). Disentangling the impact of nutrient load and climate changes on Baltic Sea hypoxia and eutrophication since 1850. *Climate Dynamics*, 53(1), 1145–1166. <https://doi.org/10.1007/s00382-018-4296-y>
- Mortimer, C. H. (1941). The exchange of dissolved substances between mud and water in lakes. *Journal of Ecology*, 29(2), 280–329. <https://doi.org/10.2307/2256395>
- Ning, W., Ghosh, A., Jilbert, T., Slomp, C. P., Khan, M., Nyberg, J., et al. (2016). Evolving coastal character of a Baltic Sea inlet during the Holocene shoreline regression: Impact on coastal zone hypoxia. *Journal of Paleolimnology*, 55(4), 319–338. <https://doi.org/10.1007/s10933-016-9882-6>
- Norbäck Ivarsson, L., Andrén, T., Moros, M., Andersen, T. J., Lönn, M., & Andrén, E. (2019). Baltic sea coastal eutrophication in a thousand year perspective. *Frontiers in Environmental Science*, 7, 88. <https://doi.org/10.3389/fenvs.2019.00088>

- Paillard, D., Labeyrie, L., & Yiou, P. (1996). Macintosh program performs time-series analysis. *Eos, Transactions American Geophysical Union*, 77(39), 379. <https://doi.org/10.1029/96EO00259>
- Rabalais, N. N., Diaz, R. J., Levin, L. A., Turner, R. E., Gilbert, D., & Zhang, J. (2010). Dynamics and distribution of natural and human-caused hypoxia. *Biogeosciences*, 7(2), 585–619. <https://doi.org/10.5194/bg-7-585-2010>
- Reed, D. C., Slomp, C. P., & Gustafsson, B. G. (2011). Sedimentary phosphorus dynamics and the evolution of bottom-water hypoxia: A coupled benthic–pelagic model of a coastal system. *Limnology & Oceanography*, 56(3), 1075–1092. <https://doi.org/10.4319/lo.2011.56.3.1075>
- Savchuk, O. P., Gustafsson, B. G., & Müller-Karulis, B. (2012). BALTSEM: A marine model for decision support within the Baltic Sea Region. *Baltic nest institute technical report*, 7 (p. 55).
- Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., et al. (2007). *Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria blooms and complicate management in the Baltic Sea* (pp. 186–194). *Ambio*. [https://doi.org/10.1579/0044-7447\(2007\)36\[186:iefenc\]2.0.co;2](https://doi.org/10.1579/0044-7447(2007)36[186:iefenc]2.0.co;2)
- Van Cappellen, P., & Ingall, E. D. (1994). Benthic phosphorus regeneration, net primary production, and ocean anoxia: A model of the coupled marine biogeochemical cycles of carbon and phosphorus. *Paleoceanography*, 9(5), 677–692. <https://doi.org/10.1029/94PA01455>
- Wang, J., Yang, B., Ljungqvist, F. C., Luterbacher, J., Osborn, T. J., Briffa, K. R., & Zorita, E. (2017). Internal and external forcing of multidecadal Atlantic climate variability over the past 1,200 years. *Nature Geoscience*, 10(7), 512–517. <https://doi.org/10.1038/NGEO2962>
- Warden, L., Moros, M., Neumann, T., Shennan, S., Timpson, A., Manning, K., et al. (2017). Climate induced human demographic and cultural change in northern Europe during the mid-Holocene. *Scientific Reports*, 7(1), 1–11. <https://doi.org/10.1038/s41598-017-14353-5>
- Ziegler, M., Jilbert, T., de Lange, G. J., Lourens, L. J., & Reichert, G. J. (2008). Bromine counts from XRF scanning as an estimate of the marine organic carbon content of sediment cores. *Geochemistry, Geophysics, Geosystems*, 9(5). <https://doi.org/10.1029/2007GC001932>
- Zillén, L., & Conley, D. J. (2010). Hypoxia and cyanobacteria blooms are they really natural features of the late Holocene history of the Baltic Sea? *Biogeosciences*, 7(8), 2567–2580. <https://doi.org/10.5194/bg-7-2567-2010>
- Zillén, L., Conley, D. J., Andrén, T., Andrén, E., & Björck, S. (2008). Past occurrences of hypoxia in the Baltic Sea and the role of climate variability, environmental change and human impact. *Earth-Science Reviews*, 91(1–4), 77–92. <https://doi.org/10.1016/j.earscirev.2008.10.001>

## References From the Supporting Information

- Al-Hamdani, Z., Reker, J., Alanen, U., Andersen, J. H., Bendtsen, J., Bergström, U., et al. (2007). Towards benthic marine landscapes in the Baltic Sea. *BALANCE interim report*, 10.
- Almroth-Rosell, E., Eilola, K., Hordoir, R., Meier, H. E. M., & Hall, P. O. J. (2011). Transport of fresh and resuspended particulate organic material in the Baltic Sea: A model study. *Journal of Marine Systems*, 87(1), 1–12. <https://doi.org/10.1016/j.jmarsys.2011.02.005>
- Almroth-Rosell, E., Eilola, K., Kuznetsov, I., Hall, P. O., & Meier, H. M. (2015). A new approach to model oxygen dependent benthic phosphate fluxes in the Baltic Sea. *Journal of Marine Systems*, 144, 127–141. <https://doi.org/10.1016/j.jmarsys.2014.11.007>
- Andersson, L. (1996). Trends in nutrients and oxygen concentrations in the Skagerrak-Kattegat. *Journal of Sea Research*, 35(1–3), 63–71. [https://doi.org/10.1016/S1385-1101\(96\)90735-2](https://doi.org/10.1016/S1385-1101(96)90735-2)
- Boudreau, B. P. (1997). *Diagenetic models and their implementation* (p. 414). Springer.
- Conley, D. J., Humborg, C., Rahm, L., Savchuk, O. P., & Wulff, F. (2002). Hypoxia in the Baltic Sea and basin-scale changes in phosphorus biogeochemistry. *Environmental Science & Technology*, 36(24), 5315–5320. <https://doi.org/10.1021/es025763w>
- Eilola, K., Meier, H. E. M., & Almroth, E. (2009). On the dynamics of oxygen, phosphorus and cyanobacteria in the Baltic Sea; A model study. *Journal of Marine Systems*, 75(1–2), 163–184. <https://doi.org/10.1016/j.jmarsys.2008.08.009>
- Gustafsson, B. G. (2001). Quantification of water, salt, oxygen and nutrient exchange of the Baltic Sea from observations in the Arkona Basin. *Continental Shelf Research*, 21(13–14), 1485–1500. [https://doi.org/10.1016/S0278-4343\(01\)00014-0](https://doi.org/10.1016/S0278-4343(01)00014-0)
- Gustafsson, B. G., & Stigebrandt, A. (2007). Dynamics of nutrients and oxygen/hydrogen sulfide in the Baltic Sea deep water. *Journal of Geophysical Research: Biogeosciences*, 112(G2). <https://doi.org/10.1029/2006JG000304>
- Gustafsson, B. G., & Westman, P. (2002). On the causes for salinity variations in the Baltic Sea during the last 8500 years. *Paleoceanography*, 17(3), 12–11214. <https://doi.org/10.1029/2000PA000572>
- Gustafsson, E., Savchuk, O. P., Gustafsson, B. G., & Müller-Karulis, B. (2017). Key processes in the coupled carbon, nitrogen, and phosphorus cycling of the Baltic Sea. *Biogeochemistry*, 134(3), 301–317. <https://doi.org/10.1007/s10533-017-0361-6>
- Gustafsson, Ö., Gelting, J., Andersson, P., Larsson, U., & Roos, P. (2013). An assessment of upper ocean carbon and nitrogen export fluxes on the boreal continental shelf: A 3-year study in the open Baltic Sea comparing sediment traps, 234Th proxy, nutrient, and oxygen budgets. *Limnology and Oceanography: Methods*, 11(9), 495–510. <https://doi.org/10.4319/om.2013.11.495>
- Hermans, M., Lenstra, W. K., van Helmond, N. A. G. M., Behrends, T., Egger, M., Séguret, M. J. M., et al. (2019). Impact of natural re-oxygenation on the sediment dynamics of manganese, iron and phosphorus in a euxinic Baltic Sea basin. *Geochimica et Cosmochimica Acta*, 246, 174–196. <https://doi.org/10.1016/j.gca.2018.11.033>
- Jönsson, A., Danielsson, Å., & Rahm, L. (2005). Bottom type distribution based on wave friction velocity in the Baltic Sea. *Continental Shelf Research*, 25(3), 419–435. <https://doi.org/10.1016/j.csr.2004.09.011>
- Meier, H. M. (2005). Modeling the age of Baltic Seawater masses: Quantification and steady state sensitivity experiments. *Journal of Geophysical Research: Oceans*, 110(C2). <https://doi.org/10.1029/2004JC002607>
- Mort, H. P., Slomp, C. P., Gustafsson, B. G., & Andersen, T. J. (2010). Phosphorus recycling and burial in Baltic Sea sediments with contrasting redox conditions. *Geochimica et Cosmochimica Acta*, 74(4), 1350–1362. <https://doi.org/10.1016/j.gca.2009.11.016>
- Pers, C., & Rahm, L. (2000). Changes in apparent oxygen removal in the Baltic proper deep water. *Journal of Marine Systems*, 25(3–4), 421–429. [https://doi.org/10.1016/S0924-7963\(00\)00031-2](https://doi.org/10.1016/S0924-7963(00)00031-2)
- Press, W. H., Teukolsky, S. A., Flannery, B. P., & Vetterling, W. T. (1992). *Numerical recipes in Fortran 77* (2nd ed.). Cambridge University Press.
- Reed, D. C., Gustafsson, B. G., & Slomp, C. P. (2016). Shelf-to-basin iron shuttling enhances vivianite formation in deep Baltic Sea sediments. *Earth and Planetary Science Letters*, 434, 241–251. <https://doi.org/10.1016/j.epsl.2015.11.033>
- Savchuk, O. P. (2002). Nutrient biogeochemical cycles in the Gulf of Riga: Scaling up field studies with a mathematical model. *Journal of Marine Systems*, 32(4), 253–280. [https://doi.org/10.1016/S0924-7963\(02\)00039-8](https://doi.org/10.1016/S0924-7963(02)00039-8)
- Savchuk, O. P., & Wulff, F. (2009). Long-term modeling of large-scale nutrient cycles in the entire Baltic Sea. In *Eutrophication in coastal ecosystems* (pp. 209–224). Springer. [https://doi.org/10.1007/978-90-481-3385-7\\_18](https://doi.org/10.1007/978-90-481-3385-7_18)
- Savchuk, O. P., Wulff, F., Hille, S., Humborg, C., & Pollehne, F. (2008). The Baltic Sea a century ago: A reconstruction from model simulations, verified by observations. *Journal of Marine Systems*, 74(1–2), 485–494. <https://doi.org/10.1016/j.jmarsys.2008.03.008>

- Seifert, T., Tauber, F., & Kayser, B. (2001). *A high resolution spherical grid topography of the Baltic Sea—revised edition* (pp. 25–29). Baltic Sea Science Congress.
- Stigebrandt, A. (1991). Computations of oxygen fluxes through the sea surface and the net production of organic matter with application to the Baltic and adjacent seas. *Limnology & Oceanography*, *36*(3), 444–454. <https://doi.org/10.4319/lo.1991.36.3.0444>
- Stigebrandt, A., & Gustafsson, B. G. (2003). Response of the Baltic Sea to climate change—Theory and observations. *Journal of Sea Research*, *49*(4), 243–256. [https://doi.org/10.1016/S1385-1101\(03\)00021-2](https://doi.org/10.1016/S1385-1101(03)00021-2)
- Tamelander, T., Spilling, K., & Winder, M. (2017). Organic matter export to the seafloor in the Baltic Sea: Drivers of change and future projections. *Ambio*, *46*(8), 842–851. <https://doi.org/10.1007/s13280-017-0930-x>
- Vollenweider, R. A. (1969). Possibilities and limits of elementary models concerning budget of substances in lakes. *Archiv für Hydrobiologie*, *66*(1), 1–36.