

On the Gel'fand-Calderón inverse problem in two dimensions

Eemeli Blåsten

Academic dissertation

*To be presented, with the permission of the
Faculty of Science of the University of Helsinki, for public examination
in Auditorium CK112, Exactum, on April 30, 2013, at 10 o'clock.*

Helsinki 2013

University of Helsinki
Faculty of Science
Department of Mathematics and Statistics
Supervisor Lassi Päivärinta

Supervisor: Professor Lassi Päivärinta
Department of Mathematics and Statistics
University of Helsinki
Helsinki, Finland

Pre-examiners: Professor Giovanni Alessandrini
Dipartimento di Matematica e Informatica
Università di Trieste
Trieste, Italia

Professor Alberto Ruiz
Departamento de Matemáticas
Universidad Autónoma de Madrid
Madrid, España

Opponent: Professor Yaroslav Kurylev
Department of Mathematics
University College London
London, United Kingdom

ISBN 978-952-10-8698-4 (paperback)

ISBN 978-952-10-8699-1 (PDF)

Unigrafia

E-thesis (<http://www.e-thesis.fi>)

Helsinki 2013

Acknowledgments

First and foremost I would like to thank my PhD advisor Lassi Päivärinta for introducing me to Bukhgeim's original article and having a good enough intuition to see that I was going the right way. I wish to thank Alberto Ruiz and Giovanni Alessandrini for their precious time spent reading the manuscript very carefully and giving excellent suggestions.

I owe my officemate Esa Vesalainen and numerous other colleagues a lot for the many invaluable discussions during the past years. Without them, I would be stuck in my own way of thinking. A special thanks to Pedro Caro for suggesting me to use boundary data instead of assuming well-posedness.

I also want to thank my wife Wang Ruiling for her patience and our wonderful time together. She said that no matter what problems I will encounter, I would find a way around them.

My special thanks to the Finnish Inverse Problems Society for arranging the yearly Inverse Problems Days. They are a wonderful workshop to meet other Finnish researchers and appreciate how close the community is. It made me very happy about starting graduate studies in this field.

Lastly, I thank the Academy of Finland for its financial support through the Finnish Center of Excellence in Inverse Problems Research.

Contents

1	Introduction	1
1.1	Abstract	1
1.2	History and related work	1
1.3	The main result and sketch of the proof	4
2	Function spaces	6
2.1	Banach-valued Lorentz spaces	6
2.2	Interpolation of Lorentz spaces	10
2.3	Lorentz-Sobolev spaces	16
2.4	Lorentz-Bessel potential spaces	18
2.5	The Cauchy operator	21
3	Using the stationary phase method	24
3.1	The main term	24
3.2	Handling the error term	25
4	Bukhgeim type solutions	28
4.1	A Carleman estimate	28
4.2	Bukhgeim’s oscillating solutions	34
5	The problem setting	36
5.1	Hadamard’s criteria and the DN operator	36
5.2	Cauchy data	38
5.3	Uniqueness and stability for the inverse problem	40
6	Future work	45
6.1	Function space properties of $H^{s,(p,q)}(\Omega)$ and $W^{s,(p,q)}(\Omega)$	45
6.2	Doing it in $W^{s,p}$	46
6.3	Non-compactly supported potentials	46
6.4	No smoothness	48
6.5	A reconstruction formula	49
7	Calculations	51
	References	59
	Index of Notation	63
	Index of Subjects	64

1 Introduction

1.1 Abstract

We prove uniqueness and stability for the inverse boundary value problem of the 2D Schrödinger equation. We assume only that the potentials are in $H^{s,(2,1)}(\Omega)$, $s > 0$, which is slightly smaller than the Sobolev space $H^{s,2}(\Omega)$. The thesis consists of two parts.

In the first part, we define the spaces $H^{s,(p,q)}$ of distributions whose fractional derivatives are in the Lorentz space $L^{(p,q)}$. We prove the embedding $H^{1,(n,1)} \hookrightarrow C^0$ and an interpolation identity.

The inverse problem is considered in the second part of the thesis. We prove a new Carleman estimate for $\bar{\partial}$. This estimate has a decay rate of $\tau^{-1} \ln \tau$. After that we use Bukhgeim's oscillating exponential solutions, Alessandrini's identity and stationary phase to get information about the difference of the potentials from the difference of the Cauchy data.

1.2 History and related work

This short survey of results concerning inverse boundary value problems for the conductivity and Schrödinger equations is based mostly on introductions in [5] and [36]. We mention also a few papers from recent years that we have personally heard of. The majority of the results cited below were proven for the conductivity equation or the Schrödinger equation having a potential coming from a related conductivity equation.

The inverse problem of the Schrödinger equation, also known as the Gel'fand or Gel'fand-Calderón inverse problem (see [19]), is the following one:

$$\text{Given } C_q = \{(u|_{\partial\Omega}, \partial_\nu u|_{\partial\Omega}) \mid \Delta u + qu = 0\} \text{ deduce } q. \quad (1.1)$$

In other words, given measurements of the solutions u only on the boundary $\partial\Omega$ of an object or area Ω , what can we say about the potential q inside of Ω ? The Schrödinger equation can model acoustic, electromagnetic and quantum waves. Hence this inverse problem models inverse scattering of time harmonic waves in these situations.

One of the important early papers on inverse boundary value problems is by Calderón [11]. He considered an isotropic body Ω from which one would like to deduce the electrical conductivity γ by doing electrical measurements on the boundary. If we keep the voltage u fixed as f on the boundary, then

the stationary state of u can be modeled by the boundary value problem

$$\begin{aligned}\nabla \cdot (\gamma \nabla u) &= 0, & \Omega \\ u &= f, & \partial\Omega.\end{aligned}\tag{1.2}$$

The weighted normal derivative $\gamma \partial_\nu u$ is the current flux going out of Ω . Calderón asked whether knowing the boundary measurements, or Dirichlet-Neumann map $\Lambda_\gamma : f \mapsto \gamma \partial_\nu u|_{\partial\Omega}$, is enough to determine the conductivity γ inside the whole domain Ω . This is called the *Calderón problem*. He showed the injectivity of a linearized problem near $\gamma \equiv 1$.

The inverse problem for the conductivity equation can be reduced to that of the Schrödinger equation. To transform the conductivity equation into the equation $\Delta v + qv = 0$, it is enough to do the change of variables $u = \gamma^{-\frac{1}{2}}v$, $q = -\gamma^{-\frac{1}{2}}\Delta\gamma^{\frac{1}{2}}$. The Dirichlet-Neumann map for the new equation can be recovered from the boundary data of the old one: $\Lambda_q = \gamma^{-\frac{1}{2}}(\Lambda_\gamma + \frac{1}{2}\frac{\partial\gamma}{\partial\nu})\gamma^{-\frac{1}{2}}$.

Sylvester and Uhlmann solved the problem in dimensions n at least three for smooth conductivities bounded away from zero [46]. They constructed *complex geometric optics solutions*, that is, solutions of the form

$$u_j = e^{x \cdot \zeta_j} \left(1 + O\left(\frac{1}{|\zeta_j|}\right)\right),\tag{1.3}$$

where the complex vectors ζ_j satisfy

$$\begin{aligned}\zeta_1 &= i(k + m) + l, \\ \zeta_2 &= i(k - m) - l,\end{aligned}\tag{1.4}$$

where $l, k, m \in \mathbb{R}^n$ are perpendicular vectors satisfying $|l|^2 = |k|^2 + |m|^2$. Using a well-known orthogonality relation for the potentials q_1 and q_2 , called the Alessandrini identity [2], they got

$$0 = \int (q_1 - q_2) u_1 u_2 dx = \int (q_1 - q_2) e^{2ix \cdot k} \left(1 + O\left(\frac{1}{|m|}\right)\right) dx,\tag{1.5}$$

and after taking $|m| \rightarrow \infty$ they saw that the Fourier transforms of q_1 and q_2 are the same, so the potentials are so too. Note that the only part that requires $n \geq 3$ in this solution is the existence of the three vectors l, k, m .

Some papers solve the Calderón problem in dimension two with various assumptions. Namely Kohn and Vogelius [29] [30], Alessandrini [2], Nachman [36] and finally Astala and Päivärinta [5]. The first three of these require the conductivity to be piecewise analytic. Nachman required two derivatives to convert the conductivity equation into the Schrödinger equation. The paper of Astala and Päivärinta solved Calderón's problem most generally:

there were no requirements on the smoothness of the conductivity. It just had to be bounded away from zero and infinity, which is physically realistic.

There are also some results for the inverse boundary value problem of the Schrödinger equation whose potential is not assumed to be of the conductivity type. Jerison and Kenig proved, according to [12], that if $q \in L^p(\Omega)$ with $p > \frac{n}{2}$, $n \geq 3$, then the Dirichlet-Neumann map Λ_q determines the potential q uniquely. The case $n = 2$ was open until the paper of Bukhgeim. In [3], he introduced new kinds of solutions to the Schrödinger equation, which allow the use of stationary phase. This led to an elegant solution of this long standing open problem. There is a point in the argument that requires differentiability of the potentials. Imanuvilov and Yamamoto published the paper [27] in arXiv after the writing of this thesis. They seem to have fixed that problem and hence proven uniqueness for $q \in L^p(\Omega)$, $p > 2$.

Some more recent results in two dimensions have concerned partial data, stability and reducing smoothness requirements for the conductivities and potentials. Notable results of partial data include Imanuvilov, Uhlmann, Yamamoto [26] and Guillarmou and Tzou [23]. In the first paper the authors consider the Schrödinger equation in a plane domain and in the second one on a Riemann surface with boundary. The results of both papers state that knowing the Cauchy data on any open subset on the boundary determines the potential uniquely if it is smooth enough.

Stability seemed to be proven first for the inverse problem of the conductivity equation. Liu [31] showed it for potentials of the conductivity type. Barceló, Faraco and Ruíz [6] showed stability for Hölder continuous conductivities. Clop, Faraco and Ruíz generalized it to $W^{\alpha,p}$, $\alpha > 0$, in [13]. For the Schrödinger equation, there's the result of Novikov and Santacesaria for C^2 potentials in [39].

Lastly, we cite very briefly some reconstruction methods. This paragraph is certainly very incomplete as reconstruction was not the focus of the thesis. Nachman gave the first result for the conductivity equation for $n \geq 3$ in [35] and later for $n = 2$ in [36]. In the recent paper [4], the authors show a numerical reconstruction method for piecewise smooth conductivities in 2D. For a more in-depth survey, see the introduction in that same paper. The case of the Schrödinger equation in the plane seems to be more elusive. Bukhgeim mentioned a reconstruction formula at the end of [3], but as far as we know, there are no published numerical methods for reconstructing the potential in 2D. There is a reconstruction formula using only the boundary data explicitly in [40] though.

1.3 The main result and sketch of the proof

We will give a top-down sketch for proving uniqueness and stability. Before that, we will describe the inverse problem. Let q_1 and q_2 be two potentials for the Schrödinger equations $(\Delta + q_j)u = 0$. We define the *boundary data* C_{q_j} as the collection of pairs $(u|_{\Omega}, \nu \cdot \partial_{\nu} u|_{\Omega})$ of boundary values and boundary derivatives of all solutions u . If we assume that the operators $\Delta + q_j$ are well posed in Hadamard's sense, then the two sets of boundary data become the *Dirichlet-Neumann maps* $\Lambda_{q_j} : u|_{\Omega} \mapsto \nabla u|_{\Omega}$, where $\Delta u + q_j u = 0$. The problem is, what can we tell about $q_1 - q_2$ if we know C_{q_1}, C_{q_2} ? We will show the following:

Theorem. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $M > 0$ and $0 < s < \frac{1}{2}$. Then there is a positive real number C such that if $\|q_j\|_{s,(2,1)} \leq M$ then*

$$\|q_1 - q_2\|_{L^{(2,\infty)}(\Omega)} \leq C (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4}. \quad (1.6)$$

Here $q_j \in H^{s,(2,1)}(\Omega)$, which can be considered as a slightly smaller space than $H^{s,2}(\Omega)$, and $d(C_{q_1}, C_{q_2})$ is the distance between C_{q_1} and C_{q_2} in a certain sense. It is basically

$$\sup \left\{ \left| \int_{\Omega} u_1(q_1 - q_2)u_2 \right| \mid \Delta u_j + q_j u_j = 0, u_j \in W^{1,2}(\Omega), \|u_j\| = 1 \right\}, \quad (1.7)$$

but, using Green's formula, the integral over Ω can be transformed to

$$\dots = \int_{\Omega} u_2 \Delta u_1 - u_1 \Delta u_2 dm = \int_{\partial\Omega} u_2 \nu \cdot \nabla u_1 - u_1 \nu \cdot \nabla u_2 d\sigma, \quad (1.8)$$

which are measurements done on the boundary. Hence, our goal is to estimate $\|q_1 - q_2\|$ by expressions involving $\int_{\Omega} u_1(q_1 - q_2)u_2$. This is achieved by choosing special solutions u_1, u_2 , which allow the use of a stationary phase method. Another powerful tool we will use is Carleman estimates. They will take care of the error term, which comes from the fact that the solutions u_1 and u_2 are not analytic.

The top-down idea starts as follows. Stationary phase arguments show that

$$\|q_1 - q_2\| \leftarrow \left\| \frac{2\tau}{\pi} e^{i\tau(z^2 + \bar{z}^2)} * (q_1 - q_2) \right\| \quad (1.9)$$

as $\tau \rightarrow \infty$. We will show that there are solutions such that $u_1 u_2 \rightarrow e^{i\tau(z^2 + \bar{z}^2)}$. This construction was first shown by Bukhgeim [3]. Those solutions will in

fact look like $u_1 = e^{i\tau(z-z_0)^2} f_1$, $u_2 = e^{i\tau(\bar{z}-\bar{z}_0)^2} f_2$, where z_0 is the variable outside the convolution, and $f_j \rightarrow 1$. Hence we get

$$\begin{aligned} \|q_1 - q_2\| &\leq \left\| q_1 - q_2 - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right\| + \left\| \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right\| \\ &\leq \left\| q_1 - q_2 - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right\| + \left\| \frac{2\tau}{\pi} \int_{\Omega} u^{(1)}(q_1 - q_2) u^{(2)} dm \right\| \\ &\quad + \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} (q_1 - q_2) (1 - f_1 f_2) dm \right\|, \end{aligned} \quad (1.10)$$

where $R = (z - z_0)^2 + (\bar{z} - \bar{z}_0)^2$.

The first term in the equation of the above paragraph can be estimated by $\tau^{-s/2} \|q_1 - q_2\|_{H^{s,2}}$ because of stationary phase. The second one is easy because of the definition of $d(C_{q_1}, C_{q_2})$. It has the upper bound

$$d(C_{q_1}, C_{q_2}) \|u_1\| \|u_2\| \sim e^{c\tau} d(C_{q_1}, C_{q_2}) \quad (1.11)$$

because of the form of the solutions. The last term is the hardest. By using a suitable cut-off function, we can estimate it above by

$$\tau^{1-s/3} \|q_1 - q_2\|_{H^{s,(2,1)}(\Omega)} \|1 - f_1 f_2\|_{H^{s,(2,\infty)}(\Omega)}. \quad (1.12)$$

We need to show that $\|1 - f_1 f_2\|_s = o(\tau^{s/3-1})$ as $\tau \rightarrow \infty$ to get uniqueness. This is the part that requires new results. It all boils down to Carleman estimates. Section 4.1 with theorem 4.1.1 and corollaries 4.1.5 and 4.1.10 are all about proving them. The new estimates are

$$\begin{aligned} \|r\|_{H^{(2,\infty)}} &\leq C_{\Omega} \tau^{-1} (1 + \ln \tau) \left\| e^{i\tau(\bar{z}-\bar{z}_0)^2} \bar{\partial} e^{-i\tau(\bar{z}-\bar{z}_0)^2} r \right\|_{H^{1,(2,1)}} \\ \|r\|_{C^0} &\leq C_{\Omega} \tau^{-1/3} \left\| e^{i\tau(\bar{z}-\bar{z}_0)^2} \bar{\partial} e^{-i\tau(\bar{z}-\bar{z}_0)^2} r \right\|_{H^{1,(2,1)}} \\ \|r\|_{H^{s,(2,\infty)}} &\leq C_{\Omega} \tau^{-1} (1 + \ln \tau) \left\| e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r \right\|_{H^{s,(2,1)}} \\ \|r\|_{M^s} &\leq C_{\Omega} \tau^{-1/3} \left\| e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r \right\|_{H^{s,(2,1)}} \end{aligned} \quad (1.13)$$

where $H^{s,(p,q)}$ is a slight generalization of $H^{s,p}$, and M^s is a space whose functions have smoothness s and can be embedded into C^0 . We will prove the estimates in the integral form, that is, having the Cauchy operator on the left-hand side. Choosing $r = f_j - 1$ implies that $\|1 - f_1 f_2\|_s = O(\tau^{-1} \ln \tau)$. Hence, whenever $s > 0$, the error term (1.12) tends to zero as τ grows.

Combining all the upper bounds, we have

$$\|q_1 - q_2\| \leq \tau^{-\beta s} + e^{c\tau} d(C_{q_1}, C_{q_2}) \quad (1.14)$$

with some $\beta, c > 0$. A suitable choice of τ implies the claim.

2 Function spaces

2.1 Banach-valued Lorentz spaces

Definition 2.1.1. Let A be a vector space and $X \subset \mathbb{R}^n$ measurable. Then the mapping $f : X \rightarrow A$ is a *simple function* if

$$f(x) = \sum_{k=0}^N a_k \chi_{E_k}(x) \quad (2.1)$$

for all $x \in X$ and some $N \in \mathbb{N}$, $a_k \in A$ and disjoint measurable $E_k \subset \mathbb{R}^n$. We use the Lebesgue measure in \mathbb{R}^n where not specified explicitly.

Definition 2.1.2. Let A be a Banach space and $X \subset \mathbb{R}^n$ measurable. A function $X \rightarrow A$ is *strongly measurable* if there is a sequence of simple functions $f_m : X \rightarrow A$ such that

$$f(x) = \lim_{m \rightarrow \infty} f_m(x) \quad (2.2)$$

for almost all $x \in X$.

Definition 2.1.3. Let A be a Banach space, $\Omega \subset \mathbb{R}^n$ open and $f : \Omega \rightarrow A$ strongly measurable. Then the *distribution function of f* , $\lambda \mapsto m(f, \lambda)$, defined on the non-negative reals, is

$$m(f, \lambda) = m\{x \in \Omega \mid |f(x)|_A > \lambda\}. \quad (2.3)$$

The *non-increasing rearrangement of f* is the map $f^* : \mathbb{R}_+ \cup \{0\} \rightarrow \mathbb{R}_+ \cup \{0\}$ given by

$$f^*(s) = \inf\{\lambda \geq 0 \mid m(f, \lambda) \leq s\}. \quad (2.4)$$

Definition 2.1.4. Let A be a Banach space, $\Omega \subset \mathbb{R}^n$ open, $1 < p < \infty$ and $1 \leq q \leq \infty$. Then the *seminormed Lorentz space $L^{p,q}(\Omega, A)$* is the following set

$$\begin{aligned} & \{f : \Omega \rightarrow A \mid f \text{ strongly measurable, } \|f\|_{L^{p,q}(\Omega, A)} < \infty\} \\ \|f\|_{L^{p,q}(\Omega, A)} &= \left(\int_0^\infty (s^{1/p} f^*(s))^q \frac{ds}{s} \right)^{1/q} \quad \text{if } q < \infty, \\ \|f\|_{L^{p,q}(\Omega, A)} &= \sup_{s \geq 0} s^{1/p} f^*(s) \quad \text{if } q = \infty, \end{aligned} \quad (2.5)$$

equipped with the equivalence $f = g$ if $f(x) = g(x)$ for almost all $x \in \Omega$.

The (normed) Lorentz space $L^{(p,q)}(\Omega, A)$ is defined as

$$\begin{aligned} & \{f : \Omega \rightarrow A \mid f \text{ strongly measurable, } \|f\|_{L^{(p,q)}(\Omega,A)} < \infty\} \\ \|f\|_{L^{(p,q)}(\Omega,A)} &= \left(\int_0^\infty (t^{1/p} f^{**}(t))^q \frac{dt}{t} \right)^{1/q} \quad \text{if } q < \infty, \\ \|f\|_{L^{(p,q)}(\Omega,A)} &= \sup_{s \geq 0} t^{1/p} f^{**}(t) \quad \text{if } q = \infty, \end{aligned} \quad (2.6)$$

where $f^{**}(t) = \frac{1}{t} \int_0^t f^*(s) ds$. Again, we set $f = g$ if they are equal almost everywhere.

Remark 2.1.5. The spaces $L^{p,\infty}(\Omega, A)$ and $L^{(p,\infty)}(\Omega, A)$ are sometimes written $L^{p^*}(\Omega, A)$ and are called *weak L^p -spaces*.

Remark 2.1.6. We often leave the domain Ω out of the notation, so write $L^{p,q}(A)$ and $L^{(p,q)}(A)$ for these spaces. On the other hand, sometimes we leave the range out. Whether the set is the domain or range should be clear from the context.

Theorem 2.1.7. *Let A be a Banach space, $\Omega \subset \mathbb{R}^n$ open, $1 < p < \infty$ and $1 \leq q \leq \infty$. Then $L^{p,q}(\Omega, A)$ is a complete semi-normed space and $L^{(p,q)}(\Omega, A)$ is a Banach space. Moreover $L^{p,q} \equiv L^{(p,q)}$ with*

$$\|f\|_{p,q} \leq \|f\|_{(p,q)} \leq \frac{p}{p-1} \|f\|_{p,q}. \quad (2.7)$$

The spaces have the following properties:

- If $1 \leq q \leq Q \leq \infty$ then $L^{(p,q)} \hookrightarrow L^{(p,Q)}$ and $L^{(p,p)} = L^p$
- $\| |f|^r \|_{p,q} = \|f\|_{pr,qr}^r$ for $r \geq 1$.
- Simple functions are dense in $L^{(p,q)}$ if $q < \infty$
- Countably valued $L^{(p,\infty)}$ functions are dense in $L^{(p,\infty)}$

Proof. Note that if $f : \Omega \rightarrow A$ is strongly measurable, then $|f|_A : \Omega \rightarrow \mathbb{R}$ is measurable. Hence most of the proofs follow exactly like in the complex-valued case, for example in chapter 1.4. of Grafakos [20]. The following all refer to that book. Completeness and equivalence follow from 1.4.11, 1.4.12. The inclusions follow from 1.4.10 and the L^p equality from 1.4.5(12). The proof of the exponential scaling of the norm is given by 1.4.7.

Densities will be proven using a different source. The spaces $L^{(p,q)}(\Omega, A)$ of this theorem can be gotten using real interpolation on the Banach couple $(L^{p_0}(\Omega, A), L^{p_1}(\Omega, A))$ with some $1 < p_0 < p < p_1 < \infty$ according to theorem

5.2.1 in [7]. Simple functions are dense in the spaces $L^p(\Omega, A)$ for $1 \leq p < \infty$ by corollary III.3.8 in [17], hence they are so in the intersection $L^{p_0} \cap L^{p_1}$ too. The latter is dense in $L^{(p,q)}(\Omega, A)$ when $q < \infty$ by theorem 3.4.2 of [7]. This inclusion is a bounded linear operator, so simple functions are dense in $L^{(p,q)}(\Omega, A)$.

Let $f \in L^{(p,\infty)}(\Omega, A)$. Split Ω into a countable number of disjoint bounded and measurable sets Ω_j . According to corollary 3 of section II.1 in [15], there are countably valued measurable functions $s_j : \Omega_j \rightarrow A$ such that

$$|f(x) - s_j(x)|_A < \epsilon 2^{-j} \min(1, \|\chi_j\|_{(p,\infty)}^{-1}) \quad (2.8)$$

for all $x \in \Omega_j$. We write $\chi_j = \chi_{\Omega_j}$. Note that $s_j \in L^{(p,\infty)}(\Omega_j, A)$. Extend s_j by zero to the whole domain Ω and let $s(x) = \sum_j s_j(x)$. Now

$$\begin{aligned} \|f - s\|_{L^{(p,\infty)}(\Omega)} &\leq \sum_{j=1}^{\infty} \|(f - s_j)\chi_j\|_{L^{(p,\infty)}(\Omega)} \\ &\leq \sum_{j=1}^{\infty} \|\chi_j\|_{L^{(p,\infty)}(\Omega)} \sup_{x \in \Omega_j} |f(x) - s_j(x)|_A < \epsilon \sum_{j=1}^{\infty} 2^{-j} = \epsilon. \end{aligned} \quad (2.9)$$

Moreover, s is a countable sum of countably valued measurable functions, so it satisfies our claim. \square

Lemma 2.1.8 (Minkowski's integral inequality). *Let A be Banach, $\Omega \subset \mathbb{R}^n$ and $S \subset \mathbb{R}^m$ both open. Moreover let $1 < p < \infty$ and $1 \leq q \leq \infty$. Let $f : \Omega \times S \rightarrow A$ be strongly measurable. If $f(\cdot, y) \in L^{(p,q)}(\Omega, A)$ for almost all $y \in S$ and $y \mapsto \|f(\cdot, y)\|_{(p,q)}$ is in $L^1(S, \mathbb{R})$, then*

$$x \mapsto \int_S f(x, y) dm(y) \quad (2.10)$$

is in $L^{(p,q)}(\Omega, A)$ and

$$\left\| \int_S f(\cdot, y) dm(y) \right\|_{L^{(p,q)}(\Omega, A)} \leq C_p \int_S \|f(\cdot, y)\|_{L^{(p,q)}(\Omega, A)} dm(y) \quad (2.11)$$

where $C_p < \infty$ depends only on p .

Proof. Denote $g(x) = \int_S |f(x, y)|_A dm(y)$, so $g : \Omega \rightarrow \mathbb{R} \cup \{\infty\}$ is measurable by Fubini's theorem, for example 8.8.a in [44]. We will first show that the real valued $g \in (L^{(p',q')}(\Omega))^*$, where $a^{-1} + a'^{-1} = 1$ for $a = p, q$. This will imply

that $g \in L^{(p,q)}(\Omega)$ by theorem 1.4.17 in [20] and lemma 2 in [14] because they show that

$$(L^{(p',q')}(\Omega))^* \cap \{\text{measurable functions}\} \subset L^{(p,q)}(\Omega) \quad (2.12)$$

assuming that the measure is non-atomic, which m is. The right-hand side of the next estimate will be finite, hence we may use Fubini's theorem. It implies, with the generalized Hölder's inequality of O'Neil [41], that

$$\begin{aligned} \left\| w \mapsto \int_{\Omega} g w d m \right\|_{(L^{(p',q')}(\Omega))^*} &= \sup_{\|w\|_{(p',q')}=1} \left| \int_{\Omega} g(x) w(x) d m(x) \right| \\ &\leq \sup_{\|w\|_{(p',q')}=1} \int_{\Omega} g(x) |w(x)| d m(x) \\ &= \sup_{\|w\|_{(p',q')}=1} \int_S \int_{\Omega} |f(x, y)|_A |w(x)| d m(x) d m(y) \\ &\leq \sup_{\|w\|_{(p',q')}=1} \int_S \|f(\cdot, y)\|_{(p,q)} \|w\|_{(p',q')} d m(y) = RHS < \infty \end{aligned} \quad (2.13)$$

by the assumptions on f . Hence $g \in L^{(p,q)}(\Omega, \mathbb{R})$ and so $y \mapsto f(x, y)$ is integrable for almost all x . It remains to show that $x \mapsto \int_S f(x, y) d m(y)$ is strongly measurable, since then

$$\left\| \int_S f(\cdot, y) d m(y) \right\|_{L^{(p,q)}(\Omega, A)} \leq \|g\|_{L^{(p,q)}(\Omega, \mathbb{R})} \leq C_p \|g\|_{(L^{(p',q')}(\Omega, \mathbb{R}))^*}, \quad (2.14)$$

and so it is in $L^{(p,q)}(\Omega, A)$.

Let $S_m : \Omega \times S \rightarrow A$ be simple functions such that $S_m(x, y) \rightarrow f(x, y)$ almost everywhere. We may assume that $|S_m(x, y)|_A \leq |f(x, y)|_A$ by considering $t_m S_m |S_m|_A^{-1}$ instead of S_m , where t_m are simple real-valued functions rising to $|f|_A$. We may also assume that S_m has bounded support. Define $s_{x,m}(y) = S_m(x, y)$. Now $s_{x,m}$ is a simple function on S , $s_{x,m}(y) \rightarrow f(x, y)$ for almost all y for almost all x , and $|s_{x,m}(y)|_A \leq |f(x, y)|_A \in L^1(S)$ for almost all x . Hence, for almost all x , we get

$$\int_S f(x, y) d m(y) = \lim_{m \rightarrow \infty} \int_S s_{x,m}(y) d m(y) \quad (2.15)$$

by dominated convergence. The latter integrals are strongly measurable, so the claim follows. \square

2.2 Interpolation of Lorentz spaces

We use definitions like in [7] when interpolating. In particular $(\cdot, \cdot)_{[\theta]}$ represents complex interpolation. We give a short definition and a few examples. After them, we interpolate Banach-valued Lorentz spaces. The proof is an almost exact replica of theorem 5.1.2 in [7], where Bergh and Löfström interpolate Banach valued L^p spaces.

Definition 2.2.1. Let A_0, A_1 be topological vector spaces and assume that there is a Hausdorff topological vector space \mathcal{H} such that $A_0, A_1 \hookrightarrow \mathcal{H}$. Then A_0 and A_1 are *compatible*.

Definition 2.2.2. Let A_0 and A_1 be Banach spaces which are subspaces of a Hausdorff topological vector space \mathcal{H} . Then (A_0, A_1) is said to be a *compatible Banach couple*, or a *Banach couple* for short.

Remark 2.2.3. Compatible couples are normally defined like this: If \mathcal{C} is a subcategory of all normed vector spaces, then (A_0, A_1) is a *compatible couple in \mathcal{C}* if these conditions hold: i) A_0 and A_1 are compatible, ii) $A_0 \cap A_1 \in \mathcal{C}$ and iii) $A_0 + A_1 \in \mathcal{C}$. Our definition satisfies this in the category of Banach spaces by lemma 2.3.1 in [7].

Definition 2.2.4. Let $S = \{z \in \mathbb{C} \mid 0 < \operatorname{Re} z < 1\}$ and $\overline{A} = (A_0, A_1)$ be a compatible Banach couple. Then $\mathcal{F}(\overline{A})$ consist of the all the functions $f : \overline{S} \rightarrow A_0 + A_1$ satisfying

- f is bounded and continuous when $A_0 + A_1$ is equipped with the norm $\|a\|_{A_0+A_1} = \inf_{a=a_0+a_1} \|a_0\|_{A_0} + \|a_1\|_{A_1}$
- f is analytic on S
- the maps $t \mapsto f(it)$, $t \mapsto f(1+it)$ are continuous $\mathbb{R} \rightarrow A_0$, $\mathbb{R} \rightarrow A_1$, respectively, and they tend to zero as $|t| \rightarrow \infty$

We equip $\mathcal{F}(\overline{A})$ with the norm

$$\|f\|_{\mathcal{F}(A_0, A_1)} = \max \left(\sup_{t \in \mathbb{R}} \|f(it)\|_{A_0}, \sup_{t \in \mathbb{R}} \|f(1+it)\|_{A_1} \right). \quad (2.16)$$

Remark 2.2.5. $\mathcal{F}(\overline{A})$ is a Banach space by theorem 4.1.1 of [7].

Definition 2.2.6. Let (A_0, A_1) be a Banach couple and $0 \leq \theta \leq 1$. Then

$$(A_0, A_1)_{[\theta]} = \{a \in A_0 + A_1 \mid a = f(\theta) \text{ for some } f \in \mathcal{F}(A_0, A_1)\} \quad (2.17)$$

and we equip it with the norm

$$\|a\|_{[\theta]} = \|a\|_{(A_0, A_1)_{[\theta]}} = \inf \{ \|f\|_{\mathcal{F}(A_0, A_1)} \mid f(\theta) = a, f \in \mathcal{F}(A_0, A_1) \}. \quad (2.18)$$

The structure $((A_0, A_1)_{[\theta]}, \|\cdot\|_{[\theta]})$ is called a *complex interpolation space*.

Theorem 2.2.7. Let $\bar{A} = (A_0, A_1)$ and $\bar{B} = (B_0, B_1)$ be Banach couples and $0 \leq \theta \leq 1$. Then $\bar{A}_{[\theta]}$ and $\bar{B}_{[\theta]}$ are Banach spaces with continuous embeddings¹ $A_0 \cap A_1 \hookrightarrow \bar{A}_{[\theta]} \hookrightarrow A_0 + A_1$ and the same for B . Moreover if

$$\begin{aligned} T : A_0 &\rightarrow B_0 && \text{with norm } M_0 \\ T : A_1 &\rightarrow B_1 && \text{with norm } M_1 \end{aligned} \quad (2.19)$$

then $T : \bar{A}_{[\theta]} \rightarrow \bar{B}_{[\theta]}$ with norm at most $M_0^{1-\theta} M_1^\theta$.

Proof. See theorem 4.1.2 in [7] and the definitions of intermediate spaces and exact interpolation functors 2.4.1, 2.4.3 in that same book. \square

Theorem 2.2.8 (Multilinear interpolation). Let \bar{A} , \bar{B} and \bar{X} be Banach couples and $0 \leq \theta \leq 1$. Assume that $T : (A_0 \cap A_1) \times (B_0 \cap B_1) \rightarrow (X_0 \cap X_1)$ is multilinear and

$$\begin{aligned} \|T(a, b)\|_{X_0} &\leq M_0 \|a\|_{A_0} \|b\|_{B_0} \\ \|T(a, b)\|_{X_1} &\leq M_1 \|a\|_{A_1} \|b\|_{B_1} \end{aligned} \quad (2.20)$$

for $a \in A_0 \cap A_1$ and $b \in B_0 \cap B_1$. Then T can be uniquely extended to a multilinear mapping $\bar{A}_{[\theta]} \times \bar{B}_{[\theta]} \rightarrow \bar{X}_{[\theta]}$ with $\|T(a, b)\|_{X_{[\theta]}} \leq M_0^{1-\theta} M_1^\theta \|a\|_{A_{[\theta]}} \|b\|_{B_{[\theta]}}$.

Proof. See theorem 4.4.1 in [7]. \square

Example 2.2.9. Let $0 \leq \theta \leq 1$. Let's prove that $(A, A)_{[\theta]} = A$ with equal norm to get a hold of the definitions. Let $a \in (A, A)_{[\theta]}$. Then there is $f \in \mathcal{F}(A, A)$ such that $a = f(\theta)$. We may assume that $\|f\|_{\mathcal{F}} \leq (1 + \epsilon) \|a\|_{[\theta]}$ by the definition of the norm in $(A, A)_{[\theta]}$. Now $a = f(\theta) \in A + A = A$, and

$$\begin{aligned} \|a\|_A &= \|f(\theta)\|_A \leq \max \left(\sup \|f(it)\|_A, \sup \|f(1 + it)\|_A \right) \\ &= \|f\|_{\mathcal{F}} \leq (1 + \epsilon) \|a\|_{[\theta]} \end{aligned} \quad (2.21)$$

because of the Phragmén-Lindelöf principle. This is allowed since f is bounded on \bar{S} . Taking similar $f \in \mathcal{F}$ while letting $\epsilon \rightarrow 0$ gives $\|a\|_A \leq \|a\|_{[\theta]}$.

Now let $a \in A$. Let's construct a suitable $f \in \mathcal{F}(A, A)$. Let

$$f(z) = e^{\epsilon(z-\theta)^2} a = e^{\epsilon((\operatorname{Re} z - \theta)^2 - (\operatorname{Im} z)^2 + 2i \operatorname{Im} z (\operatorname{Re} z - \theta))} a. \quad (2.22)$$

The function f is clearly continuous and bounded on \bar{S} and analytic on S . The continuity from the boundary to the respective spaces follows since we have just one Banach space. Finally, $\|f(it)\|_A = \exp(\epsilon(\theta^2 - t^2)) \|a\|_A \rightarrow 0$ as

¹ $A_0 \cap A_1$ is equipped with the norm $\|a\|_{A_0 \cap A_1} = \max(\|a\|_{A_0}, \|a\|_{A_1})$ and $A_0 + A_1$ is equipped with $\|a\|_{A_0 + A_1} = \inf_{a=a_0+a_1} \|a_0\|_{A_0} + \|a_1\|_{A_1}$

$|t| \rightarrow \infty$. The same holds for $f(1 + it)$, so $f \in \mathcal{F}(A, A)$. Also $f(\theta) = a$, so $a \in (A, A)_{[\theta]}$. Now

$$\begin{aligned} \|a\|_{[\theta]} &\leq \|f\|_{\mathcal{F}} = \max(\sup \|f(it)\|_A, \|f(1 + it)\|_A) \\ &= \max(\sup e^{\epsilon(\theta^2 - t^2)}, \sup e^{\epsilon((1-\theta)^2 - t^2)}) \|a\|_A \leq e^\epsilon \|a\|_A. \end{aligned} \quad (2.23)$$

Letting $\epsilon \rightarrow 0$ shows that $\|a\|_{[\theta]} \leq \|a\|_A$.

Example 2.2.10. We also have $(L^1, L^\infty)_{[\frac{1}{p}]} = L^p$. The proof is based on choosing

$$f = e^{\epsilon(z^2 - \frac{1}{p^2})} |a|^{p(1-z)} \frac{a}{|a|} \quad (2.24)$$

and using *the three lines theorem*. For details, check theorem 5.1.1 in [7].

Remark 2.2.11. It would seem that the direction $(A_0, A_1)_{[\theta]} \hookrightarrow X$ requires often the use of complex analysis, while the other one doesn't. In example 2.2.9, we used the Phragmén-Lindelöf principle when proving that $(A, A)_{[\theta]} \hookrightarrow A$. In example 2.2.10, the three lines theorem comes into play when showing that $(L^1, L^\infty)_{[\theta]} \hookrightarrow L^p$. Lastly, the proof of the next theorem will require properties of the Poisson kernel of S when showing that same direction.

We will not write out the domain \mathbb{R}^n . The proof works for any domain.

Theorem 2.2.12. *Let (A_0, A_1) be a compatible Banach couple, $1 < p_j < \infty$ and $1 \leq q_j < \infty$. Let $0 < \theta < 1$ and $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$, $\frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$. Then*

$$\begin{aligned} L^{(p, p \min(\frac{q_0}{p_0}, \frac{q_1}{p_1}))}((A_0, A_1)_{[\theta]}) \\ \subset (L^{(p_0, q_0)}(A_0), L^{(p_1, q_1)}(A_1))_{[\theta]} \\ \subset L^{(p, q)}((A_0, A_1)_{[\theta]}) \end{aligned} \quad (2.25)$$

and

$$(L^{(p, \infty)}(A_0), L^{(p, \infty)}(A_1))_{[\theta]} = L^{(p, \infty)}((A_0, A_1)_{[\theta]}) \quad (2.26)$$

with corresponding norm estimates.

Proof. Since (A_0, A_1) is a Banach couple, so are the other pairs of spaces in the theorem. We may interpolate. The idea is to take $a \in L^{(\cdot, \cdot)}((A_0, A_1)_{[\theta]})$ and then, for each x , take an analytic $A_0 + A_1$ -valued function $g_x(z)$ satisfying $g_x(\theta) = a(x)$. After that we show that $x \mapsto g_x(z)$ is a strongly measurable function, so $z \mapsto g(z)$ would actually be in $\mathcal{F}(L^{(\cdot, \cdot)}(A_0), L^{(\cdot, \cdot)}(A_1))$. Simple functions are dense in all of these spaces when $q, q_j < \infty$ and countably simple functions are so for $q = \infty$ by theorem 2.1.7. Using these makes the above much easier.

Consider the case of $q_j, q < \infty$ first. Note that $p < \infty$, so the simple functions must have support with finite measure. Let

$$\Xi = \left\{ s : \mathbb{R}^n \rightarrow A_0 \cap A_1 \mid \exists N \in \mathbb{N}, a_k \in A_0 \cap A_1, E_k \subset \mathbb{R}^n, m(E_j) < \infty, \right. \\ \left. E_j \cap E_k = \emptyset \text{ for } j \neq k, \text{ and such that } s(x) = \sum_{k=0}^N a_k \chi_{E_k}(x) \right\} \quad (2.27)$$

It is enough to assume that $a \in \Xi$. This is because of the following. The set $A_0 \cap A_1$ is dense in $(A_0, A_1)_{[\theta]}$ by theorem 4.2.2. of [7]. Hence Ξ is dense in $L^{(p,q)}((A_0, A_1)_{[\theta]})$. Moreover Ξ is dense in $L^{(p_0, q_0)}(A_0) \cap L^{(p_1, q_1)}(A_1)$, hence in $(L^{(p_0, q_0)}(A_0), L^{(p_1, q_1)}(A_1))_{[\theta]}$ too by that same theorem.

Let $a \in \Xi \subset L^{(p,q)}((A_0, A_1)_{[\theta]})$. To simplify notation we assume that $\|a\|_{(p, p \min(q_0/p_0, q_1/p_1))} = 1$ and write

$$a(x) = \sum_{k=0}^N a_k \chi_{E_k}(x). \quad (2.28)$$

Let $\epsilon > 0$. We have $a(x) \in (A_0, A_1)_{[\theta]}$ for each $x \in \mathbb{R}^n$. Then, for $x \in \mathbb{R}^n$, there exists $g_x \in \mathcal{F}(A_0, A_1)$ such that $\|g_x\|_{\mathcal{F}(A_0, A_1)} \leq (1 + \epsilon) |a(x)|_{(A_0, A_1)_{[\theta]}}$ and $g_x(\theta) = a(x)$. If $a(x) = a(y)$, take $g_x = g_y$. Define

$$\phi(z) = g(z) |a|_{(A_0, A_1)_{[\theta]}}^{p(\frac{1}{p_0} - \frac{1}{p_1})(z-\theta)}. \quad (2.29)$$

Now, given any $z \in \overline{S}$, $\phi(z)$ is strongly measurable² $\mathbb{R}^n \rightarrow A_0 + A_1$, ϕ is analytic $S \rightarrow L^{(p_0, q_0)}(A_0) + L^{(p_1, q_1)}(A_1)$, continuous on \overline{S} , $\phi(it) \in L^{(p_0, q_0)}(A_0)$, $\phi(1 + it) \in L^{(p_1, q_1)}(A_1)$, they are continuous and tend to zero as $|t| \rightarrow \infty$. Hence $\phi \in \mathcal{F}(L^{(p_0, q_0)}(A_0), L^{(p_1, q_1)}(A_1))$. Moreover $\phi(\theta) = a$. Now

$$\|a\|_{(L^{(p_0, q_0)}(A_0), L^{(p_1, q_1)}(A_1))_{[\theta]}} \leq \|\phi\|_{\mathcal{F}(L^{(p_0, q_0)}(A_0), L^{(p_1, q_1)}(A_1))} \\ = \max \left(\sup_{t \in \mathbb{R}} \|\phi(it)\|_{L^{(p_0, q_0)}(A_0)}, \sup_{t \in \mathbb{R}} \|\phi(1 + it)\|_{L^{(p_1, q_1)}(A_1)} \right). \quad (2.30)$$

Let's estimate the first supremum. Note that $\| |g|^r \|_{p,q} = \|g\|_{pr, qr}^r$ by theorem

²Because in fact $g(z) = \sum_{k=0}^N b_k(z) \chi_{E_k}$, where $b_k \in \mathcal{F}(A_0, A_1)$ gives $b_k(\theta) = a_k$.

2.1.7 and $\|g\|_{p,q} \leq \|g\|_{(p,q)} \leq \frac{p}{p-1} \|g\|_{p,q}$. Then

$$\begin{aligned}
\|\phi(it)\|_{L^{(p_0,q_1)}(A_0)} &= \left\| |g(it)|_{A_0} |a|_{(A_0,A_1)_{[\theta]}}^{-\theta p(\frac{1}{p_0} - \frac{1}{p_1})} \right\|_{L^{(p_0,q_0)}(\mathbb{R})} \\
&\leq \left\| \|g\|_{\mathcal{F}(A_0,A_1)} |a|_{(A_0,A_1)_{[\theta]}}^{-\theta p(\frac{1}{p_0} - \frac{1}{p_1})} \right\|_{L^{(p_0,q_0)}(\mathbb{R})} \leq (1 + \epsilon) \left\| |a|_{(A_0,A_1)_{[\theta]}}^{1 - \theta p(\frac{1}{p_0} - \frac{1}{p_1})} \right\|_{L^{(p_0,q_0)}(\mathbb{R})} \\
&= (1 + \epsilon) \left\| |a|_{(A_0,A_1)_{[\theta]}}^{p/p_0} \right\|_{L^{(p_0,q_0)}(\mathbb{R})} \leq C_{p_0} (1 + \epsilon) \left\| |a|_{(A_0,A_1)_{[\theta]}} \right\|_{L^{(p,pq_0/p_0)}(\mathbb{R})}^{p/p_0} \\
&= C_{p_0} (1 + \epsilon) \|a\|_{L^{(p,pq_0/p_0)}((A_0,A_1)_{[\theta]})}^{p/p_0}. \quad (2.31)
\end{aligned}$$

We get similarly

$$\|\phi(1 + it)\|_{L^{(p_1,q_1)}(A_1)} \leq \dots \leq C_{p_1} (1 + \epsilon) \|a\|_{L^{(p,pq_1/p_1)}((A_0,A_1)_{[\theta]})}^{p/p_1}. \quad (2.32)$$

Reducing the second parameter of the Lorentz spaces gives a smaller space, and we made the assumption of $\|a\|_{(p,p \min(q_0/p_0, q_1/p_1))} = 1$, so

$$\|a\|_{(L^{(p_0,q_1)}(A_0), L^{(p_1,q_1)}(A_1))_{[\theta]}} \leq C_{p_0,p_1} \|a\|_{L^{(p,p \min(\frac{q_0}{p_0}, \frac{q_1}{p_1}))}((A_0,A_1)_{[\theta]})}. \quad (2.33)$$

The other direction requires Minkowski's integral inequality of lemma 2.1.8 and the inequality

$$\begin{aligned}
|f(\theta)|_{(A_0,A_1)_{[\theta]}} &\leq \left(\frac{1}{1 - \theta} \int_{-\infty}^{\infty} |f(i\tau)|_{A_0} P_0(\theta, \tau) d\tau \right)^{1-\theta} \\
&\quad \cdot \left(\frac{1}{\theta} \int_{-\infty}^{\infty} |f(1 + i\tau)|_{A_1} P_1(\theta, \tau) d\tau \right)^{\theta} \quad (2.34)
\end{aligned}$$

proven in lemma 4.3.2 of [7]. Here $f \in \mathcal{F}(A_0, A_1)$ and

$$P_j(s + it, \tau) = \frac{e^{-\pi(\tau-t)} \sin \pi s}{\sin^2 \pi s + (\cos \pi s - e^{ij\pi - \pi(\tau-t)})^2} \quad (2.35)$$

is the Poisson kernel of the strip S .

Let $a \in (L^{(p_0,q_0)}(A_0), L^{(p_1,q_1)}(A_1))_{[\theta]}$. Then there is a corresponding analytic $f \in \mathcal{F}(L^{(p_0,q_0)}(A_0), L^{(p_1,q_1)}(A_1))$ such that $f(\theta) = a$ and whose norm is bounded by $\|f\|_{\mathcal{F}} \leq (1 + \epsilon) \|a\|_{[\theta]}$. Note that $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and $\frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$, so the generalized Hölder's inequality given in theorem 3.4 of [41] allows us

to take the norms of the factors in the product. Everything is then ready:

$$\begin{aligned}
\|a\|_{L^{(p,q)}((A_0,A_1)_{[\theta]})} &= \left\| |f(\theta)|_{(A_0,A_1)_{[\theta]}} \right\|_{L^{(p,q)}(\mathbb{R})} \\
&\leq \left\| \left(\frac{1}{1-\theta} \int_{-\infty}^{\infty} |f(i\tau)|_{A_0} P_0(\theta, \tau) d\tau \right)^{1-\theta} \right\|_{\left(\frac{p_0}{1-\theta}, \frac{q_0}{1-\theta}\right)} \\
&\quad \cdot \left\| \left(\frac{1}{\theta} \int_{-\infty}^{\infty} |f(1+i\tau)|_{A_1} P_1(\theta, \tau) d\tau \right)^{\theta} \right\|_{\left(\frac{p_1}{\theta}, \frac{q_1}{\theta}\right)} \\
&\leq C_{p_0,p_1,\theta} \left\| \int_{-\infty}^{\infty} |f(i\tau)|_{A_0} P_0(\theta, \tau) d\tau \right\|_{(p_0,q_0)}^{1-\theta} \\
&\quad \cdot \left\| \int_{-\infty}^{\infty} |f(1+i\tau)|_{A_1} P_1(\theta, \tau) d\tau \right\|_{(p_1,q_1)}^{\theta} \\
&\leq C_{p_0,p_1,\theta} \left(\int_{-\infty}^{\infty} \|f(i\tau)\|_{L^{(p_0,q_0)}(A_0)} P_0(\theta, \tau) d\tau \right)^{1-\theta} \\
&\quad \cdot \left(\int_{-\infty}^{\infty} \|f(1+i\tau)\|_{L^{(p_1,q_1)}(A_1)} P_1(\theta, \tau) d\tau \right)^{\theta} \\
&\leq C_{p_0,p_1,\theta} \|f\|_{\mathcal{F}(L^{(p_0,q_0)}(A_0), L^{(p_1,q_1)}(A_1))} \\
&\leq C_{p_0,p_1,\theta} (1+\epsilon) \|a\|_{(L^{(p_0,q_0)}(A_0), L^{(p_1,q_1)}(A_1))_{[\theta]}} < \infty. \quad (2.36)
\end{aligned}$$

The last claim follows similarly, except that we use

$$\begin{aligned}
\Xi &= \left\{ s \in L^{(p,\infty)}(A_0 \cap A_1) \mid \exists a_k \in A_0 \cap A_1, E_k \subset \mathbb{R}^n, m(E_j) < \infty, \right. \\
&\quad \left. E_j \cap E_k = \emptyset \text{ for } j \neq k, \text{ and such that } s(x) = \sum_{k=0}^{\infty} a_k \chi_{E_k}(x) \right\}, \quad (2.37)
\end{aligned}$$

which is dense in $L^{(p,\infty)}(A_0 \cap A_1)$ by theorem 2.1.7. We get density in $L^{(p,\infty)}(A_0) \cap L^{(p,\infty)}(A_1)$ and $L^{(p,\infty)}((A_0, A_1)_{[\theta]})$ because $A_0 \cap A_1$ is dense in $(A_0, A_1)_{[\theta]}$. All other steps are the same, but with simpler expressions. \square

Remark 2.2.13. The same proof works for Lorentz spaces defined on a domain.

Remark 2.2.14. If $\frac{q_0}{p_0} = \frac{q_1}{p_1}$, then the theorem shows that

$$(L^{(p_0,q_0)}(A_0), L^{(p_1,q_1)}(A_1))_{[\theta]} = L^{(p,q)}((A_0, A_1)_{[\theta]}). \quad (2.38)$$

Maybe a better choice of f could prove this without assuming anything from our parameters.

Remark 2.2.15. Why can't we have $p_0 \neq p_1$ when $q_0 = \infty$? Maybe we could, but this proof won't work then. The problem is to find a set Ξ of quite "simple" functions which would be dense in all the spaces considered at the same time. On the other hand, Adams and Fournier claim this result in 7.56 [1], assuming that $A_0 = A_1$. In that case it would follow from reiteration with real interpolation e.g. by theorem 4.7.2 of [7].

Remark 2.2.16. By Cwikel $(L^{p_0}(A_0), L^{p_1}(A_1))_{\theta, q}$ is not necessarily a Lorentz space [14]. So it is not possible to use reiteration to prove our claim in general if $A_0 \neq A_1$.

2.3 Lorentz-Sobolev spaces

Definition 2.3.1. Let $X \subset \mathbb{R}^n$ be any nonempty set and A be a Banach space. Then the *space of bounded continuous A -valued functions* is

$$BC(X, A) = \{f : X \rightarrow A \mid f \text{ is continuous and bounded}\}, \quad (2.39)$$

equipped with the norm $\|f\|_{BC(X, A)} = \sup_{z_0 \in X} |f(z_0)|_A$.

Remark 2.3.2. This is a Banach space.

Definition 2.3.3. Let $\Omega \subset \mathbb{R}^n$ open, $1 < p < \infty$, $1 \leq q \leq \infty$ and $k \in \mathbb{N}$. Define the *Lorentz-Sobolev space* $W^{k, (p, q)}(\Omega)$ as follows:

$$W^{k, (p, q)}(\Omega) = \{f \in L^{(p, q)}(\Omega) \mid D^\alpha f \in L^{(p, q)}(\Omega) \text{ for } |\alpha| \leq k\} \quad (2.40)$$

with norm

$$\|f\|_{W^{k, (p, q)}(\Omega)} = \|f\|_{L^{(p, q)}(\Omega)} + \sum_{\|\alpha\| \leq k} \|D^\alpha f\|_{L^{(p, q)}(\Omega)}. \quad (2.41)$$

where D^α is the distribution derivative in Ω .

Theorem 2.3.4. *Let $\Omega \subset \mathbb{R}^n$ be an open set satisfying the cone and segment conditions³, $1 < p < \infty$, $1 \leq q \leq \infty$ and $k \in \mathbb{N}$. Then the space $W^{k, (p, q)}(\Omega)$ is a well defined Banach space with the following properties:*

1. *The restrictions of $C_0^\infty(\mathbb{R}^n)$ test functions to Ω are dense in $W^{k, (p, q)}(\Omega)$ for $q < \infty$*
2. *We have the continuous embedding $W^{k, (\frac{n}{k}, 1)}(\Omega) \hookrightarrow BC(\bar{\Omega})$ for $k \geq 1$*

³See for example 4.5 and 4.6 in [1]. The cone condition prevents cusps while the segment condition ensures that the domain is never on both sides of the boundary, i.e. $] - 1, 0[\cup] 0, 1[$ is not allowed. Bounded Lipschitz domains have this property.

Proof. The proofs are more or less the same as for the usual Sobolev spaces $W^{k,p}(\Omega)$, but using the result

$$\|fg\|_{L^1(\Omega)} \leq \|f\|_{L^{(p,q)}(\Omega)} \|g\|_{L^{(p',q')}(\Omega)} \quad \frac{1}{p} + \frac{1}{p'} = 1 \quad \frac{1}{q} + \frac{1}{q'} = 1 \quad (2.42)$$

proven in [41] instead of the usual Hölder's inequality. We will refer to Adams and Fournier [1]. Completeness follows like in 3.3 using the completeness of $L^{(p,q)}(\Omega)$. The density of test functions follows by proving 3.22, 3.16, 2.29 and 2.19. Mimicking the proof of theorem 2.19 requires the density of simple functions, which requires $q < \infty$.

The density of the restriction of test functions to Ω , the property that $|x|^{k-n} \in L^{(\frac{n}{n-k}, \infty)}$ and 4.15 imply the embedding $\text{Id} : W^{k,(\frac{n}{k},1)}(\Omega) \hookrightarrow L^\infty(\Omega)$, and hence also into $BC(\Omega)$. We still need to show that the elements can be extended uniquely to the boundary. But this follows directly from the fact that elements in $W^{k,(\frac{n}{k},1)}(\Omega)$ can be approximated by restrictions of test functions, and those can be extended uniquely. \square

Remark 2.3.5. The idea for such spaces did arise quite naturally after proving the estimate in theorem 4.1.1. The fact that $W^{1,p} \hookrightarrow C^{1-2/p}$ for $p > 2$ and $W^{1,2} \hookrightarrow \cap_{q < \infty} L^q$ gives a natural hint for the embedding proved here because $L_{loc}^{2+\epsilon} \hookrightarrow L_{loc}^{(2,1)} \hookrightarrow L_{loc}^2$. Moreover, the embeddings into L^∞ can be proven using integrals with kernels having weak singularities. Operators with kernels in weak L^p spaces work well with Lorentz spaces because of O'Neil's inequality [41].

Remark 2.3.6. The idea of combining Lorentz and Sobolev spaces is not new. See for example [28] and [38]. In fact, in the first one, the authors consider functions $f \in W_{loc}^{1,1}$ such that $\nabla f \in L^{(n,1)}$, and show that those are continuous. They also prove that if X is a rearrangement invariant Banach space, then $\{u \mid \nabla u \in X\} \hookrightarrow AC^n$ if and only if $X \hookrightarrow L^{(n,1)}$. Here AC^n denotes the space of n -absolutely continuous functions, see [32].

Remark 2.3.7. Is it reasonable to require both f and ∇f in the same $L^{(p,q)}(\Omega)$? Consider for example $|x|^{-a}$ in the spaces $W^{k,(p,\infty)}$. We have $|x|^{-a} \in L^{(p,\infty)}$ if and only if $p = \frac{n}{a}$, but $|\nabla |x|^{-a}| = a|x|^{-a-1} \in L^{(p^*,\infty)}$ if and only if $p^* = \frac{n}{a+1}$. This would suggest that $W^{1,(p,\infty)}$ should be defined by taking the norms $\|f\|_{(p,\infty)} + \|\nabla f\|_{(p^*,\infty)}$, where $\frac{1}{p^*} = \frac{1}{n} + \frac{1}{p}$. This is the famous Sobolev conjugate. One could wonder whether the Sobolev embedding theorems give a good choice for the norm also in the case of the usual $W^{1,p}$ -spaces. Moreover, our results will have mixed norms like $\|f\|_\infty + \|\nabla f\|_{(2,1)}$ in the estimates in 2D. Anyway, this is not a big problem when working on a bounded domain. Hence we shall be content with having the same Lorentz space for all the derivatives.

2.4 Lorentz-Bessel potential spaces

Definition 2.4.1. Let $1 < p < \infty$, $1 \leq q \leq \infty$ and $s \in \mathbb{R}$. Define the Lorentz-Bessel potential space $H^{s,(p,q)}(\mathbb{R}^n)$ as follows:

$$H^{s,(p,q)}(\mathbb{R}^n) = \{f \in L^{(p,q)}(\mathbb{R}^n) \mid \mathcal{F}^{-1}((1+|\cdot|^2)^{s/2} \mathcal{F}f) \in L^{(p,q)}(\mathbb{R}^n)\} \quad (2.43)$$

with norm

$$\|f\|_{H^{s,(p,q)}(\mathbb{R}^n)} = \|\mathcal{F}^{-1}((1+|\cdot|^2)^{s/2} \mathcal{F}f)\|_{L^{(p,q)}(\mathbb{R}^n)}. \quad (2.44)$$

If $\Omega \subset \mathbb{R}^n$ is an open set, then $H^{s,(p,q)}(\Omega)$ consists of the restrictions of $H^{s,(p,q)}(\mathbb{R}^n)$ distributions to Ω . Hence

$$H^{s,(p,q)}(\Omega) = \{f|_{\Omega} \mid f \in H^{s,(p,q)}(\mathbb{R}^n)\}, \quad \|f\|_{H^{s,(p,q)}(\Omega)} = \inf_{g|_{\Omega}=f} \|g\|_{H^{s,(p,q)}(\mathbb{R}^n)}. \quad (2.45)$$

The elements of these spaces are considered as distributions in \mathbb{R}^n and Ω respectively.

Theorem 2.4.2. Let $\Omega \subset \mathbb{R}^n$ be an open set, $s \in \mathbb{R}$, $1 < p < \infty$ and $1 \leq q \leq \infty$. Then the space $H^{s,(p,q)}(\Omega)$ is a well defined Banach space and we have the equivalence $W^{k,(p,q)}(\mathbb{R}^n) = H^{k,(p,q)}(\mathbb{R}^n)$ for $k \in \mathbb{N}$.

Proof. Completeness in the whole domain follows from the completeness of $L^{(p,q)}(\mathbb{R}^n)$ and standard argument like in 6.2.2 of [7]. We write $H(\mathbb{R}^n) = H^{s,(p,q)}(\mathbb{R}^n)$ and $H(\Omega) = H^{s,(p,q)}(\Omega)$. Let $(f_k) \subset H(\Omega)$ be a sequence such that the series

$$\sum_{k=0}^{\infty} \|f_k\|_{H(\Omega)} \quad (2.46)$$

converges in \mathbb{R} . By the definition of the norm of $H(\Omega)$, we have a sequence $g_k \in H(\mathbb{R}^n)$ such that $\|g_k\|_{H(\mathbb{R}^n)} \leq 2 \|f_k\|_{H(\Omega)}$ and $g_k|_{\Omega} = f_k$. Now

$$\sum_{k=0}^{\infty} \|g_k\|_{H(\mathbb{R}^n)} \leq 2 \sum_{k=0}^{\infty} \|f_k\|_{H(\Omega)} < \infty, \quad (2.47)$$

and since $H(\mathbb{R}^n)$ is complete, the series $\sum g_k$ converges to $g \in H(\mathbb{R}^n)$ in the norm. Now

$$\left\| \sum_{k=0}^m f_k - g|_{\Omega} \right\|_{H(\Omega)} \leq \left\| \sum_{k=0}^m g_k - g \right\|_{H(\mathbb{R}^n)} \longrightarrow 0 \quad (2.48)$$

as $m \rightarrow \infty$. Hence $\sum f_k$ converges to $g|_{\Omega} \in H(\Omega)$. All norm-convergent series converge, so $H(\Omega)$ is complete.

The equivalence of $W^{k,(p,q)}$ and $H^{k,(p,q)}$ follows by standard arguments, for example like in theorem 6.2.3 of [7]. The only modification is that the Mihlin multiplier theorem, 6.1.6 in [7], gives boundedness in $L^{(p,q)}(\mathbb{R}^n)$ too by real interpolation. \square

Remark 2.4.3. We didn't prove that $W^{k,(p,q)}(\Omega) = H^{k,(p,q)}(\Omega)$. This would require showing that there is an extension operator $W^{k,(p,q)}(\Omega) \rightarrow W^{k,(p,q)}(\mathbb{R}^n)$. This would seem to be true for regular enough Ω , for example by using Calderón's construction in theorem 12 of [10] and the fact that Calderón-Zygmund operators map $L^{(p,q)}$ to $L^{(p,q)}$ by real interpolation. Anyway, it's easy to see that $H^{k,(p,q)}(\Omega) \subset W^{k,(p,q)}(\Omega)$ by the theorem above.

Remark 2.4.4. One could define $W^{s,(p,q)}(\Omega)$ by interpolation, but it is not a-priori clear whether we would get $H^{s,(p,q)}(\Omega)$. It is true when Ω has a total extension operator.

Remark 2.4.5. All these kinds of spaces are not new. See for example [28] and 5.1, 5.3 in [38].

Definition 2.4.6. Let A and B be normed linear spaces. Then B is a *retract* of A , if there are bounded linear operators $\mathcal{I} : B \rightarrow A$ and $\mathcal{P} : A \rightarrow B$ such that $\mathcal{P} \circ \mathcal{I}$ is the identity in B .

$$\begin{array}{ccc}
 B & \xrightarrow{\text{Id}} & B \\
 & \searrow \mathcal{I} & \nearrow \mathcal{P} \\
 & & A
 \end{array} \tag{2.49}$$

Remark 2.4.7. This is extremely useful when interpolating. If (B_0, B_1) are retracts of (A_0, A_1) with the mappings \mathcal{I} and \mathcal{P} , then $(B_0, B_1)_{[\theta]}$ is a retract of $(A_0, A_1)_{[\theta]}$ with the same mappings. This is a direct consequence of the interpolation property.

Theorem 2.4.8. Let $s \in \mathbb{R}$, $1 < p < \infty$ and $1 \leq q \leq \infty$. Then $H^{s,(p,q)}(\mathbb{R}^n)$ is a retract of $L^{(p,q)}(\mathbb{R}^n, \ell_2^s)$. Moreover the mappings \mathcal{I} and \mathcal{P} do not depend on any of the parameters.

$$\begin{array}{ccc}
 H^{s,(p,q)} & \xrightarrow{\text{Id}} & H^{s,(p,q)} \\
 & \searrow \mathcal{I} & \nearrow \mathcal{P} \\
 & & L^{(p,q)}(\ell_2^s)
 \end{array} \tag{2.50}$$

Proof. The proof is a word by word replica of theorem 6.4.3 in [7]. The only change is the fact that the Mihlin multiplier theorem gives boundedness in $L^{(p,q)}$ by real interpolation. \square

The symbol $(\cdot, \cdot)_{[\theta]}$ represents complex interpolation as in [7]. See definition 2.2.6. We are ready to interpolate $H^{s,(p,q)}$.

Corollary 2.4.9. *Let $s_0 < s_1$ be real numbers and $1 < p < \infty$, $1 \leq q \leq \infty$. Let $0 < \theta < 1$. Then*

$$(H^{s_0,(p,q)}(\mathbb{R}^n), H^{s_1,(p,q)}(\mathbb{R}^n))_{[\theta]} = H^{s,(p,q)}(\mathbb{R}^n) \quad (2.51)$$

with equivalent norms, where $s = (1 - \theta)s_0 + \theta s_1$.

Proof. Let \mathcal{I} and \mathcal{P} be the injection and projection of theorem 2.4.8. According to remark 2.4.7 the interpolation space $(H^{s_0,(p,q)}, H^{s_1,(p,q)})_{[\theta]}$ is a retract of $(L^{(p,q)}(\ell_2^{s_0}), L^{(p,q)}(\ell_2^{s_1}))_{[\theta]}$. By theorem 2.2.12 and the vector-valued result of theorem 5.6.3 in [7] we know that the latter is just $L^{(p,q)}(\ell_2^s)$, of which $H^{s,(p,q)}$ is a retract. Hence

$$\begin{aligned} (H^{s_0,(p,q)}, H^{s_1,(p,q)})_{[\theta]} &= \mathcal{P}\mathcal{I}(H^{s_0,(p,q)}, H^{s_1,(p,q)})_{[\theta]} \\ &\subset \mathcal{P}(L^{(p,q)}(\ell_2^{s_0}), L^{(p,q)}(\ell_2^{s_1}))_{[\theta]} = \mathcal{P}L^{(p,q)}(\ell_2^s) \subset H^{s,(p,q)}. \end{aligned} \quad (2.52)$$

On the other hand

$$\begin{aligned} H^{s,(p,q)} &= \mathcal{P}\mathcal{I}H^{s,(p,q)} \subset \mathcal{P}L^{(p,q)}(\ell_2^s) \\ &= \mathcal{P}(L^{(p,q)}(\ell_2^{s_0}), L^{(p,q)}(\ell_2^{s_1}))_{[\theta]} \subset (H^{s_0,(p,q)}, H^{s_1,(p,q)})_{[\theta]}. \end{aligned} \quad (2.53)$$

The operators are bounded, which implies that the norms are equivalent. \square

Remark 2.4.10. Without assuming that $p_0 = p_1 = p$ and $q_0 = q_1 = q$, but with $q_0, q_1 < \infty$, we would have gotten

$$H^{s,(p,p \min(\frac{q_0}{p_0}, \frac{q_1}{p_1}))} \hookrightarrow (H^{s_0,(p_0,q_0)}, H^{s_1,(p_0,q_0)})_{[\theta]} \hookrightarrow H^{s,(p,q)} \quad (2.54)$$

where $s = (1 - \theta)s_0 + \theta s_1$, $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ and $\frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$. This gives an equality if $\frac{p_0}{q_0} = \frac{p_1}{q_1}$.

Remark 2.4.11. We would need an extension operator for the same result in a domain. Without it, we can still see that

$$H^{s,(p,q)}(\Omega) \hookrightarrow (L^{(p,q)}(\Omega), H^{1,(p,q)}(\Omega))_{[s]} \hookrightarrow (L^{(p,q)}(\Omega), W^{1,(p,q)}(\Omega))_{[s]} \quad (2.55)$$

for $0 < s < 1$.

2.5 The Cauchy operator

We start by proving some estimates for the Cauchy-operators \mathcal{C} , $\overline{\mathcal{C}}$. They are the right inverse operators for $\overline{\partial}$, ∂ respectively.

Definition 2.5.1. For $f \in \mathcal{E}'(\mathbb{C})$, we define

$$\mathcal{C}f = \frac{1}{\pi z} * f, \quad \overline{\mathcal{C}}f = \frac{1}{\pi \bar{z}} * f. \quad (2.56)$$

Whenever f is not compactly supported, we assume that it is in a certain normed space, in which it can be approximated by compactly supported distributions. Moreover we have to show that \mathcal{C} and $\overline{\mathcal{C}}$ are bounded operators in those cases.

Lemma 2.5.2. For $f \in L^{(2,1)}(\mathbb{C})$ we have

$$\mathcal{C} : L^{(2,1)}(\mathbb{C}) \rightarrow BC(\mathbb{C}), \quad \|\mathcal{C}f\|_{BC} \leq \frac{2}{\sqrt{\pi}} \|f\|_{L^{(2,1)}}, \quad (2.57)$$

and if $f \in L^1(\mathbb{C})$ we have

$$\mathcal{C} : L^1(\mathbb{C}) \rightarrow L^{(2,\infty)}(\mathbb{C}), \quad \|\mathcal{C}f\|_{L^{(2,\infty)}} \leq \frac{2}{\sqrt{\pi}} \|f\|_{L^1} \quad (2.58)$$

Proof. The Cauchy operators are well-defined here because functions in L^1 and $L^{(2,1)}$ can be approximated by test functions according to lemma 2.1.7. We shall use Young's inequality for Lorentz spaces. Note that the kernel $\frac{1}{\pi z} \in L^{(2,\infty)}(\mathbb{C})$ with norm $\|\frac{1}{\pi z}\|_{L^{(2,\infty)}(\mathbb{C})} = \frac{2}{\sqrt{\pi}}$.

The first estimate is given in [41] with proof as an exercise. Using Hölder's inequality and $\|k * f\|_{\infty} \leq \int_0^{\infty} f^*(t)k^*(t)dt \leq \int_0^{\infty} f^{**}(t)k^{**}(t)dt$ proven in corollary 1.8 of [41] we get it easily:

$$\begin{aligned} \|k * f\|_{L^{\infty}(\mathbb{C})} &\leq \int_0^{\infty} k^{**}(t)f^{**}(t)dt = \int_0^{\infty} t^{1/2}k^{**}(t) \cdot t^{-1/2}f^{**}(t)dt \\ &\leq \int_0^{\infty} t^{1/2}f^{**}(t)\frac{dt}{t} \sup_{t>0} t^{1/2}k^{**}(t) = \|f\|_{L^{(2,1)}(\Omega)} \|k\|_{L^{(2,\infty)}(\mathbb{C})}. \end{aligned} \quad (2.59)$$

Test functions map to continuous functions by according to Vekua, I.6.2 [47]. They are dense in $L^{(2,1)}(\Omega)$ because of theorem 2.3.4. Hence the estimate implies continuity.

The second estimate follows from Young's inequality and real interpolation. For $p = 1, \infty$ we have $\|k * f\|_p \leq \|k\|_p \|f\|_1$. By the properties of real interpolation between L^p -spaces, for example 7.26 in [1]⁴, we get

$$\|u\|_{L^{(2,\infty)}} = \|u\|_{(L^1, L^{\infty})_{\frac{1}{2}, \infty}}, \quad (2.60)$$

⁴They write $L^{p,q}$ for our space $L^{(p,q)}$.

which implies the claim. \square

Remark 2.5.3. The continuity of $\mathcal{C}\phi$, ϕ a test function, can be seen in another way. \mathcal{C} maps $L^{(2,1)}$ locally into the space $W^{1,(2,1)}$ of 2.3.4. This space can be embedded into BC .

We need to take care of boundary terms when considering the left inverses of $\partial, \bar{\partial}$. The next lemma will be used for that.

Lemma 2.5.4. *Let Ω be a bounded Lipschitz domain and $g \in L^1(\partial\Omega)$. Then*

$$\left\| \frac{1}{2\pi} \int_{\partial\Omega} \frac{g(z')}{z - z'} \eta(z') d\sigma(z') \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq \pi^{-\frac{3}{2}} \|g\|_{L^1(\partial\Omega)} \quad (2.61)$$

Remark 2.5.5. We write $\nu(z)$ for the normal vector at z pointing outwards the region of integration in \mathcal{C} . The complex one is $\eta(z) = \nu_1(z) + i\nu_2(z)$.

Proof. By Minkowski's integral inequality we see that

$$\left\| \frac{1}{2\pi} \int_{\partial\Omega} g(z') h(z - z') \eta(z') d\sigma(z') \right\|_{L^p(\mathbb{C})} \leq \frac{1}{2\pi} \|g\|_{L^1(\partial\Omega)} \|h\|_{L^p(\mathbb{C})} \quad (2.62)$$

for $1 \leq p \leq \infty$. By the properties of real interpolation between L^p -spaces, 7.26 in [1], we have

$$\|u\|_{L^{(2,\infty)}} = \|u\|_{(L^1, L^\infty)_{\frac{1}{2}, \infty}}. \quad (2.63)$$

Note that $z \mapsto \frac{1}{\pi z}$ is in $L^{(2,\infty)}(\mathbb{C})$ with norm $\left\| \frac{1}{\pi z} \right\|_{L^{(2,\infty)}(\mathbb{C})} = \frac{2}{\sqrt{\pi}}$. This implies the claim by real interpolation's interpolation property applied to the operator $h \mapsto \frac{1}{2\pi} \int_{\partial\Omega} g(z') h(\cdot - z') \eta(z') d\sigma(z')$. \square

Lemma 2.5.6. *Let Ω be a bounded Lipschitz domain in \mathbb{C} . The mapping $\bar{\partial}\mathcal{C}$ is the identity on $\mathcal{E}'(\mathbb{C})$ and $\partial\mathcal{C} : L^{(p,q)}(\mathbb{C}) \rightarrow L^{(p,q)}(\mathbb{C})$ for $1 < p < \infty$, $1 \leq q \leq \infty$. Let $f \in W^{1,1}(\Omega)$. Then all the terms of the following expression are in $L^{(2,\infty)}(\mathbb{C})$ and*

$$\mathcal{C}\chi_\Omega \bar{\partial}f(z) = \chi_\Omega f(z) + \frac{1}{2\pi} \int_{\partial\Omega} \eta(z') \frac{\text{Tr } f(z')}{z - z'} d\sigma(z'), \quad (2.64)$$

in the distribution sense. Here $\chi_\Omega g$ is always extended as zero outside of Ω .

Proof. The first claim follows from $\bar{\partial} \frac{1}{\pi z} = \delta_0$, which can be proven by taking the Fourier transform, or like in Vekua [47], chapter I, §4, equation (4.9). Note that $\partial\mathcal{C}$ is the well known Beurling operator Π , see for example Vekua [47] chapter I, §9. That book gives the result $\Pi : L^p \rightarrow L^p$ for $1 < p < \infty$. Using real interpolation we get it for $L^{(p,q)} \rightarrow L^{(p,q)}$, $1 < p < \infty$, $1 \leq q \leq \infty$.

We see that $\mathcal{C} : L^1(\mathbb{C}) \rightarrow L^{(2,\infty)}(\mathbb{C})$ by lemma 2.5.2. The boundary integral is also in $L^{(2,\infty)}(\mathbb{C})$ by lemma 2.5.4 since $\text{Tr} : W^{1,1}(\Omega) \rightarrow L^1(\partial\Omega)$. We have also $f \in W^{1,1}(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow L^{(2,\infty)}(\Omega)$ by Sobolev embedding, so $\chi_\Omega f \in L^{(2,\infty)}(\mathbb{C})$ when extended as zero.

Let $\phi \in C_0^\infty(\mathbb{C})$. Then

$$\begin{aligned} \langle \bar{\partial}(\chi_\Omega f), \phi \rangle &= -\langle \chi_\Omega f, \bar{\partial}\phi \rangle = -\int_\Omega f \bar{\partial}\phi dm \\ &= -\int_{\partial\Omega} \text{Tr} f \frac{\eta}{2} \phi d\sigma + \int_\Omega \bar{\partial}f \phi dm = \langle -\frac{\eta}{2} \text{Tr} f d\sigma_{\partial\Omega} + \chi_\Omega \bar{\partial}f, \phi \rangle \end{aligned} \quad (2.65)$$

by integrating by parts. See for example Nečas [37], theorem 3.1.2. After that, we get

$$\begin{aligned} \chi_\Omega f &= \delta_0 * (\chi_\Omega f) = \bar{\partial} \frac{1}{\pi z} * (\chi_\Omega f) = \frac{1}{\pi z} * \bar{\partial}(\chi_\Omega f) \\ &= \frac{1}{\pi z} * \left(-\frac{\eta}{2} \text{Tr} f d\sigma_{\partial\Omega} + \chi_\Omega \bar{\partial}f \right) = -\frac{1}{2\pi} \int_{\partial\Omega} \eta(z') \frac{\text{Tr} f(z')}{z - z'} d\sigma(z') + \mathcal{C} \chi_\Omega \bar{\partial}f, \end{aligned} \quad (2.66)$$

because $\chi_\Omega f \in \mathcal{E}'(\mathbb{C})$. □

3 Using the stationary phase method

3.1 The main term

Lemma 3.1.1 (Mean-value inequality). *Let $f : X \rightarrow \mathbb{C}$, $X \subset \mathbb{C}$ be convex, $f \in C^1(\overline{X})$. Then for all $x, y \in X$*

$$|f(x) - f(y)| \leq \begin{cases} \left\| \sqrt{|\partial_1 f|^2 + |\partial_2 f|^2} \right\|_{L^\infty(X)} |x - y| \\ \sqrt{2} \left\| \sqrt{|\partial f|^2 + |\bar{\partial} f|^2} \right\|_{L^\infty(X)} |x - y| \end{cases}. \quad (3.1)$$

Proof. By theorem 7.20 in Rudin [44]

$$\begin{aligned} |f(x) - f(y)| &= \left| \int_0^1 \frac{d}{dt} f(tx + (1-t)y) dt \right| \\ &= \left| \int_0^1 (\operatorname{Re} \nabla f \cdot (x_1 - y_1, x_2 - y_2) + i \operatorname{Im} \nabla f \cdot (x_1 - y_1, x_2 - y_2)) dt \right| \\ &\leq \int_0^1 \sqrt{|\operatorname{Re} \nabla f|^2 + |\operatorname{Im} \nabla f|^2} |x - y| dt \leq \|\nabla f\|_{L^\infty(X)} |x - y|. \end{aligned} \quad (3.2)$$

Note that $|\operatorname{Re} \nabla f|^2 + |\operatorname{Im} \nabla f|^2 = |\partial_1 f|^2 + |\partial_2 f|^2 = 2(|\partial f|^2 + |\bar{\partial} f|^2)$, from which the claim follows. \square

Lemma 3.1.2. *Let $s \geq 0$ and $\xi \in \mathbb{C}$. Then*

$$\left| 1 - e^{-i(\xi^2 + \bar{\xi}^2)} \right| \leq 2^{1+s/2} |\xi|^s. \quad (3.3)$$

Proof. The two cases to consider are $|\xi| < \frac{1}{\sqrt{2}}$ and $|\xi| \geq \frac{1}{\sqrt{2}}$. We use lemma 3.1.1 to get the first case.

$$\begin{aligned} &\sup_{|\xi| \leq 2^{-1/2}} \frac{\left| 1 - e^{-i(\xi^2 + \bar{\xi}^2)} \right|}{|\xi|^s} \\ &\leq \sup_{|\xi| \leq 2^{-1/2}} \sqrt{2} \left\| \sqrt{\left| -2i z e^{-i(z^2 + \bar{z}^2)} \right|^2 + \left| -2i \bar{z} e^{-i(z^2 + \bar{z}^2)} \right|^2} \right\|_{L^\infty(B(2^{-1/2}))} |\xi|^{1-s} \\ &\leq \sqrt{2} \cdot 2 \cdot \sqrt{2} \cdot 2^{-1/2} (2^{-1/2})^{1-s} \leq 2^{1+s/2} \end{aligned} \quad (3.4)$$

The second case follows because $\xi^2 + \bar{\xi}^2 \in \mathbb{R}$.

$$\sup_{|\xi| \geq 2^{-1/2}} \frac{\left| 1 - e^{-i(\xi^2 + \bar{\xi}^2)} \right|}{|\xi|^s} \leq \sup_{|\xi| \geq 2^{-1/2}} \frac{2}{|\xi|^s} = 2^{1+s/2} \quad (3.5)$$

\square

From now on we denote $R = (z - z_0)^2 + (\bar{z} - \bar{z}_0)^2$, where z_0 is a point in \mathbb{C} . The stationary phase method is based on the Fourier transform of a complex Gaussian. See lemma 7.2 for details.

Lemma 3.1.3 (Stationary phase). *Let $\tau > 0$ and $s \geq 0$. If $Q \in H^{s,2}(\mathbb{C})$ then*

$$\left\| Q - \frac{2\tau}{\pi} \int_{\mathbb{C}} e^{i\tau R} Q(z) dm(z) \right\|_{L^2(\mathbb{C}, z_0)} \leq 2\tau^{-s/2} \|Q\|_{H^{s,2}(\mathbb{C})}. \quad (3.6)$$

Proof. A direct calculations using the Fourier transform

$$\mathcal{F}\left(\frac{2\tau}{\pi} e^{i\tau(z^2 + \bar{z}^2)}\right)(\xi) = \frac{1}{2\pi} e^{-i\frac{\xi^2 + \bar{\xi}^2}{16\tau}} \quad (3.7)$$

of lemma 7.2 and lemma 3.1.2:

$$\begin{aligned} \left\| Q - \frac{2\tau}{\pi} \int_{\mathbb{C}} e^{i\tau R} Q(z) dm(z) \right\|_{L^2(\mathbb{C}, z_0)} &= \left\| \widehat{Q} - e^{-i\frac{\xi^2 + \bar{\xi}^2}{16\tau}} \widehat{Q} \right\|_{L^2(\mathbb{C})} \\ &\leq 4^{-s} \tau^{-s/2} \left\| \frac{\left| 1 - e^{-i\left(\left(\frac{\xi}{4\sqrt{\tau}}\right)^2 + \left(\frac{\bar{\xi}}{4\sqrt{\tau}}\right)^2\right)} \right|}{\left| \frac{\xi}{4\sqrt{\tau}} \right|^s} |\xi|^s \widehat{Q} \right\|_{L^2(\mathbb{C})} \\ &\leq 2^{1-3s/2} \tau^{-s/2} \left\| |\xi|^s \widehat{Q} \right\|_{L^2(\mathbb{C})} \leq 2\tau^{-s/2} \|Q\|_{H^{s,2}(\mathbb{C})}. \end{aligned} \quad (3.8)$$

□

3.2 Handling the error term

Since the potential q of our Schrödinger equation won't be zero, our solutions will have an error term. It has to be handled separately when using stationary phase. Note that R depends only on $z_0 - z$, but the integral involving the error will not be a convolution operator. That's because the error term r will depend on z_0 too. Nevertheless, we may prove an L^∞ estimate for this operator, and ignore whether r or Q would depend on z_0 .

If $Q, r \in W^{1,p}$ with $p > 2$, we could get the estimate

$$\left\| \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} Q r(z) dm(z) \right\|_2 \leq C_{\Omega} \|Q\|_{1,p} \sup_{z_0} \|r\|_{1,p} \quad (3.9)$$

for example as in [8], which follows ideas of Bukhgeim [3]. We could try to follow that reasoning in our case of $Q \in H^{1,(2,1)}$ and $r \in H^{1,(2,\infty)}$. It would

start like this

$$\begin{aligned} \int 2\tau i e^{i\tau R} Q r dm &= \int \frac{\partial e^{i\tau R}}{z_0 - z} Q r dm = \pi \chi_\Omega e^{i\tau R} Q r \\ &+ \frac{1}{2} \int_{\partial\Omega} \eta(z') \frac{e^{i\tau R(z'-z_0)} Q r(z')}{z_0 - z'} d\sigma(z') - \int \frac{e^{i\tau R}}{z_0 - z} \partial(Qr) dm \end{aligned} \quad (3.10)$$

where we have used lemma 2.5.6. It is true even when r depends on z_0 . But then we would have to estimate $r_{z_0}(z_0)$ in the first term. This is no problem when $r \in L^\infty$, which we will have. However, for reasons related to the limitations of the interpolation result of theorem 2.2.12, we may only use the $H^{1,(2,\infty)}$ -norm of r . This space does not embed into L^∞ . Hence, if we want to follow this path, we should analyze what happens in theorem 4.1.1 and corollary 4.1.5 as the outer variable approaches z_0 . In any case, that's not needed here. The final rate $\tau^{1-s/3}$ given by the next approach is as good as τ^{1-s} , because we are more interested in the case $s \rightarrow 0$.

Theorem 3.2.1. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $\tau > 0$, $0 < s < 1$ and $z_0 \in \mathbb{C}$. If $Q \in H^{s,(2,1)}(\Omega)$ and $r \in H^{s,(2,\infty)}(\Omega)$, then*

$$\left| \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} Q r(z) dm(z) \right| \leq C_\Omega \tau^{1-s/3} \|Q\|_{s,(2,1)} \|r\|_{s,(2,\infty)}. \quad (3.11)$$

Proof. The claim follows by using complex interpolation on the multilinear mapping

$$T : (Q, r) \mapsto \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} Q r(z) dm(z), \quad T : H^{s,(2,1)}(\Omega) \times H^{s,(2,\infty)}(\Omega) \rightarrow \mathbb{C}. \quad (3.12)$$

Consider the case corresponding to $s = 0$ first. By Hölder's inequality for Lorentz spaces, theorem 3.5 in [41], we get directly

$$\left| \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} Q r(z) dm(z) \right| \leq \frac{2\tau}{\pi} \|Qr\|_1 \leq C_\Omega \tau \|Q\|_{(2,1)} \|r\|_{(2,\infty)}. \quad (3.13)$$

Take $h \in C_0^\infty(\Omega)$ as given by lemma 7.10 to prove the other limiting case. We split the integral like this

$$\begin{aligned} &\int_\Omega \frac{2\tau}{\pi} e^{i\tau R} Q r(z) dm(z) \\ &= \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} (\bar{z} - \bar{z}_0) h Q r(z) dm(z) + \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} (1 - (\bar{z} - \bar{z}_0) h) Q r(z) dm(z) \\ &= \frac{1}{i\pi} \int_\Omega \bar{\partial} e^{i\tau R} h Q r(z) dm(z) + \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} (1 - (\bar{z} - \bar{z}_0) h) Q r(z) dm(z). \end{aligned} \quad (3.14)$$

Estimate the second term first. The generalized Hölder's inequality implies

$$\begin{aligned}
\left| \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (1 - (\bar{z} - \bar{z}_0)h) Qr(z) dm(z) \right| &\leq C_{\Omega} \tau \| (1 - (\bar{z} - \bar{z}_0)h) Qr \|_1 \\
&\leq C_{\Omega} \tau \| 1 - (\bar{z} - \bar{z}_0)h \|_{(2,1)} \| Qr \|_{(2,\infty)} \\
&\leq C_{\Omega} \tau \| 1 - (\bar{z} - \bar{z}_0)h \|_{(2,1)} \| Q \|_{\infty} \| r \|_{(2,\infty)} \\
&\leq C_{\Omega} \tau \| 1 - (\bar{z} - \bar{z}_0)h \|_{(2,1)} \| Q \|_{1,(2,1)} \| r \|_{1,(2,\infty)} \quad (3.15)
\end{aligned}$$

by the embedding $H^{1,(2,1)}(\Omega) \hookrightarrow W^{1,(2,1)}(\Omega) \hookrightarrow L^{\infty}(\Omega)$ of theorems 2.3.4 and 2.4.2. Integrate the first term by parts. Then

$$\begin{aligned}
\left| \int_{\Omega} \bar{\partial} e^{i\tau R} h Qr(z) dm(z) \right| &= \left| \int_{\Omega} e^{i\tau R} \bar{\partial} (h Qr)(z) dm(z) \right| \\
&\leq \| \bar{\partial} h \|_{(2,1)} \| Qr \|_{(2,\infty)} + \| h \|_{\infty} \left(\| \bar{\partial} Q \|_{(2,1)} \| r \|_{(2,\infty)} + \| Q \|_{(2,1)} \| \bar{\partial} r \|_{(2,\infty)} \right) \\
&\leq \| \bar{\partial} h \|_{(2,1)} \| Q \|_{\infty} \| r \|_{(2,\infty)} + 2 \| h \|_{\infty} \| Q \|_{1,(2,1)} \| r \|_{1,(2,\infty)} \\
&\leq C_{\Omega} \left(\| \bar{\partial} h \|_{(2,1)} + \| h \|_{\infty} \right) \| Q \|_{1,(2,1)} \| r \|_{1,(2,\infty)} \quad (3.16)
\end{aligned}$$

because of $H^{1,(2,1)}(\Omega) \hookrightarrow W^{1,(2,1)}(\Omega) \hookrightarrow L^{\infty}(\Omega)$. Corollary 7.10 gives

$$\tau \| 1 - (\bar{z} - \bar{z}_0)h(z) \|_{L^{(2,1)}(\Omega)} + \| h \|_{L^{\infty}(\Omega)} + \| \bar{\partial} h \|_{L^{(2,1)}(\Omega)} \leq C_{\Omega} \tau^{2/3}. \quad (3.17)$$

The claim follows by interpolation as $H^{s,(p,q)}(\Omega) \hookrightarrow (L^{(p,q)}(\Omega), H^{1,(p,q)}(\Omega))_{[s]}$. This can be seen as follows: Let $f \in H^{s,(p,q)}(\Omega)$ and take $g \in H^{s,(p,q)}(\mathbb{C})$ such that $g|_{\Omega} = f$ and $\|g\|_{H^{s,(p,q)}(\mathbb{C})} \leq 2 \|f\|_{H^{s,(p,q)}(\Omega)}$. Now

$$\begin{aligned}
\|f\|_{(L^{(p,q)}(\Omega), H^{1,(p,q)}(\Omega))_{[s]}} &= \|g|_{\Omega}\|_{(L^{(p,q)}(\Omega), H^{1,(p,q)}(\Omega))_{[s]}} \\
&\leq \|g\|_{(L^{(p,q)}(\mathbb{C}), H^{1,(p,q)}(\mathbb{C}))_{[s]}} = \|g\|_{H^{s,(p,q)}(\mathbb{C})} \leq 2 \|f\|_{H^{s,(p,q)}(\Omega)}. \quad (3.18)
\end{aligned}$$

The fact that $(\mathbb{C}, \mathbb{C})_{[s]} = \mathbb{C}$ is the last small missing piece of the proof. \square

4 Bukhgeim type solutions

4.1 A Carleman estimate

Remember that we write $R = (z - z_0)^2 + (\bar{z} - \bar{z}_0)^2$. The next theorem is the heart of this whole thesis. The primary goal is to have the right-hand side vanish as fast as possible as τ grows. This requirement makes us study the function spaces $H^{s,(p,q)}$, which were the basis of chapter 2.2. The rest is quite straightforward after the next theorem. Continue by proving estimates for $\mathcal{C}(e^{-i\tau R}\chi_\Omega \bar{\mathcal{C}}(e^{i\tau R}\chi_\Omega qf))$, use them to find solutions to $(\Delta + q)e^{i\tau(z-z_0)^2}f = 0$ and use stationary phase to estimate $\|q_1 - q_2\|$ using the boundary data. Only technical details prevent this from being trivial.

Theorem 4.1.1 (The main Carleman estimate). *Let $\Omega \subset \mathbb{C}$ bounded and Lipschitz, $z_0 \in \mathbb{C}$ and $\tau > 1$. If $a \in BC(\bar{\Omega})$ and $\bar{\partial}a \in L^{(2,1)}(\Omega)$ then $\mathcal{C}(e^{-i\tau R}\chi_\Omega a) \in L^{(2,\infty)}(\mathbb{C}) \cap BC(\mathbb{C})$ and we have the norm estimates*

$$\begin{aligned} \|\mathcal{C}(e^{-i\tau R}\chi_\Omega a)\|_{L^{(2,\infty)}(\mathbb{C})} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \left(\|\bar{\partial}a\|_{L^{(2,1)}(\Omega)} + \|a\|_{BC(\bar{\Omega})} \right), \\ \|\mathcal{C}(e^{-i\tau R}\chi_\Omega a)\|_{BC(\mathbb{C})} &\leq C_\Omega \tau^{-1/3} \left(\|\bar{\partial}a\|_{L^{(2,1)}(\Omega)} + \|a\|_{BC(\bar{\Omega})} \right). \end{aligned} \quad (4.1)$$

Proof. The mapping properties of $\mathcal{C}(e^{i\tau R}\chi_\Omega \cdot)$ follow from the corresponding mapping properties of each term, all given by the same lemmas that imply the norm estimates used here.

Let $h \in W^{1,1}(\Omega)$. Now

$$\begin{aligned} \mathcal{C}(e^{i\tau R}\chi_\Omega a) &= \mathcal{C}(e^{i\tau R}(1 - (\cdot - \bar{z}_0)h)\chi_\Omega a) + \mathcal{C}(e^{i\tau R}(\cdot - \bar{z}_0)h\chi_\Omega a) \\ &= \mathcal{C}(e^{i\tau R}(1 - (\cdot - \bar{z}_0)h)\chi_\Omega a) + \frac{1}{2i\tau} \mathcal{C}(\bar{\partial}(e^{i\tau R})h\chi_\Omega a) \end{aligned} \quad (4.2)$$

We get

$$\begin{aligned} \mathcal{C}(\chi_\Omega \bar{\partial}(e^{i\tau R})ha) &= \chi_\Omega e^{i\tau R} h(z)a \\ &\quad + \frac{1}{2\pi} \int_{\partial\Omega} \eta(z') \frac{e^{i\tau R} \text{Tr } h(z')a(z')}{z - z'} d\sigma(z') \\ &\quad - \mathcal{C}(\chi_\Omega e^{i\tau R} \bar{\partial}ha) - \mathcal{C}(\chi_\Omega e^{i\tau R} h \bar{\partial}a) \end{aligned} \quad (4.3)$$

by lemma 2.5.6 because $e^{i\tau R}ha \in W^{1,1}(\Omega)$.

Take h as in lemma 7.4 to prove the first estimate. By lemma 2.5.2

$$\begin{aligned} & \left\| \mathcal{C}(e^{i\tau R}(1 - (\bar{\cdot} - \bar{z}_0)h)\chi_\Omega a) \right\|_{L^{(2,\infty)}(\mathbb{C})} \\ & \leq \frac{2}{\sqrt{\pi}} \left\| (1 - (\bar{\cdot} - \bar{z}_0)h)a \right\|_{L^1(\Omega)} \\ & \leq \frac{2}{\sqrt{\pi}} \left\| 1 - (\bar{\cdot} - \bar{z}_0)h \right\|_{L^1(\Omega)} \|a\|_{L^\infty(\Omega)}. \end{aligned} \quad (4.4)$$

Next

$$\left\| \chi_\Omega e^{i\tau R} h(z)a \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq \|h\|_{L^{(2,\infty)}(\Omega)} \|a\|_{L^\infty(\Omega)}, \quad (4.5)$$

and by lemma 2.5.4

$$\begin{aligned} & \left\| \frac{1}{2\pi} \int_{\partial\Omega} \eta(z') \frac{e^{i\tau R} \text{Tr} h(z')a(z')}{z - z'} d\sigma(z') \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq \pi^{-\frac{3}{2}} \left\| \text{Tr} ha \right\|_{L^1(\partial\Omega)} \\ & \leq \pi^{-\frac{3}{2}} \|h\|_{L^1(\partial\Omega)} \|a\|_{L^\infty(\partial\Omega)} \leq \pi^{-\frac{3}{2}} T_\Omega \|h\|_{W^{1,1}(\Omega)} \|a\|_{BC(\bar{\Omega})}, \end{aligned} \quad (4.6)$$

where $T_\Omega < \infty$ is the norm of the trace mapping $\text{Tr} : W^{1,1}(\Omega) \rightarrow L^1(\partial\Omega)$. Again, by lemma 2.5.2, we get

$$\left\| \mathcal{C}(\chi_\Omega e^{i\tau R} \bar{\partial} ha) \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq \frac{2}{\sqrt{\pi}} \left\| \bar{\partial} h \right\|_{L^1(\Omega)} \|a\|_{L^\infty(\Omega)}, \quad (4.7)$$

and according to [41] we have we have $\|ab\|_1 \leq \|a\|_{(2,\infty)} \|b\|_{(2,1)}$. Hence

$$\left\| \mathcal{C}(\chi_\Omega e^{i\tau R} h \bar{\partial} a) \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq \frac{2}{\sqrt{\pi}} \|h\|_{L^{(2,\infty)}(\Omega)} \left\| \bar{\partial} a \right\|_{L^{(2,1)}(\Omega)}. \quad (4.8)$$

Combining everything, using the Sobolev embedding $W^{1,1} \hookrightarrow L^2 \hookrightarrow L^{(2,\infty)}$ and the inequality

$$\tau \left\| 1 - (\bar{z} - \bar{z}_0)h \right\|_{L^1(\Omega)} + \|h\|_{W^{1,1}(\Omega)} \leq C_\Omega(1 + \ln \tau) \quad (4.9)$$

of lemma 7.4 gives the first estimate.

For the second one, let $h \in C_0^\infty(\Omega)$ be as in corollary 7.10. Then $\chi_\Omega h = h$. Continue from (4.2) and (4.3). The boundary term vanishes,

$$\begin{aligned} & \left\| \mathcal{C}(e^{i\tau R}(1 - (\bar{\cdot} - \bar{z}_0)h)a) \right\|_{BC(\mathbb{C})} \\ & \leq \frac{2}{\sqrt{\pi}} \left\| (1 - (\bar{\cdot} - \bar{z}_0)h)a \right\|_{L^{(2,1)}(\Omega)} \\ & \leq \frac{2}{\sqrt{\pi}} \left\| 1 - (\bar{\cdot} - \bar{z}_0)h \right\|_{L^{(2,1)}(\Omega)} \|a\|_{L^\infty(\Omega)}, \end{aligned} \quad (4.10)$$

$$\|\chi_\Omega e^{i\tau R} h(z) a\|_{BC(\mathbb{C})} \leq \|h\|_{BC(\Omega)} \|a\|_{BC(\Omega)}, \quad (4.11)$$

$$\|\mathcal{C}(e^{i\tau R} \bar{\partial} h a)\|_{BC(\mathbb{C})} \leq \frac{2}{\sqrt{\pi}} \|\bar{\partial} h\|_{L^{(2,1)}(\Omega)} \|a\|_{BC(\bar{\Omega})}, \quad (4.12)$$

and

$$\|\mathcal{C}(e^{i\tau R} h \bar{\partial} a)\|_{BC(\mathbb{C})} \leq \frac{2}{\sqrt{\pi}} \|h\|_{L^\infty(\Omega)} \|\bar{\partial} a\|_{L^{(2,1)}(\Omega)}. \quad (4.13)$$

The estimate

$$\tau \|1 - (\bar{z} - \bar{z}_0) h(z)\|_{L^{(2,1)}(\Omega)} + \|h\|_{L^\infty(\Omega)} + \|\bar{\partial} h\|_{L^{(2,1)}(\Omega)} \leq C_\Omega \tau^{2/3} \quad (4.14)$$

of corollary 7.10 gives the rest. \square

Remark 4.1.2. Dependency and measurability on z_0 will be taken care of later. It will turn out that the operator is continuous with respect to it. Actually, we would like to have the dependence in $L^2(\mathbb{C})$ or $L^{(2,\infty)}(\mathbb{C})$ for non-compactly supported potentials.

Remark 4.1.3. It seems possible to get the map $W^{1,p} \rightarrow L^p$ with exponent $\tau^{-\frac{1}{2} - \frac{1}{p}}$ (no logarithm) when $p > 2$. But it would require Bloch spaces, *BMO* in a domain and a related interpolation identity. See section 6.2.

Remark 4.1.4. This is a Carleman estimate for $\bar{\partial}$. Write $r = e^{i\tau R} \mathcal{C}(e^{-i\tau R} a)$ and consider all the derivatives in $\mathcal{D}'(\Omega)$, where $\chi_\Omega \equiv 1$. Now

$$a = e^{i\tau R} \bar{\partial} e^{-i\tau R} r = e^{i\tau(\bar{z} - \bar{z}_0)^2} \bar{\partial} e^{-i\tau(\bar{z} - \bar{z}_0)^2} r. \quad (4.15)$$

Hence we have the following Carleman estimates:

$$\begin{aligned} \|r\|_{(2,\infty)} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \left\| e^{i\tau(\bar{z} - \bar{z}_0)^2} \bar{\partial} e^{-i\tau(\bar{z} - \bar{z}_0)^2} r \right\|_{1,(2,1)} \\ \|r\|_{C^0} &\leq C_\Omega \tau^{-1/3} \left\| e^{i\tau(\bar{z} - \bar{z}_0)^2} \bar{\partial} e^{-i\tau(\bar{z} - \bar{z}_0)^2} r \right\|_{1,(2,1)} \end{aligned} \quad (4.16)$$

Corollary 4.1.5. *Let Ω be a bounded Lipschitz domain, $z_0 \in \mathbb{C}$ and $\tau > 1$. Let $q \in L^{(2,1)}(\Omega)$ or $q \in W^{1,(2,1)}(\Omega)$. Write $Sf = \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f))$. Then*

$$\begin{aligned} \|Sf\|_{L^{(2,\infty)}(\mathbb{C})} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|q\|_{L^{(2,1)}(\Omega)} \|f\|_{L^\infty(\Omega)}, \\ \|Sf\|_{H^{1,(2,\infty)}(\mathbb{C})} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|q\|_{W^{1,(2,1)}(\Omega)} \|f\|_{W^{1,(2,1)}(\Omega)}, \\ \|Sf\|_{BC(\mathbb{C})} &\leq C_\Omega \tau^{-1/3} \|q\|_{L^{(2,1)}(\Omega)} \|f\|_{L^\infty(\Omega)}, \\ \|Sf\|_{H^{1,(2,1)}(\Omega)} &\leq C_\Omega \tau^{-1/3} \|q\|_{W^{1,(2,1)}(\Omega)} \|f\|_{W^{1,(2,1)}(\Omega)}, \end{aligned} \quad (4.17)$$

with corresponding mapping properties.

Proof. The mapping properties follow from those of theorem 4.1.1 and lemma 2.5.2. We will need the facts that $\partial\mathcal{C}, \bar{\partial}\mathcal{C} : L^{(p,q)}(\mathbb{C}) \rightarrow L^{(p,q)}(\mathbb{C})$ and that $\partial\bar{\mathcal{C}}, \bar{\partial}\mathcal{C}$ are the identity in $\mathcal{E}'(\mathbb{C})$. These are given by lemma 2.5.6. We can then proceed. If there's no domain in the index of the norm, then that norm is taken in Ω . Else it is taken where denoted. Now

$$\begin{aligned} & \left\| \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{L^{(2,\infty)}(\mathbb{C})} \\ & \leq C_\Omega \tau^{-1} (1 + \ln \tau) \left(\left\| \bar{\partial}\mathcal{C}(e^{i\tau R} \chi_\Omega qf) \right\|_{(2,1)} + \left\| \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{BC(\bar{\Omega})} \right) \\ & \leq C_\Omega \tau^{-1} (1 + \ln \tau) \|q\|_{(2,1)} \|f\|_\infty, \quad (4.18) \end{aligned}$$

and using the second estimate of theorem 4.1.1 instead of the first one, we get

$$\left\| \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{BC(\mathbb{C})} \leq C_\Omega \tau^{-1/3} \|q\|_{(2,1)} \|f\|_\infty. \quad (4.19)$$

Consider the cases where there's a derivative on the left hand side. We will need the identity $W^{1,(2,\infty)}(\mathbb{C}) = H^{1,(2,\infty)}(\mathbb{C})$ of theorem 2.4.2 and the continuous embedding $L^{(2,\infty)}(\Omega) \hookrightarrow L^1(\Omega)$. It is true because $m(\Omega) < \infty$. Then

$$\begin{aligned} & \left\| \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{W^{1,(2,\infty)}(\mathbb{C})} \leq C_\Omega \left(\left\| \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_1 \right. \\ & \left. + \left\| \partial\mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{L^{(2,\infty)}(\mathbb{C})} + \left\| e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{L^{(2,\infty)}(\mathbb{C})} \right) \\ & \leq C_\Omega \left\| \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{L^{(2,\infty)}(\mathbb{C})} \leq C_\Omega \tau^{-1} (1 + \ln \tau) \|q\|_{1,(2,1)} \|f\|_{1,(2,1)}. \quad (4.20) \end{aligned}$$

The last estimate requires a technical trick since we haven't shown that $H^{1,(2,1)}(\Omega) = W^{1,(2,1)}(\Omega)$. Let $\phi \in C_0^\infty(\mathbb{C})$ be constant 1 on $B(0, R) \supset \Omega$. Note that $BC(X) \hookrightarrow L^{(2,1)}(X)$ whenever $m(X) < \infty$, so

$$\begin{aligned} & \left\| \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{1,(2,1)} \leq \left\| \phi \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{H^{1,(2,1)}(\mathbb{C})} \\ & \leq C \left\| \phi \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{W^{1,(2,1)}(\mathbb{C})} \\ & \leq C_{\Omega,\phi} \left(\left\| \mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{BC(\text{supp } \phi)} \right. \\ & \left. + \left\| \partial\mathcal{C}(e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf)) \right\|_{L^{(2,1)}(\mathbb{C})} + \left\| e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{L^{(2,1)}(\mathbb{C})} \right) \\ & \leq C_{\Omega,\phi} \left\| \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{(2,1)} \leq C_{\Omega,\phi} \left\| \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf) \right\|_{BC(\Omega)} \\ & \leq C_{\Omega,\phi} \tau^{-1/3} \|q\|_{1,(2,1)} \|f\|_{1,(2,1)}, \quad (4.21) \end{aligned}$$

by theorems 4.1.1 and 2.3.4. The cut-off function ϕ may be chosen based only on Ω , so $C_{\Omega,\phi}$ depends only on the domain. \square

Remark 4.1.6. If $m(\Omega) = \infty$, then it would make sense to define spaces $\tilde{W}^{1,(2,1)} = \{f \mid f \in BC, \nabla f \in L^{(2,1)}\}$ to avoid the use of the embedding $BC \subset L^{(2,1)}$. But then, on the other hand, we run into problems finding which space X maps $\mathcal{C} : X \rightarrow L^{(2,1)}$.

Remark 4.1.7. The estimate can be called a Carleman estimate for the Laplacian. Write $r = \mathcal{C}(e^{-i\tau R} \overline{\mathcal{C}}(e^{i\tau R} qf))$ and consider all the derivatives in $\mathcal{D}'(\Omega)$ where $\chi_\Omega \equiv 1$. Now

$$qf = e^{-i\tau R} \partial e^{i\tau R} \overline{\partial} r = \frac{1}{4} e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r. \quad (4.22)$$

Hence we have the following Carleman estimates:

$$\begin{aligned} \|r\|_{(2,\infty)} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{(2,1)}, \\ \|r\|_{1,(2,\infty)} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{1,(2,1)}, \\ \|r\|_{BC} &\leq C_\Omega \tau^{-1/3} \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{(2,1)}, \\ \|r\|_{1,(2,1)} &\leq C_\Omega \tau^{-1/3} \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{1,(2,1)}. \end{aligned} \quad (4.23)$$

Definition 4.1.8. Let $\Omega \subset \mathbb{C}$ be a Lipschitz domain. For $0 < s < 1$ we write $M^s(\Omega) = (BC(\overline{\Omega}), H^{1,(2,1)}(\Omega))_{[s]}$.

Remark 4.1.9. This is a well defined Banach space since both BC and $W^{1,(2,1)}$ are Banach spaces that can be embedded into $BC(\overline{\Omega})$ by theorem 2.3.4.

Corollary 4.1.10. Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $\tau > 1$, $z_0 \in \mathbb{C}$ and $0 < s < 1$. Let $q \in H^{s,(2,1)}(\Omega)$. Then

$$\begin{aligned} \|\mathcal{C}(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf))\|_{H^{s,(2,\infty)}(\Omega)} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|q\|_{H^{s,(2,1)}(\Omega)} \|f\|_{M^s(\Omega)} \\ \|\mathcal{C}(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf))\|_{M^s(\Omega)} &\leq C_\Omega \tau^{-1/3} \|q\|_{H^{s,(2,1)}(\Omega)} \|f\|_{M^s(\Omega)} \end{aligned} \quad (4.24)$$

with similar mapping properties. The map $z_0 \mapsto \mathcal{C}(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega qf_{z_0}))$ is in

$$BC(\overline{\Omega}, W^{1,2}(\Omega) \cap H^{s,(2,\infty)}(\Omega) \cap M^s(\Omega)) \quad (4.25)$$

for each $f : (z, z_0) \mapsto f_{z_0}(z)$ bounded and continuous $\overline{\Omega} \times \overline{\Omega} \rightarrow \mathbb{C}$.

Proof. All the norms with W on the right-hand side of corollary 4.1.5 can be estimated above by norms with H . This follows from theorem 2.4.2. The second estimate follows directly from the definition of $M^s(\Omega)$, corollary 4.1.5, the inclusion $H^{s,(2,1)}(\Omega) \hookrightarrow (L^{(2,1)}(\Omega), H^{1,(2,1)}(\Omega))_{[s]}$ and the multilinear interpolation property of complex interpolation.

The first one requires a bit more careful considerations because we haven't shown that $(L^{(2,\infty)}(\Omega), H^{1,(2,\infty)}(\Omega))_{[s]} \hookrightarrow H^{s,(2,\infty)}(\Omega)$. Interpolation tells us that the operator maps $M^s(\Omega) \rightarrow (L^{(2,\infty)}(\mathbb{C}), H^{1,(2,\infty)}(\mathbb{C}))_{[s]} = H^{s,(2,\infty)}(\mathbb{C})$. This is because of theorem 2.4.2. Now

$$\|g|_{\Omega}\|_{H^{s,(2,\infty)}(\Omega)} = \inf_{\substack{G \in H^{s,(2,\infty)}(\mathbb{C}) \\ G|_{\Omega} = g|_{\Omega}}} \|G\|_{H^{s,(2,\infty)}(\mathbb{C})} \leq \|g\|_{H^{s,(2,\infty)}(\mathbb{C})} \quad (4.26)$$

for any $g \in H^{s,(2,\infty)}(\mathbb{C})$. The estimate follows.

It's left to prove pointwise continuity $\bar{\Omega} \rightarrow H^{1,(2,1)}(\Omega)$. This is because of the chains of bounded mappings

$$\begin{aligned} H^{1,(2,1)}(\Omega) &\hookrightarrow W^{1,(2,1)}(\Omega) \hookrightarrow W^{1,2}(\Omega) \\ H^{1,(2,1)}(\Omega) &\hookrightarrow BC(\bar{\Omega}) \cap H^{1,(2,1)}(\Omega) \hookrightarrow M^s(\Omega), \\ H^{1,(2,1)}(\Omega) &\hookrightarrow H^{1,(2,\infty)}(\Omega) \hookrightarrow H^{s,(2,\infty)}(\Omega), \end{aligned} \quad (4.27)$$

which follow from $L^{(2,1)}(\mathbb{C}) \hookrightarrow L^{(2,\infty)}(\mathbb{C})$ and $H^{1,(2,\infty)}(\mathbb{C}) \hookrightarrow H^{s,(2,\infty)}(\mathbb{C})$.

Let $z_0, z_1 \in \bar{\Omega}$ and write $f_j = f_{z_j}$ and $R_j = (z - z_j)^2 + (\bar{z} - \bar{z}_j)^2$, where z is the variable being operated on. Then, proceed as in the proof of corollary 4.1.5. Take $\phi \in C_0^\infty(\mathbb{C})$ such that $\phi \equiv 1$ on $B(0, R) \supset \Omega$. Then

$$\begin{aligned} &\|\mathcal{C}(e^{-i\tau R_0} \chi_{\Omega} \bar{\mathcal{C}}(e^{i\tau R_0} \chi_{\Omega} q f_0)) - \mathcal{C}(e^{-i\tau R_1} \chi_{\Omega} \bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1))\|_{H^{1,(2,1)}(\Omega)} \\ &\leq \|\phi \mathcal{C}(e^{-i\tau R_0} \chi_{\Omega} \bar{\mathcal{C}}(e^{i\tau R_0} \chi_{\Omega} q f_0)) - \phi \mathcal{C}(e^{-i\tau R_1} \chi_{\Omega} \bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1))\|_{W^{1,(2,1)}(\mathbb{C})} \\ &\leq \|\phi \mathcal{C}(e^{-i\tau R_0} \chi_{\Omega} (\bar{\mathcal{C}}(e^{i\tau R_0} \chi_{\Omega} q f_0) - \bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1)))\|_{W^{1,(2,1)}(\mathbb{C})} \\ &\quad + \|\phi \mathcal{C}((e^{-i\tau R_0} - e^{-i\tau R_1}) \chi_{\Omega} \bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1))\|_{W^{1,(2,1)}(\mathbb{C})}. \end{aligned} \quad (4.28)$$

Next note that $\phi \mathcal{C} \chi_{\Omega}, \phi \bar{\mathcal{C}} \chi_{\Omega} : L^\infty(\Omega) \rightarrow W^{1,(2,1)}(\mathbb{C})$ by the boundedness of Ω and lemmas 2.5.2 and 2.5.6. The first term in (4.28) can be estimates as

$$\begin{aligned} \dots &\leq C_{\Omega, \phi} \|\bar{\mathcal{C}}((e^{i\tau R_0} f_0 - e^{i\tau R_1} f_1) \chi_{\Omega} q)\|_{L^\infty(\Omega)} \\ &\leq C_{\Omega, \phi} \|e^{i\tau R_0} f_0 - e^{i\tau R_1} f_1\|_{L^\infty(\Omega)} \|q\|_{L^{(2,1)}(\Omega)} \longrightarrow 0 \end{aligned} \quad (4.29)$$

as $z_0 \rightarrow z_1$ by the uniform continuity of $e^{i\tau R} f$ in $\bar{\Omega} \times \bar{\Omega}$. The second term is handled quite similarly. We continue from (4.28)

$$\begin{aligned} \dots &\leq C_{\Omega, \phi} \|(e^{-i\tau R_0} - e^{-i\tau R_1}) \bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1)\|_{L^\infty(\Omega)} \\ &\leq C_{\Omega, \phi} \|e^{-i\tau R_0} - e^{-i\tau R_1}\|_{L^\infty(\Omega)} \|\bar{\mathcal{C}}(e^{i\tau R_1} \chi_{\Omega} q f_1)\|_{L^\infty(\Omega)} \\ &\leq C_{\Omega, \phi} \|e^{-i\tau R_0} - e^{-i\tau R_1}\|_{L^\infty(\Omega)} \|q\|_{L^{(2,1)}(\Omega)} \|f_1\|_{L^\infty(\Omega)} \longrightarrow 0 \end{aligned} \quad (4.30)$$

as $z_0 \rightarrow z_1$ because $\bar{\Omega}$ is compact and $e^{-i\tau R}$ is continuous. \square

Remark 4.1.11. The equality $(L^{(2,\infty)}(\Omega), H^{1,(2,\infty)}(\Omega))_{[s]} = H^{s,(2,\infty)}(\Omega)$ would follow from the existence of a *strong extension operator* E mapping both $E : L^{(2,\infty)}(\Omega) \rightarrow L^{(2,\infty)}(\mathbb{C})$ and $E : H^{1,(2,\infty)}(\Omega) \rightarrow H^{1,(2,\infty)}(\mathbb{C})$.

Remark 4.1.12. As in an earlier remark, we get the usual form of the Carleman estimates by writing $r = \mathcal{C}(e^{-i\tau R} \overline{\mathcal{C}}(e^{i\tau R} qf))$:

$$\begin{aligned} \|r\|_{s,(2,\infty)} &\leq C_\Omega \tau^{-1} (1 + \ln \tau) \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{s,(2,1)} \\ \|r\|_{M^s} &\leq C_\Omega \tau^{-1/3} \|e^{-i\tau(z-z_0)^2} \Delta e^{i\tau(z-z_0)^2} r\|_{s,(2,1)} \end{aligned} \quad (4.31)$$

4.2 Bukhgeim's oscillating solutions

We can now show that $(\Delta + q)u = 0$ has Bukhgeim's oscillating solutions $u = e^{i\tau(z-z_0)^2} f$, with $f - 1$ small enough. See [3] for the original article. Smallness can not be proven at the same time as existence. Instead, we fetch f from M^s and show that its norm is small in $H^{s,(2,\infty)}$. It is basically a boot-strapping argument.

Theorem 4.2.1. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $0 < s < 1$, $q \in H^{s,(2,1)}(\Omega)$ and⁵ $\tau > \max\{1, (C_\Omega \|q\|_{s,(2,1)})^3\}$. Then, for each $z_0 \in \overline{\Omega}$, there is a unique $f_{z_0} \in M^s(\Omega)$ such that*

$$f_{z_0} = 1 - \frac{1}{4} \mathcal{C} \left(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f_{z_0}) \right). \quad (4.32)$$

The map $z_0 \mapsto f_{z_0}$ is continuous $\overline{\Omega} \rightarrow W^{1,2}(\Omega) \cap H^{s,(2,\infty)}(\Omega) \cap M^s(\Omega)$ and we have the norm estimates

$$\begin{aligned} \|f_{z_0} - 1\|_{s,(2,\infty)} &\leq C_{\Omega,s} \tau^{-1} (1 + \ln \tau) \|q\|_{s,(2,1)}, \\ \|f_{z_0}\|_{1,2} &\leq C_{\Omega,s} (1 + \|q\|_{(2,1)}). \end{aligned} \quad (4.33)$$

Proof. For $\tau > \max\{1, (C_\Omega \|q\|_{s,(2,1)})^3\}$ and $z_0 \in \overline{\Omega}$ define

$$T_{z_0} : f \mapsto 1 - \frac{1}{4} \mathcal{C} \left(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f) \right). \quad (4.34)$$

We have $T_{z_0} : M^s(\Omega) \rightarrow M^s(\Omega)$ by corollary 4.1.10 and

$$\begin{aligned} \|T_{z_0} f - T_{z_0} f'\|_{M^s(\Omega)} &= \frac{1}{4} \left\| \mathcal{C} \left(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega q (f - f')) \right) \right\|_{M^s(\Omega)} \\ &\leq \frac{1}{4} C_\Omega \tau^{-1/3} \|q\|_{H^{s,(2,1)}(\Omega)} \|f - f'\|_{M^s(\Omega)}, \end{aligned} \quad (4.35)$$

so T is a contraction in $M^s(\Omega)$. Moreover, corollary 4.1.10 implies that the map $z_0 \mapsto T_{z_0} f$ is continuous. Banach's fixed point theorem shows that there

⁵Here C_Ω is the constant in the second estimate of corollary 4.1.10.

is a unique $f_{z_0} \in M^s(\Omega)$ satisfying (4.32) and it depends continuously on z_0 . See for example [45], VII §1, theorem 3.

The solution $(z_0, z) \mapsto f_{z_0}(z)$ is bounded and continuous $\bar{\Omega} \times \bar{\Omega} \rightarrow \mathbb{C}$ because $M^s(\Omega) \hookrightarrow BC(\bar{\Omega})$. Hence corollary 4.1.10 gives us the continuity $\bar{\Omega} \rightarrow W^{1,2}(\Omega) \cap H^{s,(2,\infty)}(\Omega) \cap M^s(\Omega)$. Then use the first inequality of the same corollary to get

$$\|f_{z_0} - 1\|_{s,(2,\infty)} \leq \frac{1}{4}C_\Omega\tau^{-1}(1 + \ln \tau) \|q\|_{s,(2,1)} \|f_{z_0}\|_{M^s}. \quad (4.36)$$

We have $\bar{\mathcal{C}} : L^{(2,1)} \rightarrow BC(\mathbb{C})$ and $\mathcal{C}\chi_\Omega : BC(\bar{\Omega}) \hookrightarrow L^{(2,1)}(\Omega) \rightarrow W^{1,(2,1)}(\Omega) \hookrightarrow W^{1,2}(\Omega)$ by the boundedness of Ω , the fact that $L^{(2,1)} \hookrightarrow L^2$ and lemmas 2.5.2 and 2.5.6. This implies

$$\|f_{z_0}\|_{1,2} \leq C_\Omega(1 + \|q\|_{(2,1)}) \|f\|_{M^s}. \quad (4.37)$$

The claims follow since $\|T_{z_0}\|_{M^s \rightarrow M^s} \leq \frac{1}{4}$, so $\|f_{z_0}\|_{M^s} \leq \|1\|_{M^s} + \frac{1}{4}\|f_{z_0}\|_{M^s}$, which implies $\|f_{z_0}\|_{M^s} \leq \frac{4}{3}\|1\|_{M^s}$. \square

Corollary 4.2.2. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $0 < s < 1$, $q \in H^{s,(2,1)}(\Omega)$ and $\tau > \max\{1, (C_\Omega \|q\|_{s,(2,1)})^3\}$. Let $f_{z_0} \in M^s(\Omega)$ be as in theorem 4.2.1. Then*

$$\Delta \left(e^{i\tau(z-z_0)^2} f_{z_0} \right) + q e^{i\tau(z-z_0)^2} f_{z_0} = 0 \quad (4.38)$$

in $\mathcal{D}'(\Omega)$.

Proof. Note that $\chi_\Omega \equiv 1$ in $\mathcal{D}'(\Omega)$ and keep in mind that

$$f_{z_0} = 1 - \frac{1}{4}\mathcal{C}(e^{-i\tau R}\chi_\Omega\bar{\mathcal{C}}(e^{i\tau R}\chi_\Omega q f_{z_0})). \quad (4.39)$$

We get

$$\begin{aligned} \bar{\partial} \left(e^{i\tau(z-z_0)^2} f_{z_0} \right) &= e^{i\tau(z-z_0)^2} \bar{\partial} f_{z_0} = -\frac{1}{4} e^{i\tau(z-z_0)^2} e^{-i\tau R} \chi_\Omega \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f_{z_0}) \\ &= -\frac{1}{4} e^{-i\tau(\bar{z}-\bar{z}_0)^2} \bar{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f_{z_0}). \end{aligned} \quad (4.40)$$

because $\bar{\partial}\bar{\mathcal{C}} = \partial\bar{\mathcal{C}} = \text{Id}$ in $\mathcal{E}'(\mathbb{C})$ by lemma 2.5.6. Now

$$\begin{aligned} \Delta \left(e^{i\tau(z-z_0)^2} f_{z_0} \right) &= 4\partial\bar{\partial} \left(e^{i\tau(z-z_0)^2} f_{z_0} \right) = -e^{-i\tau(\bar{z}-\bar{z}_0)^2} e^{i\tau R} \chi_\Omega q f_{z_0} \\ &= -q e^{i\tau(z-z_0)^2} f_{z_0}. \end{aligned} \quad (4.41)$$

\square

5 The problem setting

5.1 Hadamard's criteria and the DN operator

We will define what the well-posedness of the direct problem means, what is the Dirichlet-Neumann operator and then make an extension allowing us to solve the inverse problem for potentials that do not necessarily give well posed direct problems.

The Dirichlet-Neumann operator is inherently related to the trace mapping Tr from a function space on a domain to its boundary. It is well known that $\text{Tr} : H^s(\Omega) \rightarrow H^{s-\frac{1}{2}}(\partial\Omega)$ for suitable values of the smoothness index s when Ω is a bounded Lipschitz domain. We want to avoid the trouble of defining $H^s(\partial\Omega)$ when Ω is not smooth, so instead we take the “boundary values” as equivalence classes in $W^{1,2}(\Omega)$. More specifically⁶, they are in $W^{1,2}(\Omega)/W_0^{1,2}(\Omega)$ with norm

$$\|u\|_{W/W_0} = \inf_{\phi \in W_0^{1,2}(\Omega)} \|u + \phi\|_{W^{1,2}(\Omega)}. \quad (5.1)$$

Using this space instead of the spaces $H^s(\partial\Omega)$ is allowed. See for example Gagliardo [18] and Ding [16]. They imply

$$\begin{array}{ccc} H^{1/2}(\partial\Omega) & \xrightarrow{\text{Id}} & H^{1/2}(\partial\Omega) \\ & \searrow \mathcal{E} & \nearrow \text{Tr} \\ & & W^{1,2}(\Omega)/W_0^{1,2}(\Omega) \end{array} \quad (5.2)$$

where Tr, \mathcal{E} are bounded linear mappings. Equation (2.9) of [18] gives linearity for \mathcal{E} . Note that here and from now on we write $W/W_0 = W^{1,2}(\Omega)/W_0^{1,2}(\Omega)$.

The Dirichlet-Neumann map is a way to model boundary measurements. It makes sense to require the differential equation to behave nicely enough to be a physical model. One such class of problems is those satisfying Hadamard's three criteria [24]: 1) the problem must have a solution, 2) the solution must be unique, and 3) the solution should depend continuously on the data. In such case, the problem is said to be well-posed. The reasons for these criteria is also a mathematical one. We can prove mapping properties for the Dirichlet-Neumann operator if the potential gives a well-posed problem.

⁶To be more exact: given $u \in W^{1,2}(\Omega)$ there is a unique equivalence class $[u] \in W/W_0$ such that $u \in [u]$. Moreover $u + \phi \in [u]$ for all $\phi \in W_0^{1,2}(\Omega)$.

Definition 5.1.1. Let $\Omega \subset \mathbb{C}$ be open and q measurable. Then *the direct problem is well-posed* if there is $C < \infty$ such that for any $u \in W^{1,2}(\Omega)/W_0^{1,2}(\Omega)$ we have

1. there is $U \in W^{1,2}(\Omega)$ such that $\Delta U + qU = 0$, $U - u \in W_0^{1,2}(\Omega)$,
2. this U is unique
3. $u \mapsto U$ is linear and bounded $\|U\|_{W^{1,2}(\Omega)} \leq C \|u\|_{W/W_0}$

Definition 5.1.2. Let $\Omega \subset \mathbb{C}$ be open and q measurable such that the direct problem is well-posed. Then we define the *Dirichlet-Neumann operator* Λ_q as follows. For $u \in W/W_0$ we define $\Lambda_q u$ by

$$(\Lambda_q u, v) = \int_{\Omega} (-\nabla U \cdot \nabla V + qUV) dm, \quad v \in W/W_0, \quad (5.3)$$

for any $U, V \in W^{1,2}(\Omega)$ such that $U - u, V - v \in W_0^{1,2}(\Omega)$ and $\Delta U + qU = 0$.

Lemma 5.1.3. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain and $q \in L^a(\Omega)$ with $a > 1$. Then the Dirichlet-Neumann map is a well defined bounded linear operator mapping W/W_0 to its dual. Moreover it satisfies*

$$(\Lambda_q u, v) = (\Lambda_q v, u), \quad u, v \in W/W_0. \quad (5.4)$$

Proof. We start by showing that the choice of U, V in definition 5.1.2 doesn't matter. First of all, U exists and is unique on the right-hand side of (5.3) by the well-posedness of the direct problem 5.1.1. Assume that $V, V' \in W^{1,2}(\Omega)$ satisfy $V - v, V' - v \in W_0^{1,2}(\Omega)$. Then

$$\begin{aligned} & \int_{\Omega} -\nabla U \cdot \nabla V + qUV dm - \int_{\Omega} -\nabla U \cdot \nabla V' + qUV' dm \\ &= \int_{\Omega} -\nabla U \cdot \nabla (V - V') + qU(V - V') dm \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} -\nabla U \cdot \nabla \phi_n + qU \phi_n dm = \lim_{n \rightarrow \infty} \langle \Delta U + qU, \phi \rangle = 0 \end{aligned} \quad (5.5)$$

where $\phi_n \in C_0^\infty(\Omega)$ such that $\|\phi_n - (V - V')\|_{1,2} \rightarrow 0$. This sequence exists because $V - V' = V - v - (V' - v) \in W_0^{1,2}(\Omega)$. Hence (5.3) is well defined.

The mapping properties follow next. We have $W^{1,2}(\Omega) \hookrightarrow L^{\frac{2a}{a-1}}(\Omega)$ by Sobolev embedding, e.g. theorem 4.12 in [1]. Now

$$1 = \frac{1}{a} + \frac{1}{\frac{2a}{a-1}} + \frac{1}{\frac{2a}{a-1}}, \quad (5.6)$$

so

$$\begin{aligned} |(\Lambda_q u, v)| &\leq \|\nabla U\|_2 \|\nabla V\|_2 + \|q\|_a \|U\|_{\frac{2a}{a-1}} \|V\|_{\frac{2a}{a-1}} \\ &\leq C_{a,\Omega} (1 + \|q\|_a) \|U\|_{1,2} \|V\|_{1,2}. \end{aligned} \quad (5.7)$$

Then, take the infimum over $V \in W^{1,2}(\Omega)$, $V-v \in W_0^{1,2}(\Omega)$ and use condition number three of the well-posedness 5.1.1 on U to get

$$|(\Lambda_q u, v)| \leq C_{a,\Omega,q} \|u\|_{W/W_0} \|v\|_{W/W_0}. \quad (5.8)$$

To prove the last formula let $u, v \in W/W_0$ and take $U, V \in W^{1,2}(\Omega)$ such that $U-u, V-v \in W_0^{1,2}(\Omega)$, $\Delta U + qU = \Delta V + qV = 0$. These exist by the well-posedness 5.1.1. Hence

$$(\Lambda_q u, v) = \int_{\Omega} -\nabla U \cdot \nabla V + qUV dm = \int_{\Omega} -\nabla V \cdot \nabla U + qVU dm = (\Lambda_q v, u). \quad (5.9)$$

□

Remark 5.1.4. Λ_q is a well defined linear operator mapping W/W_0 to its dual. Hence the linear combinations of such operators are also well defined. In particular

$$\|\Lambda_{q_1} - \Lambda_{q_2}\| = \sup \{ |((\Lambda_{q_1} - \Lambda_{q_2})u, v)| \mid u, v \in W/W_0, \|u\| = \|v\| = 1 \}. \quad (5.10)$$

5.2 Cauchy data

We would like to extend the notion of the Dirichlet-Neumann map to cases where the direct problem is not well-posed. One such way is to make use of the *Cauchy data*

$$C_q = \{(\text{Tr } u, \partial_\nu u) \mid u \in W^{2,2}(\Omega), \Delta u + qu = 0\}, \quad (5.11)$$

but this would further require three more definitions and related properties: the trace-operator Tr , the normal derivative ∂_ν , and a way to measure the distance of two Cauchy data C_{q_1} and C_{q_2} .

We are not too interested in the direct problem, so instead we will just extend the notion of $\Lambda_{q_1} - \Lambda_{q_2}$ to situations where the Dirichlet problem $\Delta U + qU = 0$, $U-u \in W_0^{1,2}(\Omega)$ does not have a unique solution. It is based on the well known Alessandrini's identity $\int U_1(q_1 - q_2)U_2 dm = ((\Lambda_{q_1} - \Lambda_{q_2})u_1, u_2)$ for solutions $\Delta U_j + q_j U_j = 0$, $U_j - u_j \in W_0^{1,2}(\Omega)$. We shall prove it here too. But first, the generalization.

Definition 5.2.1. Let $\Omega \subset \mathbb{C}$ be open and q_1, q_2 measurable. Then the distance between the boundary data, $d(C_{q_1}, C_{q_2})$, is

$$d(C_{q_1}, C_{q_2}) = \sup \left\{ \left| \int_{\Omega} U(q_1 - q_2)V dm \right| \mid U, V \in W^{1,2}(\Omega), \right. \\ \left. \|U\| = \|V\| = 1, \Delta U + q_1 U = \Delta V + q_2 V = 0 \right\} \quad (5.12)$$

Remark 5.2.2. It is possible to acquire this data purely using knowledge from the boundary, at least part of it when $U, V \in W^{2,2}$. This *Alessandrini's identity* follows from Green's formula:

$$\int_{\Omega} U(q_1 - q_2)V dm = \int_{\Omega} -V\Delta U + U\Delta V dm = \int_{\partial\Omega} \text{Tr } U\partial_{\nu}V - \text{Tr } V\partial_{\nu}U d\sigma \quad (5.13)$$

If the reader is concerned about the lack of smoothness in the solutions of the definition of $d(C_{q_1}, C_{q_2})$, then note the following: the only solution U, V that matter for solving the inverse problem are Bukhgeim's oscillating ones. Namely those that were constructed in corollary 4.2.2. It is not hard to see that they are in $W^{2,2}$ and that their $W^{2,2}$ norms also grow exponentially.

Lemma 5.2.3. Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain and $q_1, q_2 \in L^a(\Omega)$, $a > 1$. Then $d(C_{q_1}, C_{q_2}) \leq C_{a,\Omega} \|q_1 - q_2\|_a$.

Proof. Use the Sobolev embedding $W^{1,2}(\Omega) \hookrightarrow L^{\frac{2a}{a-1}}(\Omega)$, for example in theorem 4.12, [1]. Then

$$\left| \int_{\Omega} U(q_1 - q_2)V dm \right| \leq \|U\|_{\frac{2a}{a-1}} \|q_1 - q_2\|_a \|V\|_{\frac{2a}{a-1}} \\ \leq C_{a,\Omega} \|q_1 - q_2\|_a \|U\|_{1,2} \|V\|_{1,2}. \quad (5.14)$$

□

Theorem 5.2.4. Let $\Omega \subset \mathbb{C}$ be bounded and Lipschitz, and $q_1, q_2 \in L^a(\Omega)$, $a > 1$, be such that the direct problem is well-posed. Then

$$d(C_{q_1}, C_{q_2}) \leq \|\Lambda_{q_1} - \Lambda_{q_2}\|. \quad (5.15)$$

Proof. Let $u, v \in W/W_0$. Take $U, V \in W^{1,2}(\Omega)$ such that $\Delta U + q_1 U = 0$, $U - u \in W_0^{1,2}(\Omega)$ and similarly with V, v . These exist by the well-posedness of the direct problems 5.1.1. Because the Dirichlet-Neumann map in (5.3) is

well-defined, we have

$$\begin{aligned}
((\Lambda_{q_1} - \Lambda_{q_2})u, v) &= (\Lambda_{q_1}u, v) - (\Lambda_{q_2}u, v) = (\Lambda_{q_1}u, v) - (\Lambda_{q_2}v, u) \\
&= \int_{\Omega} -\nabla U \cdot \nabla V + q_1 UV + \nabla V \cdot \nabla U - q_2 VU dm \\
&= \int_{\Omega} U(q_1 - q_2)V dm \quad (5.16)
\end{aligned}$$

by lemma 5.1.3.

Let \mathcal{P} be the canonical projection map $W^{1,2}(\Omega) \rightarrow W/W_0$. It has operator norm at most one by definition. Now choose $u = \mathcal{P}U$ and $v = \mathcal{P}V$ to get

$$\begin{aligned}
\left| \int_{\Omega} U(q_1 - q_2)V dm \right| &= |((\Lambda_{q_1} - \Lambda_{q_2})u, v)| \\
&\leq \|\Lambda_{q_1} - \Lambda_{q_2}\| \|u\|_{W/W_0} \|v\|_{W/W_0} \leq \|\Lambda_{q_1} - \Lambda_{q_2}\| \|U\|_{1,2} \|V\|_{1,2}. \quad (5.17)
\end{aligned}$$

Taking the supremum over $U, V \in W^{1,2}(\Omega)$, $\Delta U + q_1 U = \Delta V + q_2 V = 0$, $\|U\|_{1,2} = \|V\|_{1,2} = 1$ gives the result. \square

5.3 Uniqueness and stability for the inverse problem

A technical lemma first.

Lemma 5.3.1. *Let $\Omega \subset \mathbb{C}$ be bounded and Lipschitz, $f_1, f_2 \in BC(\overline{\Omega}, W^{1,2}(\Omega))$ and $\tau > 0$. Let $u_1(z) = e^{i\tau(z-z_0)^2} f_1(z)$ and $u_2(z) = e^{i\tau(\bar{z}-\bar{z}_0)^2} f_2(z)$ for $z_0 \in \overline{\Omega}$. Then*

$$\|u_j\|_{BC(W^{1,2})} \leq e^{C_{\Omega}\tau} \|f_j\|_{BC(W^{1,2})} \quad (5.18)$$

for some positive real C_{Ω} depending only on Ω .

Proof. This follows from the elementary facts $|z - z_0|, |\bar{z} - \bar{z}_0| \leq \text{diam}(\Omega)$, $\tau \leq e^{\tau}$ and $\left| \partial e^{i\tau(z-z_0)^2} \right|, \left| \bar{\partial} e^{i\tau(\bar{z}-\bar{z}_0)^2} \right| \leq 2\tau \text{diam}(\Omega) e^{\text{diam}(\Omega)^2\tau}$. \square

We are now ready to solve the inverse problem. The big goal of inverse problems for partial differential equations is to deduce the values of the coefficients inside the domain using only data from the boundary. We do not get that far, that is, we do not have a reconstruction formula. Instead, we show uniqueness and logarithmic stability. It means that if there are two potentials q_1 and q_2 which are distance ϵ apart, then their corresponding boundary data must be roughly at least $e^{-\epsilon^{-1}}$ apart. This is not much, but it is not possible to get a better modulus of continuity. Mandache showed that the inverse problem is inherently *ill-posed* [33].

We remind the general flow of the proof. For a more detailed reminder see section 1.3. We start by using stationary phase

$$\int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \longrightarrow q_1 - q_2 \quad (5.19)$$

as $\tau \rightarrow \infty$. The integral can be approximated by a term $\int_{\Omega} u_1(q_1 - q_2)u_2$, like in the definition of $d(C_{q_1}, C_{q_2})$, because of the special form of our solutions. All in all

$$\begin{aligned} q_1 - q_2 &= \left(q_1 - q_2 - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right) + \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \\ &= \left(q_1 - q_2 - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right) + \frac{2\tau}{\pi} \int_{\Omega} u^{(1)}(q_1 - q_2) u^{(2)} dm \\ &\quad + \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} (q_1 - q_2) (1 - f_1 f_2) dm \end{aligned} \quad (5.20)$$

where $u^{(1)} = e^{i\tau(z-z_0)^2} f_1$ and $u^{(2)} = e^{i\tau(\bar{z}-\bar{z}_0)^2} f_2$ are the solutions given by theorem 4.2.1 and corollary 4.2.2. The first term will be estimated by lemma 3.1.3, the second one by theorem 5.2.4, and the last one by theorem 3.2.1.

Theorem 5.3.2. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $M > 0$ and $0 < s < \frac{1}{2}$. Then there is a positive real number C such that if $q_j \in H^{s,(2,1)}(\Omega)$ and $\|q_j\|_{s,(2,1)} \leq M$ then*

$$\|q_1 - q_2\|_{L^{(2,\infty)}(\Omega)} \leq C (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4}. \quad (5.21)$$

In particular, we have uniqueness and stability for potentials in $H^{s,p}(\Omega)$ with $s > 0$, $p > 2$.

Proof. Denote $Q = q_1 - q_2$ and $R = (z - z_0)^2 + (\bar{z} - \bar{z}_0)^2$ for $z, z_0 \in \mathbb{C}$. Remember that z is the variable being operated on first. Let $\tau > 1$. By the triangle inequality

$$\begin{aligned} &\|q_1 - q_2\|_{L^{(2,\infty)}(\Omega)} \\ &\leq \left\| Q - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} Q dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)} + \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)}, \end{aligned} \quad (5.22)$$

where the norms are taken with respect to z_0 . We will use stationary phase next. Extend Q by zero outside of Ω to get Q_0 . It is in $H^{s,2}(\mathbb{C})$ by 3.5 in

[13] with norm estimate $\|Q_0\|_{H^{s,2}(\mathbb{C})} \leq C_{\Omega,s} \|Q\|_{H^{s,2}(\Omega)}$. We get

$$\begin{aligned} \left\| Q - \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q dm(z) \right\|_{L^2(\Omega, z_0)} &= \left\| Q_0 - \frac{2\tau}{\pi} \int_{\mathbb{C}} e^{i\tau R} Q_0 dm(z) \right\|_{L^2(\mathbb{C}, z_0)} \\ &\leq C_s \tau^{-s/2} \|Q_0\|_{H^{s,2}(\mathbb{C})} \leq C_{\Omega,s} \tau^{-s/2} \|Q\|_{H^{s,2}(\Omega)} \\ &\leq C_{\Omega,s} \tau^{-s/2} \|Q\|_{s,(2,1)} \leq C_{\Omega,M,s} \tau^{-s/4} \end{aligned} \quad (5.23)$$

by theorem 3.1.3. We estimated $\tau^{-s/2} \leq \tau^{-s/4}$ for later purposes. We also used the embedding $L^2 \hookrightarrow L^{(2,\infty)}$.

Consider the second term next. Take $\tau_0 = \max\{1, (C_{\Omega}M)^3\}$ as in theorem 4.2.1 and let $\tau = \frac{1}{2B_{\Omega}} \ln d(C_{q_1}, C_{q_2})^{-1}$, where $B_{\Omega} = 1 + 2C_{\Omega}$ with the C_{Ω} of lemma 5.3.1. We have $\tau > \tau_0$ if

$$d(C_{q_1}, C_{q_2}) < e^{-2B_{\Omega}\tau_0}. \quad (5.24)$$

Assume that $d(C_{q_1}, C_{q_2}) < \min\{e^{-1}, e^{-2B_{\Omega}\tau_0}\}$ for now. The other case will be taken care of in the end of the proof. Note that τ_0 grows with M .

Theorem 4.2.1 (the sign of i does not matter) gives the existence of $f^{(1)}, f^{(2)} \in BC(\bar{\Omega}, M^s(\Omega) \cap W^{1,2}(\Omega))$ such that we have

$$\begin{cases} f^{(1)} = 1 - \frac{1}{4} \mathcal{C}(e^{-i\tau R} \chi_{\Omega} \overline{\mathcal{C}}(e^{i\tau R} \chi_{\Omega} q_1 f^{(1)})), \\ f^{(2)} = 1 - \frac{1}{4} \overline{\mathcal{C}}(e^{-i\tau R} \chi_{\Omega} \mathcal{C}(e^{i\tau R} \chi_{\Omega} q_2 f^{(2)})), \end{cases} \quad (5.25)$$

for all $z_0 \in \bar{\Omega}$, and

$$\begin{cases} \sup_{z_0} \|f^{(j)} - 1\|_{H^{s,(2,\infty)}(\Omega)} \leq C_{\Omega,M,s} \tau^{-1} (1 + \ln \tau), \\ \sup_{z_0} \|f^{(j)}\|_{W^{1,2}(\Omega)} \leq C_{\Omega,M,s} < \infty. \end{cases} \quad (5.26)$$

Denote

$$\begin{cases} u_{z_0}^{(1)}(z) = e^{i\tau(z-z_0)^2} f^{(1)}(z_0, z), \\ u_{z_0}^{(2)}(z) = e^{i\tau(\bar{z}-\bar{z}_0)^2} f^{(2)}(z_0, z). \end{cases} \quad (5.27)$$

Now they satisfy $u_{z_0}^{(j)} \in BC(\bar{\Omega}, M^s(\Omega) \cap W^{1,2}(\Omega))$ and $\Delta u_{z_0}^{(j)} + q_j u_{z_0}^{(j)} = 0$ for all z_0 by corollary 4.2.2. Moreover, we have

$$\sup_{z_0} \|u_{z_0}^{(j)}\|_{W^{1,2}(\Omega)} \leq C_{\Omega,M,s} e^{C_{\Omega}\tau} \quad (5.28)$$

by lemma 5.3.1. Now, by the triangle inequality,

$$\begin{aligned} \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)} &\leq \left\| \frac{2\tau}{\pi} \int_{\Omega} u_{z_0}^{(1)}(q_1 - q_2) u_{z_0}^{(2)} dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)} \\ &+ \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q (f^{(1)} f^{(2)} - 1) dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)}. \end{aligned} \quad (5.29)$$

Use definition 5.2.1 and the equation (5.28) to get

$$\begin{aligned} & \left\| \frac{2\tau}{\pi} \int_{\Omega} u_{z_0}^{(1)}(q_1 - q_2) u_{z_0}^{(2)} dm(z) \right\|_{(2,\infty)} \\ & \leq 2\pi^{-1} \tau d(C_{q_1}, C_{q_2}) \sup_{z_0} \|u_{z_0}^{(1)}\|_{W^{1,2}(\Omega)} \sup_{z_0} \|u_{z_0}^{(2)}\|_{W^{1,2}(\Omega)} \\ & \leq C_{\Omega, M, s} d(C_{q_1}, C_{q_2}) e^{B_{\Omega}\tau}. \end{aligned} \quad (5.30)$$

For the second term, we need to show that $(f^{(1)}f^{(2)} - 1) \in H^{s,(2,\infty)}(\Omega)$ and that there is no problems with measurability for the $L^{(2,\infty)}(\Omega)$ norm. Notice that $f^{(1)}f^{(2)} - 1 = (f^{(1)} - 1)(f^{(2)} - 1) + f^{(1)} - 1 + f^{(2)} - 1$.

We will show that $H^{s,(2,\infty)}(\Omega)$ is stable with respect to multiplication by $f^{(2)} - 1$, as the latter can be extended to a function $G \in BC(\mathbb{C})$ with $\nabla G \in L^4(\mathbb{C})$. Let $\phi \in C_0^\infty(\mathbb{C})$ be such that $\phi \equiv 1$ on $\bar{\Omega}$ and it vanishes outside $B_r = B(0, r) \supset \bar{\Omega}$. Then set

$$G = -\frac{1}{4}\phi\bar{\mathcal{C}}\left(e^{-i\tau R}\chi_{\Omega}\mathcal{C}\left(e^{i\tau R}\chi_{\Omega}q_2f^{(2)}\right)\right). \quad (5.31)$$

Now $G|_{\Omega} = f^{(2)} - 1$, and according to corollary 4.1.5 we have $G \in BC(\mathbb{C})$. Next

$$\begin{aligned} \|\nabla G\|_{L^4(\mathbb{C})} & \leq \|\nabla\phi\|_{L^4(\mathbb{C})} \frac{1}{4} \|\bar{\mathcal{C}}\left(e^{-i\tau R}\chi_{\Omega}\mathcal{C}\left(e^{i\tau R}\chi_{\Omega}q_2f^{(2)}\right)\right)\|_{L^\infty(\mathbb{C})} \\ & \quad + C \|\phi\|_{L^\infty(\mathbb{C})} \|\mathcal{C}\left(e^{i\tau R}\chi_{\Omega}q_2f^{(2)}\right)\|_{L^4(\Omega)} \\ & \leq C_{\phi, \Omega} \|\mathcal{C}\left(e^{i\tau R}\chi_{\Omega}q_2f^{(2)}\right)\|_{L^\infty(\Omega)} \leq C_{\Omega} \|q_2f^{(2)}\|_{L^{(2,1)}(\Omega)} \\ & \leq C_{\Omega} \|q_2\|_{H^{s,(2,1)}(\Omega)} \|f^{(2)}\|_{M^s(\Omega)} \leq C_{\Omega, M, s} \end{aligned} \quad (5.32)$$

because $\nabla\mathcal{C}\chi_{\Omega} : L^4(\mathbb{C}) \rightarrow L^4(\mathbb{C})$ by lemma 2.5.6.

It is easy to see that $\|GF\|_{(2,\infty)} \leq \|G\|_{\infty} \|F\|_{(2,\infty)} \leq C_{\Omega, M, s} \|F\|_{(2,\infty)}$ for $F \in L^{(2,\infty)}(\mathbb{C})$. Take $F \in H^{1,(2,\infty)}(\mathbb{C})$. Now

$$\begin{aligned} \|GF\|_{1,(2,\infty)} & \leq C \left(\|FG\|_{(2,\infty)} + \|G\nabla F\|_{(2,\infty)} + \|F\nabla G\|_{(2,\infty)} \right) \\ & \leq C \left(\|G\|_{\infty} \|F\|_{(2,\infty)} + \|G\|_{\infty} \|\nabla F\|_{(2,\infty)} + \|\nabla G\|_{L^4(B_r)} \|F\|_{L^4(B_r)} \right). \end{aligned} \quad (5.33)$$

Note that $F|_{B_r} \in W^{1,(2,\infty)}(B_r) \hookrightarrow W^{1,\frac{4}{3}}(B_r) \hookrightarrow L^4(B_r)$ by Sobolev embedding (e.g. 4.12 in [1]). Moreover $\|F|_{B_r}\|_{H^{1,(2,\infty)}(B_r)} \leq C_r \|F\|_{H^{1,(2,\infty)}(\mathbb{C})}$. Hence

$$\|GF\|_{1,(2,\infty)} \leq C_{r,\Omega} (\|G\|_{\infty} + \|\nabla G\|_4) \|F\|_{1,(2,\infty)}. \quad (5.34)$$

Interpolation implies that multiplying by G is stable in $H^{s,(2,\infty)}(\mathbb{C})$, with norm increasing by at most $C_{\Omega, M, s}$ since r can be chosen based on Ω only.

Now take an extension F of $f^{(1)} - 1$ such that $\|F\|_{\mathbb{C}} \leq 2\|f^{(1)} - 1\|_{\Omega}$. This is possible by the definition of $H^{s,(2,\infty)}(\Omega)$. Then

$$\begin{aligned} \|(f^{(1)} - 1)(f^{(2)} - 1)\|_{H^{s,(2,\infty)}(\Omega)} &\leq \|FG\|_{H^{s,(2,\infty)}(\mathbb{C})} \\ &\leq C_{\Omega,M,s} \|F\|_{H^{s,(2,\infty)}(\mathbb{C})} \leq C_{\Omega,M,s} \|f^{(1)} - 1\|_{H^{s,(2,\infty)}(\Omega)}. \end{aligned} \quad (5.35)$$

This shows that $f^{(1)}f^{(2)} - 1 \in H^{s,(2,\infty)}(\Omega)$. It has the norm bound

$$\|f^{(1)}f^{(2)} - 1\|_{s,(2,\infty)} \leq C_{\Omega,M,s}\tau^{-1}(1 + \ln \tau) \quad (5.36)$$

by the previous deductions and theorem 4.2.1. Measurability with respect to z_0 is no problem since $f^{(1)}, f^{(2)} \in BC(W^{1,2})$.

Next, use theorem 3.2.1 to continue from (5.29). We have

$$\begin{aligned} &\left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q(f^{(1)}f^{(2)} - 1) dm(z) \right\|_{L^{(2,\infty)}(\Omega, z_0)} \\ &\leq C_{\Omega} \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} Q(f^{(1)}f^{(2)} - 1) dm(z) \right\|_{L^{\infty}(\Omega, z_0)} \\ &\leq C_{\Omega,s}\tau^{1-s/3} \|Q\|_{s,(2,1)} \sup_{z_0 \in \Omega} \|f^{(1)}f^{(2)} - 1\|_{s,(2,\infty)} \\ &\leq C_{\Omega,M,s}\tau^{-s/3}(1 + \ln \tau) \leq C_{\Omega,M,s}\tau^{-s/4}, \end{aligned} \quad (5.37)$$

since $\ln \tau \leq C_s\tau^{s/3-s/4}$.

We can combine all the terms now, namely those from equations (5.23), (5.30) and (5.37). Remember the choice of $\tau = \frac{1}{2B_{\Omega}} \ln d(C_{q_1}, C_{q_2})^{-1}$ and that $d(C_{q_1}, C_{q_2}) < e^{-1}$. Note that $\ln a \leq \frac{1}{b}a^b$ for $a, b > 0$, so $x^{1/2} \leq C_s(\ln \frac{1}{x})^{-s/4}$ when $0 < x < e^{-1}$. Now

$$\begin{aligned} \|q_1 - q_2\|_{(2,\infty)} &\leq C_{\Omega,M,s} (\tau^{-s/4} + d(C_{q_1}, C_{q_2})e^{B_{\Omega}\tau}) \\ &= C_{\Omega,M,s} ((\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4} + d(C_{q_1}, C_{q_2})^{1/2}) \\ &\leq C_{\Omega,M,s} (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4}. \end{aligned} \quad (5.38)$$

What if $d(C_{q_1}, C_{q_2}) \geq \min\{e^{-1}, e^{-2B_{\Omega}\tau_0}\}$? Then we would get directly $(\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4} \geq (\ln \max\{e, e^{2B_{\Omega}\tau_0}\})^{-s/4}$, so

$$\begin{aligned} \|q_1 - q_2\| &\leq 2M \leq \frac{2M}{(\ln \max\{e, e^{2B_{\Omega}\tau_0}\})^{-s/4}} (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4} \\ &= C_{\Omega,M,s} (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4} \end{aligned} \quad (5.39)$$

The boundedness of Ω implies that $L^p(\Omega) \hookrightarrow L^{(2,1)}(\Omega)$. Hence the claim is true for potentials in $H^{s,p}(\Omega)$. \square

Remark 5.3.3. If q_1 and q_2 would give well-posed direct problems as in definition 5.1.1, then theorem 5.2.4 shows that

$$\|q_1 - q_2\| \leq C_{\Omega, M, s} (\ln d(C_{q_1}, C_{q_2})^{-1})^{-s/4} \leq C_{\Omega, M, s} (\ln \|\Lambda_{q_1} - \Lambda_{q_2}\|^{-1})^{-s/4}. \quad (5.40)$$

6 Future work

6.1 Function space properties of $H^{s,(p,q)}(\Omega)$ and $W^{s,(p,q)}(\Omega)$

In section 2 we showed that $H^{k,(p,q)}(\mathbb{R}^n) = W^{k,(p,q)}(\mathbb{R}^n)$ and commented on the embedding $H^{k,(p,q)}(\Omega) \hookrightarrow W^{k,(p,q)}(\Omega)$. There is still these two very likely equalities left to prove:

1. $H^{k,(p,q)}(\Omega) = W^{k,(p,q)}(\Omega)$
2. $(L^{(p,q)}(\Omega), H^{1,(p,q)}(\Omega))_{[s]} = (L^{(p,q)}(\Omega), W^{1,(p,q)}(\Omega))_{[s]} = H^{s,(p,q)}(\Omega)$

These would follow directly from the existence of a strong extension operator mapping both $E : L^{(p,q)}(\Omega) \rightarrow L^{(p,q)}(\mathbb{R}^n)$ and $E : W^{k,(p,q)}(\Omega) \rightarrow W^{k,(p,q)}(\mathbb{R}^n)$. Assume that such a map exists. Consider the restriction $Rf = f|_{\Omega}$. Now $R \circ E = \text{Id}$ and

$$\begin{array}{ccc} W^{k,(p,q)}(\Omega) & \xrightarrow{\text{Id}} & W^{k,(p,q)}(\Omega) \\ & \searrow E & \nearrow R \\ & W^{k,(p,q)}(\mathbb{R}^n) & \\ & = & \\ & H^{k,(p,q)}(\mathbb{R}^n) & \\ & & \searrow R \\ & & H^{k,(p,q)}(\Omega) \end{array} \quad (6.1)$$

Hence $\text{Id} = R \circ E : W^{k,(p,q)}(\Omega) \hookrightarrow H^{k,(p,q)}(\Omega)$, so they are the same space. Moreover, since the operator E is a strong extension operator, the commutative diagram is preserved in interpolation. Hence $(L^{(p,q)}(\Omega), W^{1,(p,q)}(\Omega))_{[s]}$ is a retract of $H^{s,(p,q)}(\mathbb{R}^n)$, and so the first space is a subspace of $H^{s,(p,q)}(\Omega)$. The other direction follows by the definition of $H^{s,(p,q)}(\Omega)$, because it is the smallest space X for which $R : H^{s,(p,q)}(\mathbb{R}^n) \rightarrow X$.

After that these are some other things to consider: Sobolev embedding theorems, trace theorems, and any other subject in most standard books on Sobolev spaces.

6.2 Doing it in $W^{s,p}$

An earlier version of this manuscript [9] focused solely on potentials in $W^{s,p}(\Omega)$ with $p > 2$, $s > 0$. If we compare that text to this one, there is a trade-off. In the old one we had simpler function spaces, but much more parameters in the estimates. This is because of the boundary integral operator

$$T : f \mapsto \frac{1}{2\pi} \int_{\partial\Omega} \frac{f(z')}{z - z'} d\sigma(z'). \quad (6.2)$$

We couldn't quite prove that it would map $L^p(\partial\Omega) \rightarrow L^p(\Omega)$ for $2 < p < \infty$. We now know that it maps $L^1(\partial\Omega) \rightarrow L^{(2,\infty)}(\Omega) \hookrightarrow L^1(\Omega)$. Using *Bloch* spaces and their relation to $BMO(\Omega)$, as in [34] or [42], we get

$$T : L^\infty(\partial\Omega) \rightarrow BMO(\Omega). \quad (6.3)$$

Hence the result would follow by an interpolation result like

$$(L^1(\Omega), BMO(\Omega))_{[1/p]} = L^p(\Omega) \quad \text{or} \quad (L^1(\Omega), BMO(\Omega))_{1/p,p} = L^p(\Omega). \quad (6.4)$$

The latter is true when Ω has a *regular Vitali family*. See [25], [43].

There seems to be an easier way however. Note that in 4.1.1, 4.1.5 and 4.1.10 we always studied operators of the form

$$a \mapsto \mathcal{C}(e^{-i\tau R} \chi_\Omega a) \quad \text{and} \quad f \mapsto \mathcal{C}(e^{-i\tau R} \chi_\Omega \overline{\mathcal{C}}(e^{i\tau R} \chi_\Omega q f)). \quad (6.5)$$

If we work in the conventional Sobolev spaces $W^{1,p}$, we have extension operators $E : W^{1,p}(\Omega) \rightarrow W_c^{1,p}(\mathbb{C})$. Hence we may take a smooth test-function ϕ , which is constant on Ω , and consider the operators

$$a \mapsto \mathcal{C}(e^{-i\tau R} \phi a) \quad \text{and} \quad f \mapsto \mathcal{C}(e^{-i\tau R} \phi \overline{\mathcal{C}}(e^{i\tau R} \phi E q f)) \quad (6.6)$$

instead. We construct Bukhgeim's solutions $u = e^{i\tau(z-z_0)^2} f$ as before. They won't be solutions to the Schrödinger equation in the whole plane, but they are so in every open set where $\phi \equiv 1$. Calculating with those operators is easier since there is no boundary terms when integrating by parts. See also the next section.

6.3 Non-compactly supported potentials

An obvious question is whether we can do all the steps in the proof for potentials supported on the whole domain \mathbb{R}^2 . The first thing to do is to show the existence and norm estimates for the oscillating solutions. One way

to do that is to replace χ_Ω by a test function, integrate by parts as in the proof of theorem 4.1.1, and then let it tend to the constant one pointwise.

If ϕ is a test function, we are able to prove

$$\|\mathcal{E}(e^{i\tau R}\phi a)\|_{L^p(\mathbb{C})} \leq C(\|\phi\|_\infty + \|\bar{\partial}\phi\|_2)\tau^{-1/2}(\|a\|_p + \|\bar{\partial}a\|_{p^*}), \quad (6.7)$$

where $\frac{1}{p^*} = \frac{1}{2} + \frac{1}{p}$ and $2 < p < \infty$. This is proven by taking $\chi \in C_0^\infty(\mathbb{C})$ which is constant one near the origin and letting

$$h(z) = \frac{1 - \chi(\tau^{1/2}(z - z_0))}{\bar{z} - \bar{z}_0}\phi(z). \quad (6.8)$$

Then proceed as in the proof of 4.1.1 and note that

$$\|\phi - (\bar{z} - \bar{z}_0)h\|_2 + \tau^{-1}\|h\|_\infty + \tau^{-1}\|\bar{\partial}h\|_2 \leq \tau^{-1/2}. \quad (6.9)$$

Letting $\phi \rightarrow 1$ gives $\|\mathcal{E}(e^{i\tau R}a)\|_p \leq C\tau^{-1/2}(\|a\|_p + \|\bar{\partial}a\|_{p^*})$. This in turn implies

$$\begin{aligned} \|\mathcal{E}(e^{i\tau R}\bar{\mathcal{E}}(e^{-i\tau R}qf))\|_{L^p(\mathbb{C})} &\leq C\tau^{-1/2}(\|\bar{\mathcal{E}}(e^{-i\tau R}qf)\|_p + \|\bar{\Pi}(e^{-i\tau R}qf)\|_{p^*}) \\ &\leq C\tau^{-1/2}\|qf\|_{p^*} \leq C\tau^{-1/2}\|q\|_{L^2(\mathbb{C})}\|f\|_{L^p(\mathbb{C})}, \end{aligned} \quad (6.10)$$

since $\bar{\mathcal{E}} : L^{p^*} \rightarrow L^p$ and the Beurling operator $\bar{\partial}\bar{\mathcal{E}}$ is bounded on L^{p^*} . For the derivatives,

$$\begin{aligned} \|\nabla\mathcal{E}(e^{i\tau R}\bar{\mathcal{E}}(e^{-i\tau R}qf))\|_{L^p(\mathbb{C})} &\leq C\|\bar{\mathcal{E}}(e^{-i\tau R}qf)\|_{L^p(\mathbb{C})} \\ &\leq C\tau^{-1/2}(\|qf\|_p + \|f\bar{\partial}q\|_{p^*} + \|q\bar{\partial}f\|_{p^*}) \leq C\tau^{-1/2} \\ &\leq C\tau^{-1/2}(\|q\|_p + \|q\|_2 + \|\bar{\partial}q\|_2)(\|f\|_\infty + \|f\|_p + \|\bar{\partial}f\|_p) \\ &\leq C\tau^{-1/2}\|q\|_{W^{1,2}(\mathbb{C})}\|f\|_{W^{1,p}(\mathbb{C})} \end{aligned} \quad (6.11)$$

because $\nabla\mathcal{E} \cong (\text{Id}, \Pi) : L^p \rightarrow L^p$ and by Sobolev embedding. Thus we have shown the existence of Bukhgeim's oscillating solutions in the whole domain if the potential is in L^2 or $W^{1,2}$, with error term vanishing at a rate of $\tau^{-1/2}$ in L^p or $W^{1,p}$, respectively. This is not completely new, see for example [22, ch. 3, 4].

We must handle the error term of the stationary phase integral next. The space $W^{1,p}$ is a Banach algebra, so our error term is

$$\int \tau e^{i\tau R}(q_1 - q_2)r_{\tau, z_0} dm(z) \quad (6.12)$$

where $\sup_{z_0} \|r\|_{1,p} \leq C\tau^{-1/2}$. Take a suitable smooth function h cutting z_0 off. Split the integral by writing $1 = (1 - (\bar{z} - \bar{z}_0)h) + (\bar{z} - \bar{z}_0)h$. After integrating the second term by parts and using Hölder's inequality, we arrive at

$$\begin{aligned} \left| \int \tau e^{i\tau R} (q_1 - q_2) r dm \right| &\leq \dots \\ &\leq (\tau \|1 - (\bar{z} - \bar{z}_0)h\|_2 + \|\bar{\partial}h\|_2 + \|h\|_{p^*}) \|q_1 - q_2\|_{W^{1,2}} \|r\|_{W^{1,p}} \\ &\leq \tau^{\frac{2}{p+2}-\frac{1}{2}} \|q_1 - q_2\|_{W^{1,2}} \longrightarrow 0 \quad (6.13) \end{aligned}$$

as $\tau \rightarrow \infty$. Here p^* is the Hölder conjugate of p .

Assume that the potentials have a bit more smoothness and integrability, so that $\hat{q}_j \in L^1$. Write $Q = q_1 - q_2$. We get

$$\begin{aligned} \|q_1 - q_2\|_\infty &\leq \left\| Q - \int_\Omega \frac{2\tau}{\pi} e^{i\tau R} Q dm \right\|_\infty + \left\| \frac{2\tau}{\pi} \int_\Omega u^{(1)}(q_1 - q_2) u^{(2)} dm \right\|_\infty \\ &\quad + \left\| \frac{2\tau}{\pi} \int_\Omega e^{i\tau R} Q (1 - f_1 f_2) dm \right\|_\infty \\ &\leq \left\| (1 - e^{i\frac{\xi^2 + \bar{\xi}^2}{16\tau}}) \hat{Q} \right\|_1 + \left\| \frac{2\tau}{\pi} \int_\Omega u^{(1)}(q_1 - q_2) u^{(2)} dm \right\|_\infty + \tau^{\frac{2}{p+2}-\frac{1}{2}} \|Q\|_{W^{1,2}} \\ &\quad \longrightarrow \lim_{\tau \rightarrow \infty} \left\| \frac{2\tau}{\pi} \int_\Omega u^{(1)}(q_1 - q_2) u^{(2)} dm \right\|_\infty \quad (6.14) \end{aligned}$$

as $\tau \rightarrow \infty$ by dominated convergence and the previous deductions on the error term. Hence the inverse problem can be solved whenever $q_j \in W^{1,2}(\mathbb{C})$, $\hat{q}_j \in L^1(\mathbb{C})$ and the measurements imply the orthogonality relation

$$\int u^{(1)}(q_1 - q_2) u^{(2)} dm = 0. \quad (6.15)$$

Note that this does not follow from the equality of the corresponding scattering amplitudes. That would contradict the existence of certain counterexamples in [21].

6.4 No smoothness

Since we have stability for $q \in H^{s,(2,1)}(\Omega)$ with any $s > 0$, it is tempting to try to prove uniqueness for $q \in L^{(2,1)}(\Omega)$. We basically have to estimate three

terms, just like in the sketch of section 1.3:

$$\begin{aligned} & \left\| q_1 - q_2 - \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} (q_1 - q_2) dm \right\|, \\ & \left\| \frac{2\tau}{\pi} \int_{\Omega} u^{(1)}(q_1 - q_2) u^{(2)} dm \right\|, \\ & \text{and } \left\| \frac{2\tau}{\pi} \int_{\Omega} e^{i\tau R} (q_1 - q_2) (1 - f_1 f_2) dm \right\|. \end{aligned} \quad (6.16)$$

The first term is not a problem. We can use dominated convergence on the Fourier-side to see that that term vanishes as τ grows. The second term vanishes if we assume that $C_{q_1} = C_{q_2}$.

There's only the third term left. There are two choices. One is to try to show that $\|1 - f_1 f_2\| = o(\tau^{-1})$, which seems unlikely. Maybe some numerical simulations could shed a better light on this? The other option is to try to study how $f_1 f_2$ behaves with respect to z_0 , and see what kind of operator we have here. If f_j didn't depend on z_0 , then it would be a convolution operator, and we could show that the term vanishes using stationary phase. But there is dependence, so something nontrivial has to be done.

Actually, the problem seems to have been solved after the writing of this thesis. Imanuvilov and Yamamoto published the paper [27] in arXiv, and it claims to show uniqueness for $q_j \in L^p(\Omega)$, $p > 2$. They approximate the potentials by smooth functions and use results for oscillatory integral operators.

6.5 A reconstruction formula

There is a reconstruction formula by Bukhgeim in the last few lines of his article [3]. The idea is as follows. We have

$$q(z_0) \longleftarrow \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} q(z) dm(z) \quad (6.17)$$

by stationary phase. Let $u = e^{i\tau(z-z_0)^2} f$ be Bukhgeim's oscillating solution to $\Delta u + qu = 0$. Then $e^{i\tau(z-z_0)^2} = u + e^{i\tau(z-z_0)^2} (1 - f)$, so

$$q(z_0) \longleftarrow \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau(\bar{z}-\bar{z}_0)^2} qu(z) dm(z) + \int_{\Omega} \frac{2\tau}{\pi} e^{i\tau R} q(1 - f)(z) dm(z) \quad (6.18)$$

The second term tends to zero according to theorem 3.2.1. Use the fact that $qu = -\Delta u = -4\partial\bar{\partial}u$ and integrate by parts. Note that $\partial e^{i\tau(\bar{z}-\bar{z}_0)^2} = 0$. Hence

$$q(z_0) \longleftarrow \int_{\partial\Omega} \frac{4\tau}{\pi} \eta(z) e^{i\tau(\bar{z}-\bar{z}_0)^2} \bar{\partial}u(z) d\sigma(z) \quad (6.19)$$

After that, the idea is to reconstruct $\bar{\partial}u$ for Bukhgeim's solutions when we only know the boundary data. We haven't used all degrees of freedom when constructing u . This is because \mathcal{C} is only a right inverse of $\bar{\partial}$. We can always add an analytic function, and it is completely determined by its boundary values. Hence, we should look for ways to choose the analytic functions so that we may set some boundary values of u independently of q , and observe $\bar{\partial}u$.

According to [3], we may set $\operatorname{Re} u$ and $\operatorname{Re} \bar{\partial}u$, and observe $\operatorname{Im} u$ and $\operatorname{Im} \bar{\partial}u$. This is because the real part of a complex analytic function determines its imaginary part apart from a constant. In any case, this is just a theoretical tool for now. Any noise in the measurement of $\bar{\partial}u$ would get amplified and oscillated exponentially. For a recent result with an explicit boundary integral equation for $u_{\partial\Omega}$, see [40].

7 Calculations

Lemma 7.1. *If $c > 0$ then the Fourier transform of $t \mapsto e^{-ct^2/2}$ is the mapping $\xi \mapsto \frac{1}{\sqrt{c}}e^{-\xi^2/(2c)}$.*

Proof. This is a direct calculation using Cauchy's integral theorem. Let $c > 0$ and $\xi \in \mathbb{R}$. Then

$$\begin{aligned} (e^{-c\frac{t^2}{2}})^\wedge(\xi) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-c\frac{t^2}{2}-i\xi t} dt = \frac{1}{\sqrt{2\pi}} e^{-\frac{\xi^2}{2c}} \int_{-\infty}^{\infty} e^{-\left(\sqrt{\frac{c}{2}}t + \frac{i\xi}{2\sqrt{c/2}}\right)^2} dt \\ &= \frac{1}{\pi c} e^{-\frac{\xi^2}{2c}} \int_{-\infty}^{\infty} e^{-\left(s + \frac{i\xi}{2\sqrt{c/2}}\right)^2} ds = \frac{1}{\sqrt{\pi c}} e^{-\frac{\xi^2}{2c}} \int_{-\infty}^{\infty} e^{-u^2} du \quad (7.1) \\ &= \frac{1}{\sqrt{c}} e^{-\frac{\xi^2}{2c}} \end{aligned}$$

by Cauchy's integral theorem. This is justified because the function given by $z \mapsto e^{-z^2}$ is analytic and for any $A \in \mathbb{R}$ we have

$$\left| \int_s^{s+iA} e^{-z^2} dz \right| \leq \int_s^{s+iA} |e^{-z^2}| d\sigma(z) = \int_s^{s+iA} e^{A^2-s^2} d\sigma(z) = |A|e^{A^2-s^2}, \quad (7.2)$$

which tends to zero when $s \rightarrow \infty$ or $s \rightarrow -\infty$ along the real line. \square

Lemma 7.2. *Let $\tau > 0$ and define $\kappa_\tau : \mathbb{C} \rightarrow \mathbb{C}$ by $\kappa_\tau(z) = \frac{2\tau}{\pi} e^{i\tau(z^2 + \bar{z}^2)}$. It is a tempered distribution and*

$$\widehat{\kappa}_\tau(\xi) = \frac{1}{2\pi} e^{-i\frac{\xi^2 + \bar{\xi}^2}{16\tau}}. \quad (7.3)$$

Proof. The function κ_τ is bounded and measurable so it is a tempered distribution. Let $\varphi \in \mathcal{S}(\mathbb{R})$ be a Schwartz test function. Note that by lemma 7.1

$$\int_{\mathbb{R}} e^{-c\frac{t^2}{2}} \widehat{\varphi}(t) dt = \int_{\mathbb{R}} \frac{1}{\sqrt{c}} e^{-\frac{\xi^2}{2c}} \varphi(\xi) d\xi. \quad (7.4)$$

Let us choose a branch of the square root in the complex plane such that $\arg \sqrt{z} \in]-\frac{\pi}{2}, \frac{\pi}{2}]$. Now both sides of the previous equation are analytic functions of c in the right half-plane $\operatorname{Re}(c) > 0$. Hence they are equal also in the whole right half-plane.

Let $\phi, \psi \in \mathcal{S}(\mathbb{R})$ be two Schwartz test functions. By Fubini's theorem

and dominated convergence

$$\begin{aligned}
\int_{\mathbb{C}} e^{i\tau(z^2+\bar{z}^2)} (\phi(\xi_1)\psi(\xi_2))^\wedge(z) dm(z) &= \int_{\mathbb{R}^2} e^{i2\tau(x^2-y^2)} \widehat{\phi}(x)\widehat{\psi}(y) dm(x, y) \\
&= \int_{-\infty}^{\infty} e^{i2\tau x^2} \widehat{\phi}(x) dx \int_{-\infty}^{\infty} e^{-i2\tau y^2} \widehat{\psi}(y) dy \\
&= \lim_{\epsilon \rightarrow 0^+} \int_{-\infty}^{\infty} e^{-(2\epsilon-i4\tau)\frac{x^2}{2}} \widehat{\phi}(x) dx \int_{-\infty}^{\infty} e^{-(2\epsilon+i4\tau)\frac{y^2}{2}} \widehat{\psi}(y) dy \\
&= \lim_{\epsilon \rightarrow 0^+} \frac{1}{\sqrt{2\epsilon-i4\tau}} \int_{-\infty}^{\infty} e^{-\frac{\xi_1^2}{4\epsilon-i8\tau}} \phi(\xi_1) d\xi_1 \frac{1}{\sqrt{2\epsilon+i4\tau}} \int_{-\infty}^{\infty} e^{-\frac{\xi_2^2}{4\epsilon+i8\tau}} \psi(\xi_2) d\xi_2 \\
&= \frac{1}{\sqrt{4\tau}} e^{i\frac{\pi}{4}\tau} \frac{1}{\sqrt{4\tau}} e^{-i\frac{\pi}{4}\tau} \int_{-\infty}^{\infty} e^{\frac{\xi_1^2-\xi_2^2}{i8\tau}} \phi(\xi_1)\psi(\xi_2) dm(\xi_1, \xi_2) \\
&= \frac{1}{4\tau} \int_{\mathbb{C}} e^{-i\frac{\xi^2+\bar{\xi}^2}{16\tau}} \phi(\xi_1)\psi(\xi_2) dm(\xi), \quad (7.5)
\end{aligned}$$

so $\mathcal{F}\left\{\frac{2\tau}{\pi} e^{i\tau(z^2+\bar{z}^2)}\right\}(\xi) = \frac{1}{2\pi} e^{-i\frac{\xi^2+\bar{\xi}^2}{16\tau}}$. \square


Lemma 7.3. *Let $\Omega \subset \mathbb{C}$ be open, $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ non-increasing and $\epsilon \geq 0$. Then*

$$\int_{\Omega \setminus B(0, \epsilon)} f(|z|) dm(z) \leq \int_{\{\epsilon \leq |z| \leq \sqrt{\frac{m(\Omega)}{\pi} + \epsilon^2}\}} f(|z|) dm(z). \quad (7.6)$$

Proof. The idea is that the integral on the left-hand side is maximized when Ω is an annulus around the ball.

Let $r = \sqrt{m(\Omega)\pi^{-1} + \epsilon^2}$ denote the outer radius of this annulus. Write

$$\begin{aligned}
\Omega_r &= (\Omega \setminus B(0, \epsilon)) \setminus B(0, r), \\
\Omega_\cap &= (\Omega \setminus B(0, \epsilon)) \cap B(0, r), \\
A_r &= B(0, r) \setminus B(0, \epsilon) \setminus \Omega.
\end{aligned}$$



Note that

$$\begin{aligned}
m(\Omega_r) &= m(\Omega \setminus B(0, \epsilon)) - m(\Omega_\cap) \leq m(\Omega) - m(\Omega_\cap) \\
&= m(B(0, r) \setminus B(0, \epsilon)) - m(\Omega_\cap) \\
&= m(B(0, r) \setminus B(0, \epsilon) \setminus \Omega) = m(A_r), \quad (7.7)
\end{aligned}$$

and $\sup_{\Omega_r} f(|z|) \leq \inf_{A_r} f(|z|)$ because if $z \in \Omega_r$ then $|z| \geq r$, but it is smaller

than r in A_r . Hence

$$\begin{aligned} \int_{\Omega \setminus B(0,\epsilon)} f(|z|) dm(z) &= \int_{\Omega_\cap} f(|z|) dm(z) + \int_{\Omega_r} f(|z|) dm(z) \\ &\leq \int_{\Omega_\cap} f(|z|) dm(z) + \int_{A_r} f(|z|) dm(z) = \int_{B(0,r) \setminus B(0,\epsilon)} f(|z|) dm(z). \end{aligned} \quad (7.8)$$

□

Lemma 7.4. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain, $z_0 \in \mathbb{C}$ and $\tau > 0$. Then there exists $h \in W^{1,1}(\Omega)$ such that*

$$\tau \|1 - (\bar{z} - \bar{z}_0)h\|_{L^1(\Omega)} + \|h\|_{W^{1,1}(\Omega)} \leq 2\pi \left(\sqrt{\frac{m(\Omega)}{\pi}} + 2 + \ln_+ \sqrt{\frac{m(\Omega)\tau}{\pi}} \right). \quad (7.9)$$

Proof. Write $\epsilon = \tau^{-1/2}$. Define

$$H(z) = \begin{cases} \frac{1}{\bar{z}} & , |z| > \epsilon \\ \frac{|z|}{\epsilon \bar{z}} & , |z| < \epsilon \end{cases}. \quad (7.10)$$

Note that $H \in L^1_{loc} \subset \mathcal{D}'$. A straightforward integration by parts against a test function gives us

$$\partial H(z) = \begin{cases} 0 & , |z| > \epsilon \\ \frac{1}{2\epsilon|z|} & , |z| < \epsilon \end{cases}, \quad \bar{\partial} H(z) = \begin{cases} \frac{-1}{\bar{z}^2} & , |z| > \epsilon \\ \frac{-|z|}{2\epsilon \bar{z}^2} & , |z| < \epsilon \end{cases}. \quad (7.11)$$

Let $h(z) = H(z - z_0)$. Now $h \in W^{1,1}(\Omega)$ because Ω is bounded. The rest is straightforward calculations using lemma 7.3. We will show the hardest case, namely $\bar{\partial} h$. By the lemma, we have

$$\begin{aligned} \int_{\Omega} |\bar{\partial} H(z - z_0)| dm(z) &\leq \int_{\{|z| \leq \sqrt{\frac{m(\Omega)}{\pi}}\}} |\bar{\partial} H(z)| dm(z) \\ &= 2\pi \int_0^{\sqrt{\frac{m(\Omega)}{\pi}}} |\bar{\partial} H(r)| r dr = 2\pi \int_0^\epsilon \frac{dr}{2\epsilon} + 2\pi \int_\epsilon^{\sqrt{\frac{m(\Omega)}{\pi}}} \frac{dr}{r} \\ &= \pi + 2\pi \ln \sqrt{\frac{m(\Omega)}{\pi \epsilon^2}}, \end{aligned} \quad (7.12)$$

when $\epsilon < \sqrt{\frac{m(\Omega)}{\pi}}$. When not, then the upper bound is just π . This justifies \ln_+ in the estimate.

To estimate the first term, it is enough to note that

$$1 - \bar{z}H(z) = \begin{cases} 0 & , |z| > \epsilon \\ 1 - |z|\epsilon^{-1} & , |z| < \epsilon \end{cases} \quad (7.13)$$

and integrate over the ball $B(z_0, \epsilon)$. This gives us

$$\|1 - (\bar{z} - \bar{z}_0)h\|_{L^1(\Omega)} \leq \frac{\pi}{3}\epsilon^2. \quad (7.14)$$

Summing all the terms and estimating upwards a bit gives the estimate. \square

Remark 7.5. This works also for unbounded domains with finite measure.

Lemma 7.6. *Let $\Omega \subset \mathbb{C}$ be a bounded open set, $z_0 \in \mathbb{C}$ and $\tau > 0$. Write $\Omega_\tau = \Omega \setminus B(z_0, \tau)$. Then*

$$\begin{aligned} \|(\bar{z} - \bar{z}_0)^{-1}\|_{L^{(2,1)}(\Omega_\tau)} &\leq 2\sqrt{\pi} \ln \left(\frac{2m(\Omega)}{\pi\tau^2} + 1 + \sqrt{\left(\frac{2m(\Omega)}{\pi\tau^2} + 1\right)^2 - 1} \right), \\ \|(\bar{z} - \bar{z}_0)^{-2}\|_{L^{(2,1)}(\Omega_\tau)} &\leq 4\sqrt{\pi}\tau^{-1} \arctan \sqrt{\frac{m(\Omega)}{\pi\tau^2}}. \end{aligned} \quad (7.15)$$

Proof. We may assume that $z_0 = 0$ by the translation invariance of $L^{(2,1)}$ and $\Omega \mapsto m(\Omega)$. First, calculate

$$\begin{aligned} m(|z|^{-a}, \lambda) &= m\{z \in \Omega_\tau \mid |z|^{-a} > \lambda\} = m\{\Omega \setminus B(0, \tau) \cap B(0, \lambda^{-1/a})\} \\ &\leq \begin{cases} 0, & \lambda^{-1/a} \leq \tau \Leftrightarrow \tau^{-a} \leq \lambda \\ \pi(\lambda^{-2/a} - \tau^2), & \tau < \lambda^{-1/a} \leq \sqrt{\frac{m(\Omega)}{\pi}} + \tau^2 \Leftrightarrow \sqrt{\frac{m(\Omega)}{\pi}} + \tau^2 \leq \lambda < \tau^{-a} \\ m(\Omega), & \sqrt{\frac{m(\Omega)}{\pi}} + \tau^2 < \lambda^{-1/a} \Leftrightarrow \lambda < \sqrt{\frac{m(\Omega)}{\pi}} + \tau^2 \end{cases} \end{aligned} \quad (7.16)$$

then

$$(|z|^{-a})^*(s) = \inf\{\lambda \mid m(|z|^{-a}, \lambda) \leq s\} = \begin{cases} \tau^{-a}, & s = 0 \\ \sqrt{\frac{s}{\pi}} + \tau^{2-a}, & 0 < s < m(\Omega) \\ 0, & m(\Omega) \leq s \end{cases} \quad (7.17)$$

By Hölder's inequality $\|f\|_{(2,1)} \leq 2\|f\|_{2,1} = 2\int_0^\infty s^{-1/2}f^*(s)ds$, so

$$\| |z|^{-a} \|_{L^{(2,1)}(\Omega_\tau)} \leq 2 \int_0^\infty s^{-1/2} (|z|^{-a})^*(s) ds = 2 \int_0^{m(\Omega)} s^{-1/2} \left(\frac{s}{\pi} + \tau^2\right)^{-a/2} ds. \quad (7.18)$$

Case $a = 1$: We have $D \ln(x + \sqrt{x^2 - 1}) = (x^2 - 1)^{-1/2}$ for $x > 1$. Use the change of variables $u = \frac{2s}{\pi\tau^2} + 1$ to get

$$\begin{aligned} \int_0^{m(\Omega)} s^{-1/2} \left(\frac{s}{\pi} + \tau^2\right)^{-1/2} ds &= \sqrt{\pi} \frac{2}{\pi\tau^2} \int_0^{m(\Omega)} \frac{ds}{\sqrt{\left(\frac{2s}{\pi\tau^2} + 1\right)^2 - 1}} \\ &= \sqrt{\pi} \int_1^{\frac{2m(\Omega)}{\pi\tau^2} + 1} \frac{du}{\sqrt{u^2 - 1}} = \sqrt{\pi} \ln \left(\frac{2m(\Omega)}{\pi\tau^2} + 1 + \sqrt{\left(\frac{2m(\Omega)}{\pi\tau^2} + 1\right)^2 - 1} \right). \end{aligned} \quad (7.19)$$

Case $a = 2$: Use $D \arctan v = (v^2 + 1)^{-1}$ with the change of variables $v = \frac{u}{\sqrt{\pi\tau}}$ and $u = s^{1/2}$. Then

$$\begin{aligned} \int_0^{m(\Omega)} s^{-1/2} \left(\frac{s}{\pi} + \tau^2\right)^{-1} ds &= 2\pi \int_0^{\sqrt{m(\Omega)}} \frac{du}{u^2 + \pi\tau^2} = 2\pi \int_0^{\sqrt{\frac{m(\Omega)}{\pi\tau^2}}} \frac{\sqrt{\pi\tau} dv}{\pi\tau^2(v^2 + 1)} \\ &= 2\sqrt{\pi\tau}^{-1} \int_0^{\sqrt{\frac{m(\Omega)}{\pi\tau^2}}} \frac{dv}{v^2 + 1} = 2\sqrt{\pi\tau}^{-1} \arctan \sqrt{\frac{m(\Omega)}{\pi\tau^2}}. \end{aligned} \quad (7.20)$$

□

Lemma 7.7. *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain. Write*

$$\partial\Omega_\varepsilon = \{x \in \Omega \mid d(x, \partial\Omega) < \varepsilon\}. \quad (7.21)$$

Then there is $C_\Omega < \infty$ such that

$$m(\partial\Omega_\varepsilon) \leq C_\Omega \varepsilon \quad (7.22)$$

for any $\varepsilon \geq 0$.

Proof. Let $\{U_j\}$ be a finite open cover of $\partial\Omega$ such that each U_j is a cube and there exists Lipschitz functions f_j and orthonormal coordinate systems $(\zeta_{j,1}, \dots, \zeta_{j,n})$ in U_j such that

$$\begin{aligned} U_j &= \{\zeta_j \mid -s_j < \zeta_{j,k} < s_j \text{ for all } k\} \\ \Omega \cap U_j &= \{\zeta_j \mid \zeta_{j,n} < f(\zeta_{j,1}, \dots, \zeta_{j,n-1})\}. \end{aligned} \quad (7.23)$$

The existence of such an atlas follows by first taking an arbitrary open cover, the functions f_j and the coordinate axes, which are all given by the definition of a Lipschitz domain. Then cover each point in $\overline{\Omega} \cap U_j$ by a properly oriented cube $K \subset U_j$. The compactness of $\overline{\Omega}$ implies the rest.

From now on we will write $\tilde{x} = (x_1, \dots, x_{n-1})$ for $x \in \mathbb{R}^n$. Using the given coordinate systems we will see that

$$\begin{aligned} \partial\Omega_\varepsilon \cap U_j &= \{\zeta_j \mid \inf_{\xi_j} |(\check{\xi}_j, f_j(\check{\xi}_j)) - \zeta_j| < \varepsilon\} \\ &\subset \{\zeta_j \mid |(\check{\zeta}_j, f_j(\check{\zeta}_j)) - \zeta_j| < (M+1)\varepsilon\}, \end{aligned} \quad (7.24)$$

where $M = \max \|f_j\|_{C^{0,1}}$. This is true because of the following reasoning: if $\zeta_j \in \partial\Omega_\varepsilon \cap U_j$ then there is $\xi_j \in \partial\Omega \cap U_j$ such that $|\zeta_j - \xi_j| < \varepsilon$, and

$$\begin{aligned} |(\check{\zeta}_j, f_j(\check{\zeta}_j)) - \zeta_j| &\leq |\zeta_j - (\check{\zeta}_j, \xi_{j,n})| + |(\check{\zeta}_j, \xi_{j,n}) - (\check{\zeta}_j, f_j(\check{\zeta}_j))| \\ &= |\zeta_{j,n} - \xi_{j,n}| + |\xi_{j,n} - f_j(\check{\zeta}_j)| \leq |\zeta_j - \xi_j| + |f(\check{\xi}_j) - f_j(\check{\zeta}_j)| \\ &\leq (1+M)|\zeta_j - \xi_j| \leq (1+M)\varepsilon \end{aligned} \quad (7.25)$$

because $\xi_j \in \partial\Omega \cap U_j$ implies $\xi_{j,n} = f(\check{\xi}_j)$.

We are almost done. Let $\varepsilon_0 > 0$ be such that $\partial\Omega_\varepsilon \subset \cup U_j$ if $0 < \varepsilon < \varepsilon_0$. If $\varepsilon \geq \varepsilon_0$, then $m(\partial\Omega_\varepsilon) \leq m(\Omega) \leq \frac{m(\Omega)}{\varepsilon_0}\varepsilon$. The constant in front depends only on Ω since the choice of the atlas doesn't depend on ε . Hence we may assume that $\partial\Omega_\varepsilon \subset \cup U_j$.

Now

$$\begin{aligned} m(\partial\Omega_\varepsilon \cap U_j) &\leq m\{\zeta_j \mid |f_j(\check{\zeta}_j) - \zeta_{j,n}| < (M+1)\varepsilon\} \\ &\leq m\{\zeta_j \mid -s_j < \zeta_{j,k} < s_j \text{ for all } k \neq n, \text{ and } |f_j(\check{\zeta}_j) - \zeta_{j,n}| < (M+1)\varepsilon\} \\ &= (2s_j)^{n-1}(M+1)\varepsilon. \end{aligned} \quad (7.26)$$

The cover is finite, so summing all the pieces gives $m(\partial\Omega_\varepsilon) \leq C_\Omega \varepsilon$. \square

Lemma 7.8. *Let $\Omega \subset \mathbb{C}$ be bounded and Lipschitz. Then there is $C_\Omega < \infty$ such that for all $z_0 \in \mathbb{C}$ and $\varepsilon, \delta > 0$ there exists $h \in C_0^\infty(\Omega)$ with*

$$\begin{aligned} \|1 - (\bar{z} - \bar{z}_0)h\|_{L^{(2,1)}(\Omega)} &\leq C_\Omega \sqrt{\delta^2 + \varepsilon}, \\ \|h\|_{L^\infty(\Omega)} &\leq \delta^{-1}, \\ \|\bar{\partial}h\|_{L^{(2,1)}(\Omega)} &\leq C_\Omega \left(\varepsilon^{-1} \ln \left(1 + C_\Omega \varepsilon \delta^{-2} + \sqrt{(1 + C_\Omega \varepsilon \delta^{-2})^2 - 1} \right) + \delta^{-1} \right), \end{aligned} \quad (7.27)$$

Remark 7.9. ε describes how close the support of h is to $\partial\Omega$, while δ tells how close it is to z_0 .

Proof. Let $\phi \in C_0^\infty(\mathbb{C})$ be such that $\text{supp } \phi \subset B(0,1)$, $0 \leq \phi$, $\int \phi = 1$. Denote

$$\begin{aligned} \chi_\varepsilon &= \chi_{\{z \in \Omega \mid d(z, \partial\Omega) > 2\varepsilon\}} * \phi_\varepsilon \\ \chi^\delta &= \chi_{\mathbb{C} \setminus B(z_0, 2\delta)} * \phi_\delta, \end{aligned} \quad (7.28)$$

where $\phi_a(z) = a^{-2}\phi(z/a)$. It is clear that $\chi_\epsilon \in C_0^\infty(\Omega)$, $\chi^\delta \in C^\infty(\mathbb{C})$, $0 \leq \chi_\epsilon, \chi^\delta \leq 1$,

$$\begin{aligned} \chi_\epsilon(z) &= 0 \text{ if } d(z, \partial\Omega) < \epsilon, & \chi^\delta(z) &= 0 \text{ if } |z - z_0| < \delta \\ \chi_\epsilon(z) &= 1 \text{ if } d(z, \partial\Omega) > 3\epsilon, & \chi^\delta(z) &= 1 \text{ if } |z - z_0| > 3\delta \end{aligned} \quad (7.29)$$

and

$$\begin{aligned} |\partial_j \chi_\epsilon(z)| &= \left| \partial_j \int_{\{z \in \Omega | d(z, \partial\Omega) > 2\epsilon\}} \epsilon^{-2} \phi\left(\frac{z-z'}{\epsilon}\right) dm(z') \right| = \left| \epsilon^{-1} \int_{\{z \in \Omega | d(z, \partial\Omega) > 2\epsilon\}} \epsilon^{-2} (\partial_j \phi)\left(\frac{z-z'}{\epsilon}\right) dm(z') \right| \\ &\leq \epsilon^{-1} \int_{\mathbb{C}} |\partial_j \phi|(w) dm(w) = \epsilon^{-1} \|\partial_j \phi\|_1. \end{aligned} \quad (7.30)$$

Similarly, $|\partial_j \chi^\delta(z)| \leq \delta^{-1} \|\partial_j \phi\|_1$. Finally, set

$$h(z) = \frac{\chi_\epsilon(z) \chi^\delta(z)}{\bar{z} - \bar{z}_0}. \quad (7.31)$$

This is a compactly supported test function in Ω because $z_0 \notin \text{supp } \chi^\delta$ and $\chi_\epsilon \in C_0^\infty(\Omega)$. The estimate for $\|h\|_\infty$ comes directly from the support of χ^δ . For the first estimate, note that if $A \subset \mathbb{C}$ is measurable, then $\|\chi_A\|_{(p,q)} = \left(\frac{p}{q} + \frac{p}{q(p-1)}\right)^{1/q} (m(A))^{1/p}$ for $p > 1$, $q < \infty$. Hence we have

$$\begin{aligned} \|1 - (\bar{z} - \bar{z}_0)h\|_{L^{(2,1)}(\Omega)} &= \|1 - \chi_\epsilon \chi^\delta\|_{(2,1)} = \|\chi_{\{|z-z_0| < 3\delta \text{ or } d(z, \partial\Omega) < 3\epsilon\}}\|_{(2,1)} \\ &\leq 4\sqrt{m\{|z - z_0| < 3\delta \text{ or } d(z, \partial\Omega) < 3\epsilon\}} \leq 4\sqrt{9\pi\delta^2 + m(\partial\Omega_{3\epsilon})} \\ &\leq C_\Omega \sqrt{\delta^2 + \epsilon} \end{aligned} \quad (7.32)$$

by lemma 7.7.

Note that $\|fg\|_{(2,1)} \leq \|f\|_{(2,1)} \|g\|_\infty$. Use the estimates for $\partial_j \chi_\epsilon$ and $\partial_j \chi^\delta$, and keep track of the sets where $\chi_\epsilon, \chi^\delta$ are constants to get

$$\begin{aligned} \|\bar{\partial}h\|_{L^{(2,1)}(\Omega)} &\leq \left\| \frac{\chi^\delta}{\bar{z} - \bar{z}_0} \bar{\partial}\chi_\epsilon \right\|_{(2,1)} + \left\| \frac{\chi_\epsilon}{\bar{z} - \bar{z}_0} \bar{\partial}\chi^\delta \right\|_{(2,1)} + \left\| \frac{\chi_\epsilon \chi^\delta}{(\bar{z} - \bar{z}_0)^2} \right\|_{(2,1)} \\ &\leq \left\| \frac{1}{\bar{z} - \bar{z}_0} \right\|_{L^{(2,1)}(\partial\Omega_{3\epsilon} \setminus B(z_0, \delta))} \epsilon^{-1} \|\bar{\partial}\phi\|_1 + \left\| \frac{1}{\bar{z} - \bar{z}_0} \right\|_{L^{(2,1)}(B(z_0, 3\delta) \setminus B(z_0, \delta))} \delta^{-1} \|\bar{\partial}\phi\|_1 \\ &\quad + \left\| \frac{1}{(\bar{z} - \bar{z}_0)^2} \right\|_{L^{(2,1)}(\Omega \setminus B(z_0, \delta))} \end{aligned} \quad (7.33)$$

Again, by lemma 7.7, we have $m(\partial\Omega_{3\epsilon}) \leq C_\Omega \epsilon$, and $m(B(z_0, 3\epsilon) \setminus B(z_0, \epsilon)) = 8\pi\delta^2$. The rest follows directly from lemma 7.6. \square

Corollary 7.10. *Let $\Omega \subset \mathbb{C}$ be a bounded Lipschitz domain. Then, for every $\tau \geq 1$ and $z_0 \in \mathbb{C}$, there exists $h \in C_0^\infty(\Omega)$ such that*

$$\tau \|1 - (\bar{z} - \bar{z}_0)h(z)\|_{L^{(2,1)}(\Omega)} + \|h\|_{L^\infty(\Omega)} + \|\bar{\partial}h\|_{L^{(2,1)}(\Omega)} \leq C_\Omega \tau^{2/3}, \quad (7.34)$$

where C_Ω does not depend on τ or z_0 .

Proof. Take h as in lemma 7.8, and choose $\epsilon = \delta^2 = \tau^{-2/3}$. This gives us

$$C_\Omega(\tau^{2/3} + \tau^{1/3}) \leq 2C_\Omega \tau^{2/3} \quad (7.35)$$

for the right-hand side. □

References

- [1] Robert A. Adams and John J. F. Fournier. *Sobolev spaces*, volume 140 of *Pure and Applied Mathematics (Amsterdam)*. Elsevier/Academic Press, Amsterdam, second edition, 2003.
- [2] Giovanni Alessandrini. Stable determination of conductivity by boundary measurements. *Appl. Anal.*, 27(1-3):153–172, 1988.
- [3] Bukhgeim Alexander L. Recovering a potential from Cauchy data in the two-dimensional case. *J. Inverse Ill-Posed Probl.*, 16(1):19–33, 2008.
- [4] Kari Astala, Jennifer L. Mueller, Lassi Päivärinta, Allan Perämäki, and Samuli Siltanen. Direct electrical impedance tomography for nonsmooth conductivities. *Inverse Probl. Imaging*, 5(3):531–549, 2011.
- [5] Kari Astala and Lassi Päivärinta. Calderón’s inverse conductivity problem in the plane. *Ann. of Math. (2)*, 163(1):265–299, 2006.
- [6] Tomeu Barceló, Daniel Faraco, and Alberto Ruiz. Stability of Calderón inverse conductivity problem in the plane. *J. Math. Pures Appl. (9)*, 88(6):522–556, 2007.
- [7] Jöran Bergh and Jörgen Löfström. *Interpolation spaces. An introduction*. Springer-Verlag, Berlin, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [8] Eemeli Blåsten. *The Inverse Problem of the Schrödinger Equation in the Plane, A Dissection of Bukhgeim’s Result*. Licentiate thesis, University of Helsinki, Finland, 2010.
- [9] Eemeli Blåsten. Stability and uniqueness for the inverse problem of the Schrödinger equation in 2D with potentials in $W^{\epsilon,p}$. *ArXiv e-prints*, June 2011.
- [10] Alberto P. Calderón. Lebesgue spaces of differentiable functions and distributions. In Charles Bradfield Morrey, Jr., editor, *Partial Differential Equations*, volume IV of *Proc. Sympos. Pure Math.*, pages 33–49, Providence, R.I., 1961. American Mathematical Society.
- [11] Alberto P. Calderón. On an inverse boundary value problem. In *Seminar on Numerical Analysis and its Applications to Continuum Physics (Rio de Janeiro, 1980)*, pages 65–73. Soc. Brasil. Mat., Rio de Janeiro, 1980.

- [12] Sagun Chanillo. A problem in electrical prospection and an n -dimensional Borg-Levinson theorem. *Proc. Amer. Math. Soc.*, 108(3):761–767, 1990.
- [13] Albert Clop, Daniel Faraco, and Alberto Ruiz. Stability of Calderón’s inverse conductivity problem in the plane for discontinuous conductivities. *Inverse Probl. Imaging*, 4(1):49–91, 2010.
- [14] Michael Cwikel. The dual of Weak L^p . *Ann. Inst. Fourier (Grenoble)*, 25(2):xi, 81–126, 1975.
- [15] Joseph Diestel and John J. Uhl, Jr. *Vector measures*. American Mathematical Society, Providence, R.I., 1977. With a foreword by B. J. Pettis, Mathematical Surveys, No. 15.
- [16] Zhonghai Ding. A proof of the trace theorem of Sobolev spaces on Lipschitz domains. *Proc. Amer. Math. Soc.*, 124(2):591–600, 1996.
- [17] Nelson Dunford and Jacob T. Schwartz. *Linear operators. Part I*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1988. General theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, A Wiley-Interscience Publication.
- [18] Emilio Gagliardo. Caratterizzazioni delle tracce sulla frontiera relative ad alcune classi di funzioni in n variabili. *Rend. Sem. Mat. Univ. Padova*, 27:284–305, 1957.
- [19] Izraïl M. Gel’fand. Some aspects of functional analysis and algebra. In *Proceedings of the International Congress of Mathematicians, Amsterdam, 1954, Vol. 1*, pages 253–276. Erven P. Noordhoff N.V., Groningen, 1957.
- [20] Loukas Grafakos. *Classical and modern Fourier analysis*. Pearson Education, Inc., Upper Saddle River, NJ, 2004.
- [21] Piotr G. Grinevich and Roman G. Novikov. Transparent potentials at fixed energy in dimension two. Fixed-energy dispersion relations for the fast decaying potentials. *Comm. Math. Phys.*, 174(2):409–446, 1995.
- [22] Colin Guillarmou, Mikko Salo, and Leo Tzou. Inverse scattering at fixed energy on surfaces with euclidean ends. *Comm. Math. Phys.*, 303(3):761–784, 2011.
- [23] Colin Guillarmou and Leo Tzou. Calderón inverse problem with partial data on Riemann surfaces. *Duke Math. J.*, 158(1):83–120, 2011.

- [24] Jacques Hadamard. Sur les problèmes aux dérivés partielles et leur signification physique. *Princeton University Bulletin*, 13:49–52, 1902.
- [25] Robert H. Hanks. Interpolation by the real method between BMO, $L^\alpha(0 < \alpha < \infty)$ and $H^\alpha(0 < \alpha < \infty)$. *Indiana Univ. Math. J.*, 26(4):679–689, 1977.
- [26] Oleg Y. Imanuvilov, Gunther Uhlmann, and Masahiro Yamamoto. The Calderón problem with partial data in two dimensions. *J. Amer. Math. Soc.*, 23(3):655–691, 2010.
- [27] Oleg Y. Imanuvilov and Masahiro Yamamoto. Inverse boundary value problem for linear Schrödinger equation in two dimensions. *ArXiv e-prints*, August 2012.
- [28] Janne Kauhanen, Pekka Koskela, and Jan Malý. On functions with derivatives in a Lorentz space. *Manuscripta Math.*, 100(1):87–101, 1999.
- [29] Robert V. Kohn and Michael Vogelius. Determining conductivity by boundary measurements. *Comm. Pure Appl. Math.*, 37(3):289–298, 1984.
- [30] Robert V. Kohn and Michael Vogelius. Determining conductivity by boundary measurements. II. Interior results. *Comm. Pure Appl. Math.*, 38(5):643–667, 1985.
- [31] Lianfang Liu. *Stability Estimates for the Two Dimensional Inverse Conductivity Problem*. University of Rochester. Department of Mathematics, 1997.
- [32] Jan Malý. Absolutely continuous functions of several variables. *J. Math. Anal. Appl.*, 231(2):492–508, 1999.
- [33] Niculae Mandache. Exponential instability in an inverse problem for the Schrödinger equation. *Inverse Problems*, 17(5):1435–1444, 2001.
- [34] Katsushi Muramoto. Harmonic Bloch and BMO functions on the unit ball in several variables. *Tokyo J. Math.*, 11(2):381–386, 1988.
- [35] Adrian I. Nachman. Reconstructions from boundary measurements. *Ann. of Math. (2)*, 128(3):531–576, 1988.
- [36] Adrian I. Nachman. Global uniqueness for a two-dimensional inverse boundary value problem. *Ann. of Math. (2)*, 143(1):71–96, 1996.

- [37] Jindřich Nečas. *Direct Methods in the Theory of Elliptic Equations*. Springer Monographs in Mathematics. Springer Berlin Heidelberg, 2012. Translation from the 1967 French edition.
- [38] Júlio S. Neves. Spaces of Bessel-potential type and embeddings: the super-limiting case. *Math. Nachr.*, 265(1):68–86, 2004.
- [39] Roman G. Novikov and Matteo Santacesaria. A global stability estimate for the Gel’fand-Calderón inverse problem in two dimensions. *J. Inverse Ill-Posed Probl.*, 18(7):765–785, 2010.
- [40] Roman G. Novikov and Matteo Santacesaria. Global uniqueness and reconstruction for the multi-channel Gel’fand-Calderón inverse problem in two dimensions. *Bull. Sci. Math.*, 135(5):421–434, 2011.
- [41] Richard O’Neil. Convolution operators and $L(p, q)$ spaces. *Duke Math. J.*, 30(1):129–142, 1963.
- [42] Miroslav Pavlović. A remark on BMO and Bloch functions on the unit ball of \mathbb{R}^n . *Tokyo J. Math.*, 16(2):291–294, 1993.
- [43] Néstor M. Rivière. Interpolation à la Marcinkiewicz. *Rev. Un. Mat. Argentina*, 25(3–4):363–377, 1970/71. Collection of articles dedicated to Alberto González Domínguez on his sixty-fifth birthday.
- [44] Walter Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- [45] Friedrich Sauvigny. *Partial differential equations. 2*. Universitext. Springer-Verlag, Berlin, 2006. Functional analytic methods, With consideration of lectures by E. Heinz, Translated and expanded from the 2005 German original.
- [46] John Sylvester and Gunther Uhlmann. A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math. (2)*, 125(1):153–169, 1987.
- [47] Ilia N. Vekua. *Generalized analytic functions*. Pergamon Press, London, 1962. Translation from the 1959 Russian edition.

Index of Notation

- $(A_0, A_1)_{[\theta]}$; interpolation space, 10
- $BC(X, A)$; bounded continuous functions, 16
- $\mathcal{C}, \overline{\mathcal{C}}$; Cauchy operators, 21
- C_q ; boundary data, 38
- $d(C_{q_1}, C_{q_2})$ distance between boundary data, 39
- $\partial\Omega_\varepsilon$; ε -neighborhood of $\partial\Omega$, 55
- η ; complex normal vector, 22
- $\mathcal{F}(A)$, 10
- $f^*(s)$; non-increasing rearrangement, 6
- $f^{**}(t)$, 7
- $H^{s,(p,q)}$; Lorentz-Bessel potential space, 18
- Λ_q ; Dirichlet-Neumann map, 37
- $L^{p,q}, L^{(p,q)}$; Lorentz spaces, 6
- $L^{(p,q)}(\mathbb{R}^n, l_2^s)$, 19
- L^{p*} ; weak L^p -spaces, 7
- $m(f, \lambda)$; distribution function, 6
- $M^s(\Omega)$, 32
- $\|\Lambda_{q_1} - \Lambda_{q_2}\|$, 38
- ν ; normal vector, 22
- $\Pi, \overline{\Pi}, \partial\mathcal{C}, \overline{\partial\mathcal{C}}$; Beurling operator, 22
- Q ; $q_1 - q_2$, 41
- R ; the phase function $(z - z_0)^2 + (\bar{z} - \bar{z}_0)^2$, 25, 28
- τ_0 ; minimum value of τ , 42
- Tr; trace mapping, 36
- $W/W_0, W^{1,2}/W_0^{1,2}$; quotient space, 36

Index of Subjects

- Alessandrini's identity, 2, **39**, 40
- Banach couple, **10**, 11, 12
- Banach's fixed point theorem, 34
- Beurling operator, **22**, 47
- boundary data, 1, 4, **38**, 39
- boundary integral equation, 50
- boundary term, 22
- Carleman estimate, 5, **28**, 30, 32, 34
- Cauchy data, *see* boundary data
- Cauchy operator, **21**
- compatible, **10**
- complex Gaussian, 51
- counterexamples, 40, 48
- cut-off function, 53, 56
- Dirichlet-Neumann map, 4, 36, **37**,
see also boundary data
- distribution function, **6**
- ε -neighborhood, 55
- equation
- conductivity, 2
 - Schrödinger, 4
- error term estimate, **26**, 44
- extension operator, 34, 45
- functions
- countably valued, 8, 15
 - simple, **6**, 13
 - strongly measurable, **6**
- Green's formula, *see* integration by parts
- Hadamard's criteria, 36
- ill-posed, *see* counterexamples
- integration by parts, 4, **22**, 28, 39
- interpolation
- A -valued Lorentz spaces, 12
 - complex, **10**
 - Lorentz-Bessel spaces $H^{s,(p,q)}$, 20
 - multilinear operators, 11
- inverse problem
- Calderón, 2
 - Gel'fand-Calderón, 1
 - solution, 41
- L^p potential, *see* non-smooth potential
- mean value inequality, **24**
- Mihlin multiplier theorem, **19**, 20
- Minkowski's integral inequality, **8**
- non-increasing rearrangement, **6**
- non-smooth potential, 3, **49**
- orthogonality relation, *see* Alessandrini's identity
- Phragmén-Lindelöf principle, 11
- Poisson kernel, 14
- reconstruction, 3, **49**
- retract, **19**
- solutions
- Bukhgeim's, 4, **34**, 42
 - whole domain, 47
 - complex geometric optics, **2**
- spaces
- bounded continuous functions, **16**
 - complex interpolation, **10**
 - Lorentz, **6**
 - Lorentz-Bessel potential, **18**
 - Lorentz-Sobolev, **16**
 - weak- L^p , **7**
- stability, *see* inverse problem solution
- stationary phase, 4, **25**, 41, 49
- three lines theorem, 12
- trace mapping, 36
- well-posed problem, 37