

1 The spoilage flora of vacuum-packaged, sodium nitrite or potassium nitrate treated,
2 cold-smoked rainbow trout stored at 4°C or 8°C

3

4 Ulrike Lyhs *, Johanna Björkroth, Eija Hyytiä and Hannu Korkeala

5

6 Department of Food and Environmental Hygiene, Faculty of Veterinary Medicine,

7 University of Helsinki,

8 Helsinki, Finland

9

10

11

12

13 *Corresponding author: Ulrike Lyhs

14 Mailing address: Department of Food and Environmental Hygiene

15 Faculty of Veterinary Medicine

16 P.O. Box 57

17 FIN-00014 University of Helsinki

18 Finland

19 Tel: +358 9 70849 706. Fax: + 358 9 70849 718

20 E-mail address: Ulrike.Lyhs@Helsinki.Fi

21

22

23

24

25

26

27

28

29

30

31

32

33

34

1 **Abstract**

2

3 The spoilage flora of vacuum-packaged, salted, cold-smoked rainbow trout fillets,
4 with or without the addition of nitrate or nitrite, stored at 4°C and 8°C, was studied.

5 Of 620 isolates, lactic acid bacteria were the major fraction (76%), predominating in

6 all samples of spoiled product. However, the phenotypical tests used were

7 insufficient to identify the lactic acid bacteria to the species level. Gram-positive,

8 catalase-positive cocci, Gram-negative, oxidase-negative rods and Gram-negative,

9 oxidase-positive rods were found in 6%, 16% and 2% of the samples, respectively. Of

10 39 Gram-positive, catalase-positive cocci, 29 were identified as staphylococci and 10

11 as micrococci. Eighty-five isolates were found to belong to the family

12 *Enterobacteriaceae*, with 45 of those being *Serratia plymuthica*. Eleven isolates from

13 the nitrate treated samples stored at 8°C were identified as *Pseudomonas aeruginosa*.

14 The occurrence of *P. aeruginosa* and staphylococci in the nitrate-containing samples,

15 stored at 8°C, may cause problems with respect to the safety of the product. The types

16 of lactic acid and other bacteria in the spoilage flora were generally reduced by the

17 addition of nitrate or nitrite to fillets.

18

19

20 **Keywords:** Fish; Spoilage; Vacuum-packaging; Cold-smoking; Nitrite; Nitrate;

21 NaCl;

22

23

24

25

26

1 **1. Introduction**

2
3 Vacuum-packaged, cold-smoked fish products are highly perishable foods.
4 Previous studies with cold-smoked salmon have shown that increasing the salt
5 concentration and decreasing the storage temperature extend their storage life
6 (Hildebrandt and Erol, 1988; Civera et al., 1995; Truelstrup Hansen et al., 1995). During
7 their storage at chill temperatures, a complex microflora develops which is dominating by
8 lactic acid bacteria at the end of the storage period along with lower numbers of other
9 bacteria like *Enterobacteriaceae*, *Pseudomonas* spp., enterococci, micrococci and
10 yeasts (Magnússon and Traustadóttir, 1982; Schneider and Hildebrandt, 1984; Shimasaki
11 et al., 1994; Civera et al., 1995). There was variation in the composition of the
12 microflora described by the authors, probably due to the different processes applied
13 and differing smokehouse production environment (Truelstrup Hansen, 1995).
14 However, only limited work has been carried out about the exact composition and
15 characterisation of the lactic acid bacteria and other bacteria as part of the spoilage
16 flora of vacuum-packaged, cold-smoked fish products.

17 Nitrate (NO_3) has been added to the curing salt mixture of certain semi-
18 preserved pickled fish products in order to delay spoilage and to control microbial
19 activity during storage (Pedersen and Meyland, 1981; Knøchel and Huss, 1984). It
20 may also act as a reservoir of nitrite if nitrate-reducing bacteria are present
21 (Skovgaard, 1992). Nitrite is an important antimicrobial agent. It has shown to have
22 an inhibitory effect on bacterial spoilage and *Clostridium botulinum* growth and toxin
23 production also in fish (Sofos et al., 1979; Pierson and Smoot, 1987; Hyytiä et al.,
24 1997). Combinations of sodium chloride (NaCl) and sodium nitrite (NaNO_2) or
25 potassium nitrate (KNO_3) have been used as a preservative in hot-smoked fish
26 products (Pelroy et al., 1982) and cold-smoked rainbow trout (Hyytiä et al., 1997).

1 However, no reports been published about the changes in the bacterial groups causing
2 spoilage after adding of nitrate or nitrite to cold-smoked fish product.

3 As the use of nitrate or nitrite in cold-smoked fish might be
4 advantageous, this study was undertaken to characterise the spoilage flora of vacuum-
5 packaged, cold-smoked rainbow trout stored at 4°C or 8°C and to determine the
6 effect of nitrate and nitrite on the spoilage flora of the product.

7

8 **2. Materials and methods**

9

10 2.1. Samples

11

12 Rainbow trout (*Oncorhynchus mykiss*) from two Finnish fish farms was
13 used. Before brining, the trout were deheaded and filleted at a processing plant. The
14 fillets had an average weight of 600-900 g. Brining was carried out by the injection
15 method. The pressure used in the brine injection equipment (Fomaco 44/176, Fomaco
16 Food Machinery Company A/S, Køge, Denmark) was 1.6 bar. The brine
17 concentration was 21%, producing a NaCl concentration of 2.2% (w/w) in the final
18 product. The NaNO₂ and KNO₃ (Riedel-deHaën AG, Seelze, Germany)
19 concentrations of the curing solutions were 3 g/l and 13 g/l respectively, producing
20 nitrite and nitrate concentrations of 166 ppm and 686 ppm in the product after
21 preparation (Hyytiä et al., 1997). The fillets were cold-smoked overnight at 18-21°C,
22 at the processing plant, in an electronically controlled, electrically heated smoke house
23 equipped with an external smoke generator (Alpas, Alpas GmbH, Bremen, Germany).
24 After the smoking process, the fillets were vacuum-packaged, using a Multivac R
25 7000 1976 packaging machine (Multivac Verpackungsmaschinen, Wolfertschwenden,

1 Germany), in a polyethylene / polyamide film (Suomen Union Verpackungs Ltd,
2 Helsinki, Finland) with an oxygen permeability of 29-45 ml O₂/m²/ 24 h /atm (23^oC,
3 50% RH) and a water vapour permeability of 10-15 g/m²/ 24 h (38^oC, 90% RH).
4 Immediately after processing the samples were transported to the laboratory and
5 stored at either 4^oC or 8^oC.

6 2.2. Sensory evaluation

7

8 Sensory evaluation was performed once a week in order to determine
9 when the samples studied were spoiled. The sensory evaluation panel consisted of 9
10 or 10 trained panelists. The samples were evaluated for aroma and taste using the
11 method described by Amerine et al. (1965) on a scale from zero to five, in which a
12 score of two points or less indicated unacceptable product. The sample was deemed
13 spoiled if at least two judges considered it unfit. The colonies were selected from
14 samples which were deemed spoiled.

15

16 2.3. Microbiological analyses

17

18 Each 10 g cold-smoked, rainbow trout sample was homogenized with 90
19 ml of 0.1% (w/v) peptone water and 10-fold serial dilutions were used for
20 microbiological analyses. The aerobic plate count (APC) was determined by the
21 method of the Nordic Committee on Food Analysis (1986) using Plate Count Agar
22 (Difco, Detroit, Michigan, USA). At least 10 colonies from the highest dilutions that
23 yielded colonies were selected at random from the APC plates when the total bacteria
24 count was > 10⁷ cfu/g.

1 For samples stored at 4^oC, totals of 99, 110 or 100 isolates were
2 obtained from the flora from samples containing NaCl only, NaCl and KNO₃ or NaCl
3 and NaNO₂, respectively. For samples stored at 8^oC, totals of 104, 108 or 99 isolates
4 were obtained from the flora from samples containing NaCl only, NaCl and KNO₃ or
5 NaCl and NaNO₂, respectively.

6 All isolates were Gram-stained and were tested for haemolytic activity
7 (Columbia agar base, GIBCO BRL, Paisley, UK). Also, all organisms were grown on
8 brain heart infusion (BHI) agar (Difco) at 25^oC to test for catalase production (Baird-
9 Parker, 1979).

10 11 2.4. Characterisation tests

12
13 Gram-positive, catalase-negative bacteria were tested for growth in de
14 Man, Rogosa and Sharpe (MRS) broth (Difco) incubated at 25^oC for 3 days. Isolates
15 showing growth were further plated on Rogosa selective *Lactobacillus* (SL) agar
16 (Orion Diagnostica, Espoo, Finland). The plates were incubated anaerobically at
17 25^oC for 5 days using a model BR 38 gas-generating kit (Oxoid, Basingstoke, UK) in
18 an anaerobic jar. Growth on Slanetz and Bartley agar (Orion Diagnostica) was
19 observed after 2 days' incubation at 37^oC. Production of gas from glucose was
20 studied by the method of Schillinger and Lücke (1987). Acetoin production was
21 detected using the Voges-Proskauer test after 3 days incubation. Hydrolysis of
22 arginine was examined as described by Reuter (1970). Lactic acid configuration was
23 determined enzymatically using a UV method kit (Boehringer Mannheim GmbH,
24 Mannheim, Germany), according to the manufacturer`s instructions. The presence of

1 m-DPA in the cell walls was determined by the two - dimensional - thin layer paper
2 chromatography method of Harper and Davis (1979).

3 The Gram-positive, catalase-positive cocci were examined for acid
4 production from glycerol in the presence of erythromycin (Schleifer and Kloos, 1975)
5 and sensitivity to lysostaphin by the method of Kloos et al. (1974). For further
6 identification, API Staph (bio Mérieux, Marcy l'Etoile, France) was used and the
7 results were recorded after incubation at 25⁰C for 24 - 48 h.

8 Gram-negative microorganisms were examined for oxidase production
9 using Kovàcs reagent (Kovàcs, 1956). For further identification, API 20 E (bio
10 Mérieux) and API 20 NE (bio Mérieux) were used and the results were recorded after
11 incubation at 25⁰C for 24 - 48 h.

12 Yeasts were isolated on Sabouraud-Dextrose-Medium (Oxoid) and
13 Rose-Bengal-Medium (Difco) after 3 days' incubation at 25⁰C.

14
15
16 **3. Results**

17
18 Of 620 isolates, 469 Gram-positive, catalase-negative cocci or rods,
19 which grew on MRS were classified as lactic acid bacteria; 39 were Gram-positive,
20 catalase-positive cocci; 98 were Gram-negative, oxidase-negative rods; and 12
21 isolates were Gram-negative, oxidase-positive rods (Table 1).

22 Lactic acid bacteria predominated in the spoilage flora of all samples.
23 The isolated lactic acid bacteria were divisible into seven subgroups on the basis of
24 cell morphology, the formation of gas from glucose, the growth on Rogosa selective
25 *Lactobacillus* (SL) agar, the lactic acid isomers produced, the hydrolysis of arginine,

1 the production of acetoin and the presence of diaminopimelic acid in the cell walls
2 (Table 2). The 189 isolates of subgroup 1 were homofermentative rods which grew
3 well on SL agar. The second subgroup of 50 isolates were heterofermentative oval
4 cocci which did not grow on SL agar. The three isolates forming subgroup 3 were
5 heterofermentative rods with diaminopimelic acid in their cell walls. The 182 isolates
6 in subgroup 4 were heterofermentative oval cocci that grew on SL agar. The 3 isolates
7 in subgroup 5 were cocci which formed colonies with a red-pink centre on Slanetz
8 and Bartley agar but did not grow on SL agar. They produced predominantly L(+) -
9 lactic acid. They were isolated only from the samples without KNO₃ or NaNO₂ which
10 were stored at 8^oC. Subgroup 6 contained heterofermentative and subgroup 7
11 homofermentative cocci or oval cocci. Table 3 presents the distributions of the lactic
12 acid bacteria subgroups in the differently cured samples at either storage temperature.

13 Based on the production of acid from glycerol in the presence of
14 erythromycin and lysostaphin sensitivity, 29 of the 39 Gram-positive, catalase-
15 positive cocci were identified as staphylococci and 10 as micrococci. From the API
16 Staph tests, most of the staphylococcal isolates from the Gram-positive, catalase-
17 positive cocci were classifiable as *Staphylococcus epidermidis* or *Staphylococcus*
18 *hominis*. Other isolates were *Staphylococcus sciuri*, *Staphylococcus capitis*,
19 *Staphylococcus warneri*, *Staphylococcus intermedius* or *Staphylococcus lentus*. Three
20 of the ten micrococcal isolates were identified as *Micrococcus kristinae*. Micrococci
21 were mainly isolated from the nitrate-containing samples stored at 4^oC. Nine
22 staphylococci originated from the nitrate-containing samples stored at 8^oC.

23 The API 20 E tests placed 85 of the Gram-negative, oxidase-negative
24 rods in the family *Enterobacteriaceae*, 9 in the genus *Xanthomonas* and one in the
25 genus *Acinetobacter*. Most of the *Enterobacteriaceae* were identified as *Serratia*

1 *plymuthica*, *Serratia liquefaciens*, *Hafnia alvei* or *Enterobacter* spp. (Table 4). Of the
2 45 *S. plymuthica* isolates, 32 originated from the samples containing NaCl only and
3 stored at 4^oC. Of the nine *H. alvei*, six were recovered from the nitrate-containing
4 samples and two from the samples which contained NaCl only and were stored at
5 8^oC. All eight *Enterobacter amnigenus* isolates were isolated from nitrate-containing
6 samples stored at 8^oC and all five *Enterobacter sakazakii* isolates from the samples
7 which contained NaCl only and which were stored at 8^oC.

8 From the API 20 NE tests 11 of the Gram-negative, oxidase-positive
9 rods were identified as *Pseudomonas aeruginosa*. They originated from the nitrate-
10 containing samples stored at 8^oC. The remaining one isolate was identified as
11 *Ochrobacterium* and it originated from a nitrite-containing sample which was stored
12 at 4^oC. The isolates originating from the samples which contained NaCl only did not
13 include Gram-negative, oxidase-positive rods.

14 Two colonies obtained from nitrite-containing samples which were
15 stored at 8^oC grew both on Sabouraud-Dextrose-Medium and Rose-Bengal-Medium.

16

17 **4. Discussion**

18

19 The spoilage flora in all samples mainly consisted of lactic acid bacteria.
20 The dominance of lactic acid bacteria in vacuum-packaged, lightly preserved fish
21 products after a few weeks' storage at chilled temperatures has been reported
22 previously. Magnússon and Traustadóttir (1982) found lactic acid bacteria dominating
23 in vacuum-packaged cold-smoked herring, as did Schneider and Hildebrandt (1984),
24 Hildebrandt and Erol (1988), Shimasaki et al. (1994), Truelstrup Hansen (1995) and

1 Civera et al. (1995) in vacuum-packaged cold-smoked salmon and Jeppesen and Huss
2 (1993) in vacuum-packaged sugar-salted ('gravad') fish.

3 The relative proportion of lactic acid bacteria in microflora was higher
4 in the nitrite- and nitrate-containing samples than in the samples which contained
5 NaCl only at both storage temperatures. Lactic acid bacteria have previously been
6 reported to be resistant to nitrite (Doods and Collins-Thompson, 1984; Skovgaard,
7 1992). Their nitrite resistance may explain the high proportion of lactic acid bacteria
8 found in the nitrite-containing samples in this study.

9 Based on the characteristics of subgroup 1 of lactic acid bacteria, these
10 bacteria could be considered as homofermentative or facultatively heterofermentative
11 lactobacilli. The occurrence of lactobacilli with homofermentative glucose
12 metabolism has also been reported by Magnússon and Traustadóttir (1982) in
13 vacuum-packaged cold-smoked herring fillets stored for 12 weeks at chill
14 temperatures. In the present study, their proportion in the nitrate-containing samples
15 was higher than in the nitrite-containing samples and in the samples which contained
16 NaCl only (Table 3).

17 The isolates in subgroup 2 could be considered to belong to
18 *Leuconostoc/Weissella*-species. The dominance of leuconostocs has been reported
19 previously. Jeppesen and Huss (1993) studied the lactic acid bacteria from vacuum-
20 packaged, minced herring and identified all isolated lactic acid bacteria as
21 *Leuconostoc* spp. However, Mauguin and Novel (1994) found that only eight out of
22 86 lactic acid bacteria isolated from various samples of seafood belonged to the genus
23 *Leuconostoc*. In the present study, high numbers of heterofermentative lactobacilli
24 and *Leuconostoc* spp. occurred in the nitrite treated samples stored at 8⁰C (Table 3).

1 The bacteria in subgroup 3 possessing m-DPA in their cell walls
2 appeared to belong to the genus *Carnobacterium*. Carnobacteria have earlier been
3 found in vacuum-packaged ‘gravad’ fish (Leisner et al., 1994) and some other
4 vacuum-packaged fish products (Mauguin and Novel, 1994). However, Gancel et al.
5 (1997) did not find any carnobacteria in fillets of vacuum-packaged smoked and
6 salted herring and proposed smoking the fish to be the reason.

7 The other lactic acid bacterium groups formed could not be identified to
8 the species level by the phenotypical methods used. Subgroup 4, forming the second
9 largest group, could be classified as leuconostocs because of their cell morphology,
10 oval cocci, and their formation of gas from glucose. On the other hand, they grew on
11 SL agar as do heterofermentative lactobacilli. Most of them were found in the nitrite-
12 containing samples stored at either 4°C or 8°C.

13 The three isolates in subgroup 5 seemed to belong to the genus
14 *Enterococcus*. The occurrence of enterococci in vacuum-packaged cold-smoked
15 salmon has also been reported previously (Schneider and Hildebrandt, 1984;
16 Hildebrandt and Erol, 1988). Ben Embarek et al. (1994) isolated enterococci during
17 studies of bacterial survivors in *sous-vide* cooked fish fillets. The fractions of
18 lactic acid bacteria in subgroups 6 and 7 decreased after the addition of nitrite and
19 nitrate, indicating possible sensitivity to this kind of treatment. The phenotypical tests
20 were insufficient to characterise accurately the dominant lactic acid genera of these
21 bacterial groups. Species level identification of these above named bacterial groups
22 warrants further analysis such as genotyping.

23 The species identification of *Enterobacteriaceae* in this study generally
24 agrees with the results of Truelstrup Hansen (1995), who studied spoiled vacuum-
25 packaged cold-smoked salmon. However, there are no previous reports about the high

1 prevalence of *S. plymuthica* in fish products. This can be due to the fact that the fish
2 of the present study were originated in farms located in brackish water. *S. plymuthica*
3 strains isolated from water have been isolated frequently from fresh water (Nieto, et
4 al., 1990).

5 Micrococci were mainly isolated from the nitrate treated samples stored
6 at 4^oC. It is possible that nitrate might facilitate the growth of these bacteria.
7 Micrococci, as strict aerobic organisms, are presumed to use nitrate as an alternative
8 electron acceptor to oxygen under vacuum (Taylor and Shaw, 1975). The highest
9 prevalence of staphylococci was detected in the nitrate treated samples stored at 8^oC.
10 The anaerobic respiration of nitrate appears to be widespread among facultatively
11 anaerobic bacteria, such as staphylococci (Doelle, 1975). This may explain why
12 staphylococci were found in the nitrate treated samples. Of the nitrate treated samples
13 stored at 8^oC 11 isolates, were classified as *P. aeruginosa*. Since it is well known, that
14 *P. aeruginosa* can utilize nitrate as an electron acceptor it is able to grow under
15 anaerobic conditions if nitrate is present (Yamanaka et al., 1959). No *Pseudomonas*
16 spp. were isolated from any of the samples containing NaCl only, indicating the
17 attribution of vacuum packaging and sensitivity of *Pseudomonas* spp. to low oxygen
18 and high carbon dioxide levels (Clark and Takacs, 1980; Flick et al., 1991).
19 Therefore, the growth of *P. aeruginosa* may cause food hygienic problems in
20 vacuum-packed fish products treated with nitrate in the case of temperature abuse.

21 The composition of the spoilage flora was found to be affected by the
22 nitrate and nitrite treatment. Insensitivity to nitrate and nitrite, favoured by the
23 anaerobic conditions, resulted in lactic acid bacteria to constitute the major proportion
24 of the total flora in the nitrate- and nitrite-containing samples. However, the types of
25 lactic acid and other bacteria in the spoilage flora were generally reduced by the

1 addition of nitrate or nitrite to the product. The occurrence of *P. aeruginosa* and
2 staphylococci in the nitrate-containing samples stored at 8°C may cause problems
3 with respect to the safety of the product. Therefore, nitrate is not recommended as a
4 preservative in this type of fish product and the maintenance of the chill chain under
5 the temperature of 4°C should be ensured.

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Acknowledgements

The authors wish to thank Dr. Satu Sankari for her kind help and Mirja Eerikäinen for her excellent technical assistance. This research was supported by the Centre for International Mobility (CIMO) in Helsinki, the Finnish Veterinary Association and the Faculty of Veterinary Medicine of the University of Helsinki.

1

2

3 **References**

4

5 Amerine, M.A., Pangborn, R.M. and Roessler, E.B., 1965. Principles of Sensory
6 Evaluation of Food. Academic Press, New York and London, pp. 354-366.

7

8 Baird-Parker, A.C., 1979. Methods for identifying staphylococci and micrococci. In:
9 F.A. Skinner and D.W. Lovelock (Eds.), Identification Methods for Microbiologists.
10 Academic Press, London, pp. 201-210.

11

12 Ben Embarek, P.K., Jeppesen, V. and Huss, H.H., 1994. Antibacterial potential of
13 *Enterococcus faecium* strains from *sous-vide* cooked fish fillets. Food Microbiol. 11,
14 525-536.

15

16 Civera, T., Parisi, E., Amerio, G.P. and Giaccone, V., 1995. Shelf-life of vacuum-
17 packed smoked salmon: microbiological and chemical changes during storage. Arch.
18 Lebensmittelhyg. 46, 1-24.

19

20 Clark, D.S. and Takacs, J., 1980. Gases as preservatives. In: International
21 Commission on Microbiological Specifications for Food, Microbial Ecology of
22 Foods. vol. 1, Factors Affecting Life and Death of Microorganisms. Academic Press,
23 New York, p. 171.

24

1 Doelle H.W., 1975. Bacterial metabolism. 2nd ed. Academic Press, London, pp. 157-
2 192.

3

4 Dodds, K. L. and Collins-Thompson, D.L., 1984. Nitrite tolerance and nitrite
5 reduction in lactic acid bacteria associated with cured meat. Int. J. Food Microbiol. 1,
6 163-170.

7

8 Flick, G.J., Hong, G.P. and Knobl, G.M., 1991. Non-traditional methods of seafood
9 preservation. Mar. Technol. Soc. J. 25, 35-43.

10

11 Gancel, F., Dzierszynski, F. and Tailliez, R., 1997. Identification and characterization
12 of *Lactobacillus* spp. isolated from fillets of vacuum-packed smoked and salted
13 herring (*Clupea Hargenus*). J. Appl. Microbiol. 82, 722-728.

14

15 Harper, J.J. and Davis, G.H.G., 1979. Two dimensional thin-layer chromatography for
16 the amino-acid analysis of bacterial cell-walls. Int. J. Syst. Bacteriol. 29, 56-58.

17

18 Hildebrandt, G. and Erol, I., 1988. Sensorische und mikrobiologische Untersuchung
19 an vakuumverpacktem Räucherlachs in Scheiben. Arch. Lebensmittelhyg. 39, 109-
20 132.

21

22 Hyytiä, E., Eerola, S., Hielm, S. and Korkeala, H., 1997. Sodium nitrite and
23 potassium nitrate in control of nonproteolytic *Clostridium botulinum* outgrowth and
24 toxigenesis in vacuum-packed cold-smoked rainbow trout. Int. J. Food Microbiol. 37,
25 63-72.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Jeppesen, V.T. and Huss, H.H., 1993. Characteristic and antagonistic activity of lactic acid bacteria isolated from chilled fish products. *Int. J. Food Microbiol.* 18, 305-320.

Kloos W.E., Schleifer, K.H. and Tornabene, T.G., 1974. Isolation and characterization of micrococci from human skin, including 2 new species: *Micrococcus lyae* and *Micrococcus kristinae*. *Int. J. Syst. Bacteriol.* 24, 79-101.

Knøchel, S. and Huss, H.H., 1984. Ripening and spoilage of sugar salted herring with and without nitrate. II. Effect of nitrate. *J. Food Technol.* 19, 215-224.

Kovács, N., 1956. Identification of *Pseudomonas pyogenea* by the oxidase reaction. *Nature* 178, 703.

Leisner, J.J., Millan, J.C., Huss, H.H. and Larsen, C.M., 1994. Production of histamine and tyramine by lactic acid bacteria isolated from vacuum-packaged sugar-salted fish. *J. Appl. Bacteriol.* 76, 417-423.

Magnússon, H. and Traustadóttir, K., 1982. The microbial flora of vacuum-packed smoked herring fillets. *J. Food Technol.* 17, 695-702.

Mauguin, S. and Novel, G., 1994. Characterisation of lactic acid bacteria isolated from seafood. *J. Appl. Bacteriol.* 76, 616-625.

1 Nieto, T.P., López, L.R., Santos, Y., Núñez, S. and Toranzo, A.E., 1990. Isolation of
2 *Serratia plymuthica* as an opportunistic pathogen in rainbow trout, *Salmo gairdneri*
3 Richardson. J. Fish Dis. 13, 175-177.
4
5 Nordic Committee on Food Analysis, 1986. Aerobic micro-organisms. Enumeration
6 at 30°C in meat and meat products. NCFA method no 86, 2nd ed. Espoo, Finland.
7
8 Pederson, E. and Meyland, I., 1981. Nitrate, nitrite and volatile nitrosamines in
9 pickled fish prepared with addition of nitrate, Z. Lebensm. Unters. For. 173, 359-361
10
11 Pelroy, G.A., Eklund, M.W., Paranjpye, R.N., Suzuki, E.M. and Peterson, M.E., 1982.
12 Inhibition of *Clostridium botulinum* types A and E toxin formation by sodium nitrite
13 and sodium chloride in hot-process (smoked) salmon. J. Food Prot. 45, 833-841.
14
15 Pierson, M.D. and Smoot, L.A., 1987. Nitrite, nitrite alternatives, and the control of
16 *Clostridium botulinum* in cured meats. Crit. Rev. Food Sci. Nutr. 17, 141-187.
17
18 Reuter, G., 1970. Laktobazillen und eng verwandte Mikroorganismen in Fleisch und
19 Fleischerzeugnissen. 2. Mitteilung: Die Charakterisierung der isolierten
20 Laktobazillenstämme. Fleischwirtsch. 50, 954-962.
21
22 Schillinger, U. and Lücke, K.F., 1987. Identification of lactobacilli from meat and
23 meat products. Food Microbiol. 4, 199-208.
24

1 Schleifer, K.H. and Kloos, W.E., 1975. A simple test system for the separation of
2 staphylococci from micrococci. *J. Clin. Microbiol.* 1, 337-338.

3

4 Schneider, W. and Hildebrandt, G., 1984. Untersuchungen zur Lagerfähigkeit von
5 vacuumverpacktem Räucherlachs. *Arch. Lebensmittelhyg.* 35, 49-72.

6

7 Shimasaki, T., Miake, K., Tsukamasa, Y., Sugiyama, M., Minegishi, Y. and Shinano,
8 H., 1994. Effect of water activity and storage temperature on the quality and
9 microflora of smoked salmon. *Nippon Suisan Gakkaishi* 60, 569-576.

10

11 Skovgaard, N., 1992. Microbiological aspects and technological needs: technological
12 needs for nitrates and nitrites. *Food Add. Cont.* 9, 391-397.

13

14 Sofos, J.N., Busta, F.F. and Allen, C.F., 1979. Botulism control by nitrite and sorbate
15 in cured meats: a review. *J. Food Prot.* 42, 739-770.

16

17 Taylor, A.A. and Shaw, B.G., 1975. Whiltshire curing with and without nitrate. *J.*
18 *Food Technol.* 10, 157-167.

19

20 Truelstrup Hansen, L., Gill, T. and Huss, H.H., 1995. Effects of salt and storage
21 temperature on chemical, microbiological and sensory changes in cold-smoked
22 salmon. *Food Res. Int.* 28, 123-130.

23

1 Truelstrup Hansen L., 1995. Quality of chilled, vacuum-packed cold-smoked salmon.
2 Thesis, Danish Institute of Fisheries Research, Department of Seafood Research,
3 Technical University, Denmark.

4

5 Yamanka, T., Ota, A. and Okunuki, K., 1959. A nitrite reducing system reconstructed
6 with purified cytochrome components of *Pseudomonas aeruginosa*. *Biochem.*
7 *Biophys. Acta* 53, 294-308.

8

9

10

11

12

13

14

15

1
 2 Table 1. The four main bacterial groups obtained from spoiled vacuum-packed cold-smoked rainbow
 3 trout fillets produced using different curing methods and stored at 4^o C and 8^o C
 4

Curing method ^a	Storage temperature (°C)	Number of colonies isolated	Microbial group				
			Lactic acid bacteria	Gram-positive, catalase-positive cocci	Gram-negative, oxidase-negative rods	Gram-negative, oxidase-positive rods	Yeast
NaCl	4	99	50 (50%) ^b	12 (12%)	37 (37%)	0	0
NaCl and KNO ₃	4	110	94 (85%)	9 (9%)	7 (6%)	0	0
NaCl and NaNO ₂	4	100	79 (79%)	1 (1%)	19 (19%)	1 (1%)	0
NaCl	8	104	81 (80%)	5 (5%)	18 (17%)	0	0

NaCl and KNO ₃	8	108	68 (63%)	12 (11%)	17 (16%)	11 (10%)	0
NaCl and NaNO ₂	8	99	97 (97%)	0	0	0	2 (2%)

- 1
- 2 ^a Concentration of NaCl: 2.2%, KNO₃: 686 ppm and NaNO₂: 166 ppm.
- 3 ^b Proportion of strains isolated from samples with specified curing method and storage temperature.
- 4
- 5
- 6

1
2
3
4
5
6
7
8

Table 2. Characterisation of lactic acid bacteria isolated from spoiled vacuum-packed cold-smoked rainbow trout fillets produced using different curing methods and stored at 4^o C and 8^o C

Characteristic	Subgroups and number of isolates in parenthesis													
	1 (189)		2 (50)		3 (3)		4 (182)		5 (3)		6 (17)		7 (25)	
Morphology	Rods		Oval cocci		Rods		Oval cocci		Cocci		Cocci or oval cocci		Cocci or oval cocci	
Production of gas from glucose	-		+		+		+		-		+		-	
Growth on SL agar	+		-		-		+		-		-		-	
Lactic acid isomere ^a	DL /D(L) ^b		D(L)		D		DL /D(L)		L		DL		DL /L(D)	
Hydrolysis of arginine	+/- ^c		-		+		+/-		+/-		+/-		+/-	

Production of acetoin	-	-	+	-	+/-	+/-	+/-
m-DPA ^a	+/-	-	+	-	-	-	-

- 1 ^a Ten or all strains analysed from each group.
- 2 ^b Parenthesized isomers indicate < 15% of total lactic acid.
- 3 ^c +/- : positive or negative reactions.
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17

1 Table 3. Distribution of the lactic acid bacteria isolated from spoiled vacuum-packed cold-smoked
 2 rainbow trout fillets produced using different curing methods and stored at 4° C and 8° C
 3

Subgroup of lactic acid bacteria	No. of isolates	Storage temperature (°C)					
		4			8		
		NaCl ^a	NaCl and KNO ₃ ^a	NaCl and NaNO ₂ ^a	NaCl ^a	NaCl and KNO ₃ ^a	NaCl and NaNO ₂ ^a
1	189	8 (2%) ^b	56 (12%)	37 (8%)	9 (2%)	59 (12%)	20 (4%)
2	50	11 (2%)	2 (0.5%)	1 (0.5%)	8 (2%)	2 (0.5%)	26 (5%)
3	3	0	0	1 (0.5%)	1 (0.5%)	1 (0.5%)	0
4	182	27 (6%)	36 (7%)	40 (9%)	24 (5%)	5 (1%)	50 (11%)
5	3	0	0	0	3 (1%)	0	0
6	17	1 (0.5%)	0	1(0.5%)	13 (3%)	1 (0.5%)	1(0.5%)
7	25	3 (1%)	0	0	22 (5%)	0	0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

^a Concentration of NaCl: 2.2%, KNO₃: 686 ppm and NaNO₂: 166 ppm.

^b Percentage of all strains in isolated different curing method at each storage temperature.

1
2 Table 4. Distribution of the Gram-negative, oxidase-negative rods isolated from spoiled vacuum-packed cold-smoked rainbow trout fillets produced using
3 different curing methods and stored at 4° C and 8° C
4

Microorganisms	No. of isolates	Storage temperature (°C)					
		4			8		
		NaCl ^a	NaCl and KNO ₃ ^a	NaCl and NaNO ₂ ^a	NaCl ^a	NaCl and KNO ₃ ^a	NaCl and NaNO ₂ ^a
<u>Serratia plymuthica</u>	42	32	3	0	5	2	0
<u>Hafnia alvei</u>	9	0	0	1	2	6	0
<u>Enterobacter amnigenus</u>	8	0	0	0	0	8	0
<u>Xanthomonas maltophilia</u>	9	0	0	9	0	0	0
<u>Serratia liquefaciens</u>	7	3	0	0	4	0	0
<u>Enterobacter sakazakii</u>	5	0	0	0	5	0	0
<u>Morganelli morganii</u>	4	0	0	4	0	0	0
<u>Rahnella aquatilis</u>	4	0	3	0	0	1	0
<u>Citrobacter freundii</u>	3	0	0	3	0	0	0
<u>Serratia odofera</u>	1	0	0	0	1	0	0
<u>Enterobacter agglomerans</u>	1	1	0	0	0	0	0
<u>Escherichia fergusonii</u>	1	0	0	0	1	0	0
<u>Erwinia sp.</u>	1	1	0	0	0	0	0
<u>Acinetobacter sp.</u>	1	0	0	1	0	0	0

Unidentified	2	0	1	1	0	0	0
--------------	---	---	---	---	---	---	---

1

2 ^a Concentration of NaCl: 2.2%, KNO₃: 686 ppm and NaNO₂: 166 ppm.