



UNIVERSITY OF HELSINKI

<https://helda.helsinki.fi>

Protein – a scoping review for Nordic Nutrition Recommendations 2023

Geirsdóttir, Ólöf Guðný; Pajari, Anne Maria

2023

Swedish Nutrition Foundation

<http://hdl.handle.net/10138/569312>

Geirsdóttir, Ó G & Pajari, A M 2023, 'Protein – a scoping review for Nordic Nutrition Recommendations 2023', Food and Nutrition Research, vol. 67, 10261. <https://doi.org/10.29219/fnr.v67.10261>

Downloaded from Helda, University of Helsinki institutional repository. <https://helda.helsinki.fi>
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.
Please cite the original version.

REVIEW ARTICLE

Protein – a scoping review for Nordic Nutrition Recommendations 2023

Ólöf Guðný Geirsdóttir^{1*} and Anne-Maria Pajari²

¹Faculty of Food Science and Nutrition, School of Health Science, University of Iceland, Reykjavik, Iceland;

²Department of Food and Nutrition, University of Helsinki, Helsinki, Finland

Popular scientific summary

- Dietary proteins provide essential amino acids, nitrogen, and energy.
- Average requirements for protein intake are based on the maintenance of nitrogen balance, accounting for growth in children, pregnancy, and lactation.
- The amount and bioavailability of essential amino acids determines the protein quality of foods.
- Current intakes in Nordic and Baltic countries range from 15 to 19% of total energy intake.

Abstract

Proteins are needed for providing essential amino acids, nitrogen, and fuel for the body's needs in all age groups. Proteins are especially required during active growth in pregnancy, lactation, childhood, and tissue growth in general. An adequate protein intake is needed in old adults to avoid premature muscle loss. According to the current dietary surveys, protein intake in the Nordic and Baltic countries varies from 15 to 19% of the total energy intake in adults. Comprehensive data regarding children and older adults are lacking. No good measure for protein status exists, and the estimation of physiological requirements is based on N-balance studies having some weaknesses. Protein quality is assessed by considering the protein digestibility of individual indispensable amino acids and their utilization (bioavailability), which is affected by food antinutrients and processing. The evidence regarding the association of protein intake per se with health outcomes is limited or suggestive. It is difficult to separate from the effect of other nutrients or ingredients in protein-rich foods. Proteins are widespread in foods, deriving from both animal and plant sources. Animal-sourced protein production puts more strain on the environment than plant-sourced proteins and contributes significantly to greenhouse gas emissions, thereby enhancing climate change. In Nordic and Baltic countries, consumption of animal-sourced proteins is relatively high. A shift toward more plant-based protein diets would be advisable for promoting a healthy and sustainable diet.

Keywords: *dietary protein; amino acids; protein recommendations; essential amino acids; nitrogen balance; protein intake; protein turnover; protein bioavailability; protein quality*

Received: 1 December 2022; Revised: 16 May 2023; Accepted: 27 September 2023; Published: 12 December 2023

Proteins are found in many food sources, including both animals and plants. Amino acids are essential for constructing proteins and various cellular structures. In nutrition, dietary protein plays two significant roles: providing a source of nitrogen (N) and amino acids and supplying energy in a non-specific manner. Proteins are especially required during active growth in pregnancy, lactation, childhood, tissue and wound healing, and tissue building in sports. National surveys conducted using varying methods in each country have revealed that protein

consumption among adults in the Nordic and Baltic regions ranges from 15 to 19% of total energy intakes (E%) (1). Comprehensive systematic data from all Nordic and Baltic countries regarding children and older adults are lacking. Nevertheless, epidemiological studies with limited coverage have been conducted on children and elderly individuals (2–6). No good measure for protein status exists, and the estimation of physiological requirements is based on N-balance studies having some weaknesses (7). Protein quality is assessed considering the protein

digestibility of individual indispensable amino acids and their utilization (bioavailability), which is affected by food antinutrients, natural or synthetic compounds in foods that interfere with the absorption of nutrients. The evidence regarding the association of protein intake per se with health outcomes is limited or suggestive. It is difficult to separate the effect of protein from the effect of other nutrients or ingredients in protein-rich foods.

In individuals with energy balance and moderate physical activity, the protein requirement is the lowest protein intake to maintain N-balance (8). For growing children, pregnant and lactating women, or another anabolic state, the protein requirement is taken to include the needs associated with the deposition of tissues at rates consistent with good health. In the Nordic Nutritional Recommendation (NNR), the energy content from protein in a mixed diet is calculated as 17 kJ/g.

Both animal- and plant-based foods serve as dietary protein sources. Meat, fish, milk, and eggs are major animal protein sources with specific protein content ranging from 15 to 30% (wet weight), while cereals, legumes, nuts, and seeds represent primary plant protein sources with protein content varying from 5 to 15% (dry matter) in cereals to 17 to 40% (dry matter) in legumes.

Traditionally, a nitrogen-to-protein conversion factor of 6.25 is applied to determine protein content in most foods and food ingredients. It is based on two assumptions: that the total mass of proteins contains about 16% N by weight and that all N in food is derived from protein (7). The N content of various amino acids ranges from 7.7 to 32.2%, and the N content of protein in individual foods depends on the amino acid composition. Thus, the conversion factor can vary from 5.70 for soya to 6.32 for milk. Therefore, using the same conversion factor for all protein sources can lead to significant errors in the actual protein content of most foods. Foods also contain various amounts of non-protein nitrogenous compounds, which is an additional source of error. Multiplying total N content with the default value of 6.25 to quantify protein is imperfect and can lead to a 15–20% error in the actual protein content. Protein source-specific nitrogen-to-protein conversion factors have been presented and used over the years. Still, there are no standardized methods to determine them (8), which may be why most trade and food regulatory bodies, including European Union (EU) labeling legislation, continue to use the default value of 6.25.

Amino acids from dietary proteins are classified as essential amino acids that cannot be synthesized in the human body or non-essential amino acids synthesized within the body from other amino acids (transamination) if there is an adequate N supply (Table 1).

Histidine is considered essential, although it does not fulfill the criterion of reducing protein deposition and inducing negative nitrogen balance when removed from the diet. Conditionally essential amino acids are amino acids whose synthesis requires the availability of another amino acid either as the carbon donor or as the donor of an accessory group, for example, the sulfur group of cysteine (9). Under normal conditions, conditionally essential amino acids are synthesized in sufficient amounts. Still, during certain conditions, such as prematurity (10) and illness (11–13), the synthesis might not support all the body’s metabolic needs, which means they can become essential amino acids.

This scoping review aims to describe the totality of the evidence regarding the role of protein in health-related outcomes as a basis for setting up and updating dietary reference values (DRVs).

Methods

This scoping review follows the protocol developed within the NNR2023 (14). The sources of evidence used in the review follow the eligibility criteria described in the paper ‘The Nordic Nutrition Recommendations 2022 – Principles and Methodologies’, published in Food & Nutrition Research (15).

For the NNR2023 update, a literature search on proteins was performed according to principles and methodology for the NNR2023, resulting in 115 systematic reviews (SRs); 38 SRs were considered relevant for three topics for SR. The literature search included the following terms: (protein[Title] AND (systematic review[Publication Type] AND (“2011”[PDAT] : “3000”[PDAT]))) AND Humans[Filter]) AND ((“diet”[MeSH Terms] OR “diet”[All Fields]) OR (“diet”[MeSH Terms] OR “diet”[All Fields] OR “dietary”[All Fields]) OR (“food”[MeSH Terms] OR “food”[All Fields]) OR (“nutritional status”[MeSH Terms] OR (“nutritional”[All Fields] AND “status”[All Fields]) OR “nutritional status”[All Fields] OR “nutrition”[All Fields] OR “nutritional sciences”[MeSH Terms] OR (“nutritional”[All Fields] AND “sciences”[All Fields]) OR “nutritional sciences”[All Fields]) OR Nutritional[All Fields])).

Table 1. Amino acids classification

Essential amino acids	Non-essential amino acids	Conditionally essential amino acids
Isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, and histidine	Alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, proline, serine, and tyrosine	Arginine, cysteine, glutamine, glycine, proline, and tyrosine

Physiology and metabolism

For the body to utilize dietary proteins, they must be broken down into small peptides and amino acids. Protein digestion starts in the stomach with the denaturation of polypeptides by stomach acids and partial hydrolysis of peptide linkages by the enzyme pepsin (16). The polypeptides enter the duodenum, where they are subjected to pancreatic proteases and hydrolyzed to oligo-, tri-, and dipeptides and, to a lesser extent, to free amino acids. Peptide fragments are further hydrolyzed by brush border-bound aminopeptidases and carboxypeptidases on the luminal surface of enterocytes. The resulting free amino acids, dipeptides, and tripeptides are then absorbed through different transport pathways into an enterocyte, where cytosolic aminopeptidases hydrolyze the remaining di- and tripeptides to free amino acids. These amino acids enter the bloodstream and are incorporated into tissue protein and other N-containing compounds such as neurotransmitters, hormones, creatinine, and drug elimination ligands. This makes dietary protein requirement a requirement for amino acids and N.

Protein quality is primarily characterized by its essential amino acid content and the digestibility and availability of the resulting amino acids in the circulation. These factors influence their metabolism within different body protein pools (17). Hence, determining protein quality is based on protein digestibility ranking methods such as the protein digestibility-corrected amino acid score (PDCAAS) used earlier and the digestible indispensable amino acid score (DIAAS) currently recommended by the Food and Agriculture Organization of the United Nations (FAO) (7, 18). The significant differences between the two approaches are that DIAAS uses ileal digestibility instead of fecal digestibility and does not truncate the values artificially to 100%. DIAAS is defined as:

$$\text{DIAAS \%} = 100 \times (\text{mg of digestible dietary indispensable amino acid in 1 g of the dietary protein} / \text{mg of the same nutritional indispensable amino acid in 1 g of the reference protein}).$$

For calculating DIAAS, it is recommended to use actual ileal digestibility values of individual amino acids, preferably measured in humans, but if not available, then values obtained in pigs or rats. As measuring ileal digestibility is highly invasive, human data still need improvement (19). Table 2 shows DIAAS values and limiting AA for various animal and plant-based protein sources (20).

Bioavailability comprises the proportion of the amino acid in a utilizable form and the exclusion of metabolism-interfering compounds in the food. Unprocessed plant protein sources contain antinutrients such as phytates, tannins, and protease inhibitors, which interfere with the digestion of plant proteins, making them less well-digestible and less bioavailable than animal-sourced proteins (21). Antinutrients

Table 2. DIAAS % and limiting AA for animal and plant proteins (20)

Protein source	DIAAS %	Limiting AA
Corn	43	Lysine
Rice	56	Lysine
Wheat	56	Lysine
Hemp	56	Lysine
Faba bean	64	Methionine + cysteine
Oat	68	Lysine
Rapeseed	79	Lysine
Lupin	83	Methionine + cysteine
Pea	83	Methionine + cysteine
Canola	85	Lysine
Soy	103	NA
Potato	125	NA
Whey	106	NA
Egg	111	NA
Casein	137	NA
Pork	126	NA

partly explain the low DIAAS values of some plant protein sources in Table 2. The bioavailability of plant proteins varies and is generally lower than that of animal proteins. Soy and potato protein represents a good-quality plant protein with all essential amino acids, followed by legumes low in sulfur-containing amino acids and cereals low in lysine. Food matrix and food processing, for example, heating, fermentation, and extrusion, can also impact amino acid bioavailability, either increasing or decreasing it (22). It is worth emphasizing that the digestibility of individual amino acids in the same food source differs. Therefore, FAO recommends treating each amino acid as an individual nutrient when evaluating protein quality (7).

In practice, the differences in quality between proteins might be less critical in diets containing a variety of protein sources, such as in a typical Nordic mixed diet. Dietary proteins of animal origin (meat, fish, milk, and eggs) or a combination of plant proteins from, for example, legumes and cereal grains give a good distribution of essential amino acids. With the current Nordic relatively high protein intake, replacing some animal proteins with plant proteins would probably lead to lower protein intake and lower bioavailability, but still provide enough protein and essential amino acids (23). However, protein digestibility and bioavailability may become an issue in protein transition toward more plant-based diets in vulnerable groups such as older adults, as protein bioavailability is known to decrease with age (21).

Dietary protein sources contain not only proteins but also provide other critical nutrients for health. In randomized controlled trials (RCTs), replacing animal-sourced proteins with plant-based ones in a typical Nordic diet resulted in favorable changes such as increased intakes of fiber and unsaturated fatty acids and decreased intakes of

saturated fatty acids and cholesterol, which were accompanied by a decrease in blood Low density lipoprotein (LDL) cholesterol (23). However, the replacement considerably decreased intakes of vitamin B₁₂ and iodine (24), and markers of bone health indicated an enhanced bone turnover with a possible increased risk for fractures (25). When modeling with population data with a more moderate change of only replacing red and processed meat with legumes, increases in the average intakes of fiber, folate, K, Mg, Cu, and Fe were observed. In contrast, intakes of saturated fat, niacin, vitamin B₁₂, Se, and Zn were decreased (26). Overall, distribution shifts toward a higher probability of inadequate intakes of the studied nutrients were not observed.

The efficiency of the anabolic effects of dietary amino acids on body and muscle protein depends on the quantity and bioavailability of the amino acids and on the pattern of essential amino acids, which should cover the needs of all tissues. Furthermore, the kinetics of the delivery of amino acids to muscle tissues also appears to be an essential factor in modulating the anabolic efficiency of dietary amino acids in muscle protein synthesis (11). It is assumed that the postprandial rise in circulating amino acids represents the main driver of the muscle protein synthetic response to feeding. A high rate of protein digestion, amino acid absorption, and blood amino acid delivery has been previously observed with soluble proteins such as whey protein or with different hydrolyzed proteins with a high rate of digestion and amino acid delivery (18).

Some studies suggested that a spread feeding pattern with about 30 g of dietary protein during the main meals (i.e. breakfast, lunch, and dinner) could be a more effective strategy to counteract age-related muscle atrophy and strength loss in older adults (27, 28). The results of

longitudinal studies have sparked debate as they indicate that while a varied protein intake can increase muscle mass and strength, it may not necessarily improve mobility over 2 years.

Body proteins, all proteins in the body, are continually broken down and synthesized. The protein turnover of about 300 g per day in adults (29) is higher than the proteins consumed from the diet (Fig. 1). This indicates an extensive reutilization of amino acids in protein metabolism. N from the amino acids leaves the body via the urine, mainly urea, uric acid, and creatinine. N is also lost in feces, sweat, and other secretions through the skin, hair, and nails. Amino acids are needed to compensate for these losses, and amino acids are also required for protein synthesis during anabolism (20), for example, during active growth in pregnancy, lactation, childhood, tissue and wound healing, and tissue building in sports. It is usually assumed that almost all dietary N is incorporated into protein. N-balance is the difference between nitrogen intake and nitrogen output. A negative N-balance, that is, losses greater than intake, is seen during fasting, starvation, and catabolic diseases like cancer, heart failure, and chronic obstructive pulmonary disease. On a long-term basis, a healthy adult should be in N equilibrium, that is, intake and losses should be equal.

Assessment of nutrient status

There are no specific blood tests for protein deficiency or other excellent or valid measures for protein status in healthy or subclinical malnutrition. Where plasma albumin and other plasma proteins decrease in severe malnutrition, hypoalbuminemia's etiology is multifactorial and not only related to nutrition (31). Clinical studies suggest that the underlying cause of hypoalbuminemia, rather




Fig. 1. An overview of turnover of proteins and amino acids in the body, modified from Millward (30).

than solely low albumin levels, is accountable for the associated morbidity and mortality (32, 33).

N-balance has been used to establish dietary protein recommendations, but its use is controversial. This methodology indirectly determines protein turnover, and no information about whole-body N, protein turnover, or various protein metabolic pathways can be obtained. It is important to consume equicaloric diets to meet nutritional needs. The World Health Organization (WHO) has examined stable isotope studies to determine the necessary intake of essential amino acids (34). Their research shows that when the oxidation of essential amino acids increases, the intake exceeds the requirements. Therefore, it is crucial to avoid low protein intake as it may lead to underestimation of our needs by inducing protein sparing (34, 35). Over the last several years, more direct methods of measuring the turnover of various body proteins have been developed, including stable isotope tracer methods. This has enabled a mechanistic approach to the effects of different dietary proteins (36). Similar logical reasoning cannot be applied to whole-body protein turnover beyond what can already be deduced from N-balance studies. Studies of muscle protein turnover have yet to add to an understanding of muscle function because no studies are available demonstrating a correlation between, for example, muscle strength

or endurance and the dynamics of muscle protein turnover. Thus, in the future, it will be essential to use more advanced methodologies in strictly controlled long-term studies to establish mechanistic links between health outcomes and protein intake from various sources, as well as for different age groups and health conditions.

Dietary intake in Nordic and Baltic countries

According to national dietary surveys, the average protein intake among adults is high in the Nordic and Baltic countries, ranging from 15 E% in females in Denmark to 18 E% in Iceland, Norway, and Finland (1) (Tables 3a, 3b). Dietary protein intake in the Nordic and Baltic countries is mainly with high-quality protein where protein contains all essential amino acids in the right proportion required by the human body, that is, a protein with high biological value. When comparing the intakes between the countries, it needs to be mentioned that while these are the most recent reliable data on the national consumption of adults, the national dietary surveys have used different methods. Therefore, the values should be compared with caution (1).

The average protein intake among children varies between 13 and 16 E% from about 1 year of age in most European countries, including the Nordic countries (1) (Tables 4a, 4b).

Table 3a. Daily mean intakes of protein among adults and children in Nordic countries (1)

	Denmark 2011 (18–75 years)		Finland 2017 (18–74 years)		Iceland 2019–2021 (37) (18–80 years)		Norway 2010 (18–70 years)		Sweden 2010 (18–80 years)	
	Men (n = 1,464)	Women (n = 1,552)	Men (n = 780)	Women (n = 875)	Men (n = 632)	Women (n = 680)	Men (n = 862)	Women (n = 925)	Men (n = 792)	Women (n = 1,005)
Protein g/day	101	76	98	73	104	76	112	81	92	72
Protein E%	16	15	18	18	19	18	18	18	17	17
	Men age 65–75 years	Women age 65–75 years	Men age 65–74 years	Women age 65–74 years	Men age 61–80 years	Women age 61–80 years	Men age 61–80 years	Women age 60–70 years	Men age 65–80 years	Women age 65–80 years
Protein g/day	95	74	76	66	96	78	102	77	84	70
Protein E%	15	15	17	17	19	18	18	18	17	17

Table 3b. Daily mean intakes of protein among adults and children in Baltic countries (1)

	Estonia 2014 (age 18–74 years)		Latvia 2018 (age 19–64 years)		Lithuania 2019 (age 19–75 years)	
	Men (n = 907)	Women (n = 1,806)	Men (n = 470)	Women (n = 541)	Men (n = 1,348)	Women (n = 1,562)
Protein g/day	86.5	63.2	104.4	75.1	78.7	62.8
Protein E%	16.6	16.2	17.4	17.0	15.6	16.0
	Estonia 2014 (age 70–74 years)		Latvia 2018 (age 50–64 years)		Lithuania 2019 (age 65–75 years)	
	Men	Women	Men	Women	Men and women	
Protein g/day	74.6	55.3	100.8	77.1	57.2	
Protein E%	16.7	16.5	17.8	18.1	15.1	

Table 4a. Protein intake among children in Nordic countries

Denmark			Finland			Norway		Sweden		
Age (years)	Boys	Girls	Age (years)	Boys	Girls	Age (years)	Both boys and girls	Age (years)	Boys	Girls
4–5	66 ^a	58 ^a	3–4	56 ^a	51 ^a					
	14 ^b	14 ^b		17 ^b	17 ^b					
6–9	74 ^a	67 ^a	5–6	61 ^a	55 ^a					
	14 ^b	14 ^b		16 ^b	16 ^b					
10–13	85 ^a	70 ^a				10–11	68 ^a	12	85 ^a	78 ^a
	15 ^b	15 ^b					16 ^b		18 ^b	17 ^b
14–17	96 ^a	63 ^a				14–15	75 ^a	15	104 ^a	79 ^a
	16 ^b	15 ^b					16 ^b		17 ^b	16 ^b
								18	112 ^a	79 ^a
									18 ^b	16 ^b

^ag/d.^bE%.**Table 4b.** Protein intake among children in Baltic countries

Estonia			Lithuania	
Age (years)	Boys	Girls	Age (years)	Both boys and girls
2–5	50.6 ^a	47.6 ^a		
	13.9 ^b	14.0 ^b		
6–9	66.8 ^a	56.0 ^a	7–10	56.3 ^a
	14.6 ^b	13.8 ^b		15.9 ^b
10–13	73.2 ^a	54.0 ^a	11–14	65.2 ^a
	14.6 ^b	13.6 ^b		14.2 ^b
14–17	84.5 ^a	56.3 ^a	15–18(19)	70.5 ^a
	15.1 ^b	14.5 ^b		14.2 ^b

^ag/d.^bE%.

Protein intake and health outcomes relevant for Nordic and Baltic countries

Severe protein deficiency results in edema, muscle weakness, and changes to the hair and skin. Protein deficiency is often linked to energy deficiency, protein-energy malnutrition, and other nutrients based on a general nutrition deficiency.

Obesity

Many meta-analyses of short-term studies indicate favorable effects of higher versus lower protein diets on health outcomes like adiposity (38). A meta-analysis of RCTs was done in 2013 on long-term effects (≥ 12 months) of low- or high-protein diets on cardiovascular and metabolic risk factors (39). It showed that high-protein diets exerted neither specific beneficial nor detrimental effects on obesity.

The ‘early protein hypothesis’ established by Rolland-Cachera suggests that consuming a high amount of protein during infancy can potentially lead to an increased risk of obesity (40). The composition of macronutrients in infant formula differs from that in human milk.

Protein consumption stimulates the production of insulin-like growth factor 1 (IGF-1) and insulin. Studies have shown an inverse correlation between IGF-1 levels in infancy and those in late adolescence (41). The variety of proteins ingested can have varying effects on IGF-1. In particular, the consumption of cow’s milk during early infancy has been shown to enhance linear growth and increase circulating IGF-1 levels in well-nourished populations (42). Research has shown that consuming poor-quality protein without animal protein is more likely to lead to non-communicable diseases (NCDs) later in life than consuming high amounts of total protein. Additionally, children who have a higher BMI for their age or become obese during childhood are at an increased risk of developing adult obesity and associated health problems (43).

Cardiovascular diseases and diabetes

Cardiovascular disease (CVD) and type 2 diabetes (T2D) are significant causes of morbidity and mortality worldwide and are associated with high societal costs (3).

Based on three SRs of cohort studies by Tian et al. (44), Virtanen et al. (45), and Chen et al. (46), total and animal protein intake increased the risk of T2D, whereas plant protein decreased the risk of T2D. Again, it should be emphasized that low carbohydrate or high-protein diets cannot be used to assess the effect of protein per se.

The association between dietary protein intake from different sources of proteins and the risk of CVD in an SR of prospective studies is of interest where some studies showed no association between dietary protein intake and CVD risk from different sources of proteins (47, 48). According to a meta-analysis done in 2023, there were no observed cardiovascular effects resulting from a high-protein diet in which protein accounted for 18% or more of energy intake (49). Still, in subgroup analysis, there was a decreased risk of CVD mortality with an increasing plant protein intake. A *de novo* SR for NNR2023 by Lamberg et al. (50) concluded that evidence that substituting animal protein with plant protein reduces the risk of CVD mortality or T2D incidence is *limited – suggestive*.

Replacing animal protein with plant protein for nutrients, fibers, and sustainability should also be a public health strategy to lower the risk of CVD mortality and T2D.

Bone health

The role of dietary protein on bone health has been controversial. Urinary calcium loss increases in high protein intakes, but at the same time, protein increases calcium absorption and bioavailability. These seemingly contradictory effects make it uncertain what the net effect of a high-protein diet (typically >20 E% and/or >1.0 g/kg body weight [BW]) is on calcium metabolism and bone health (51). Few studies have shown that protein intake above 0.8 g/kg BW prevents bone loss and hip fracture. However, the European Food Safety Authority (EFSA) (52) found that evidence is inconclusive regarding protein and bone health (53). However, there is an interaction between low calcium (<800 mg/day) intake and protein intake, where an increased risk of fractures was related to high animal protein intake. However, under high calcium intake (>800 mg), a high animal protein intake was related to a decreased risk of fractures (25). This finding is supported by an older Norwegian study of 39,787 middle-aged men and women that showed an elevated risk of hip fractures in women with high animal (non-dairy) protein intake under low calcium intake (54). The evidence for an association between vegetable protein intake and fracture risk is inconclusive (53, 55) and this finding was supported by an older study of 32,050 postmenopausal women that showed a decreased risk of hip fracture related to a high animal protein intake, but not to vegetable protein intake (56). In older adults, Pedersen and Cederholm (57) assessed the evidence as suggestive regarding a positive

association between protein intake and bone mineral density based on one intervention study and three prospective cohort studies. The evidence was assessed as inconclusive regarding the relation of protein intake to bone loss and the risk of fractures. Interestingly, in the included randomized controlled study with calcium (500 g/day) and vitamin D supplementation (17.5 g/day) by Dawson-Hughes and Harris (53), the highest tertile of protein intake (20 E%, or 1.2 g/kg BW) was associated with less bone loss compared to the lowest tertile (14 E%, or 1.1 g/kg BW), but only in the intervention group. The habitual mean intake in the placebo group was 871 mg of calcium and about 7 µg vitamin D daily compared to 1,346 mg per day calcium in the intervention group. Thus, the possible effect of protein intake on bone health might depend on an input of calcium and vitamin D above this level.

Renal function and kidney stones

High-protein diets have been associated with increased Glomerular filtration rate (GFR), serum urea, urinary calcium excretion, and serum concentrations of uric acid. Some results have indicated caution in recommending high-protein diets. However, the evidence for associations between protein intake and kidney function and kidney stones in healthy people needs to be more conclusive (35, 56, 58, 59). Also, the EFSA (52) concluded that no maximum protein intake level could be established for those with healthy kidney function due to insufficient evidence.

Cancer

A meta-analysis from 2017 of 21 observational studies comprising 8,187 cases by Lai et al. (60) found no association between protein intake or protein source (animal vs. plant) and colorectal cancer risk. The authors reported a significant between-study heterogeneity, which may have affected the results. Liao et al. (61) utilized a large prospective US cohort comprising nearly a half-million adults with a median follow-up of 15.5 years to evaluate the effect of substituting plant protein for animal protein on the risk of colorectal cancer. The substitution was associated with up to an 11% decreased risk of colorectal cancer, primarily due to substituting plant proteins for red meat, not white meat, dairy, or egg protein. Further analyses revealed that the risk reduction was mostly limited to replacing protein from bread, cereal, and pasta for red meat protein and was statistically significant only for the distal colon and rectum. These results align with other studies and the latest World Cancer Research Fund's food-based evaluation, concluding that solid scientific evidence exists, suggesting that high intakes of red and processed meat increase, and high intakes of dairy products and whole grains decrease, the risk of colorectal cancer (62).

A meta-analysis of 11 studies with 2,537 cases and 11,432 participants by Kong et al. (2020) (63) found no

association between overall dietary protein intake and esophageal cancer risk. Still, a sub-analysis revealed an increased risk for esophageal squamous cell carcinoma.

No association was found between prostate cancer and dietary protein intake, or animal or plant protein intake in a meta-analysis of 12 studies comprising 13,483 prostate cancer cases and 286,245 participants (64). The authors claim no publication bias or between-study heterogeneity.

A meta-analysis of 46 prospective studies by Wu et al. (65) examined the association between dietary protein sources and breast cancer risk. They found that a higher intake of soy food and skim milk could decrease the risk and that a higher intake of processed meat may increase the risk of breast cancer.

Most studies on the relationship between protein intake and cancer are food-based and, therefore, cannot isolate the effect of the protein intake per se from other nutrients or ingredients in foods. Furthermore, while plant-based foods are commonly associated with a somewhat lower risk of cancers, different animal-based foods may have opposite effects on cancers, as demonstrated by red and processed meat and dairy products in colorectal cancer (62).

Mortality

Several meta-analyses found that a high animal protein intake was positively associated with cardiovascular mortality. A high plant protein intake was inversely associated with all-cause and cardiovascular mortality, especially among individuals with at least one lifestyle risk factor. The substitution of plant protein for animal protein, especially that from processed red meat, was associated with lower mortality, suggesting the importance of protein sources (48, 66–70). An SR of cohort studies from 2020 on the connection between protein sources and mortality yielded inconsistent findings (68). Nevertheless, it has been confirmed that older men (mean age 74 years) who do not consume enough protein (<0.8 g/kg/day), regardless of its source, have a slightly higher risk of all-cause and cause-specific mortality (71). Based on current literature, replacing animal protein with plant protein may have health benefits (72).

Requirement and recommended intakes

The metabolism of proteins in humans is closely related to energy intake (73). The WHO/FAO/UNU (34) define the protein requirement of an individual as ‘the lowest level of dietary protein intake that will balance the losses of nitrogen from the body, and thus maintain the body protein mass, in persons at energy balance with modest levels of physical activity, plus, in children or pregnant or lactating women, the needs associated with the deposition of tissues or the secretion of milk at rates consistent with good health’. Despite limitations in the method that

are mainly related to the accuracy of the measurements and interpretation of the results, N-balance remains the method of choice for determining the protein requirement in adults in the absence of validated or accepted alternatives and the lack of a reliable biological marker of protein status.

Recommended intake

Adults

The EFSA (52) states that 0.83 g of good-quality protein/kg BW per day based on an estimated average requirement (EAR) of 0.66/kg BW per day covers daily protein needs. The 2002 US recommendations from the Institute of Medicine (now the National Academies of Sciences, Engineering, and Medicine, NASEM) (74) for protein were also based on the meta-analysis of N-balance studies by Rand et al. (75) and cite an EAR of 0.66 g/kg BW per day and a recommended daily allowance (RDA) of 0.8 g good-quality protein/kg BW per day. These recommendations are for healthy adults based on a coefficient of variation of 12% and with no significant differences according to adult age or sex.

An SR was conducted to update the NNR2012 (57) to evaluate the potential health impacts of different protein intake levels. The evidence regarding protein intake was largely inconclusive for most outcomes, including all-cause mortality, cancer mortality and diseases, CVD, bone health, BW control, body composition, and renal function. However, the review identified a potential relationship between plant-based proteins and lower cardiovascular mortality and blood pressure rates. Thus, despite some studies finding a decreased risk of outcomes associated with vegetable protein intake (76), the authors concluded there was insufficient evidence to recommend an increased intake of plant-based sources.

Based on Nordic and Baltic dietary habits and available evidence, the current intake falls within the recommended range of 10–20% of daily energy intake (1). This protein intake adequately meets the requirements for essential amino acids. With decreasing energy intake below 8 MJ (e.g. due to decreased physical activity or during intentional weight loss), the protein E% should increase accordingly.

Infants and children

Good nutrition is vital for healthy growth and development during the first 2 years of life. Starting good nutrition practices early can help children develop healthy dietary patterns. Regarding the later risk of NCDs such as CVD, the quantity and quality of protein intake in infancy and childhood are of interest.

Breastfeeding is associated with a reduced later obesity risk relative to feeding conventional infant formula. Breastfeeding induces less weight gain during the first 2

years of life, which predicts less obesity up to adulthood (77). During the first 6 months, infants are breastfed or receive infant formula. The protein content of breast milk is considered adequate for term infants, and the protein content of infant formula is regulated by the EU legislation. According to the current regulation/directive (REGULATION [EU] No 609/2013 and COMMISSION DIRECTIVE 2013/46/EU), the protein content of infant formula should be between 0.45 and 0.7 g/100 kJ, and the protein content of follow-on formula should be between 0.45 and 0.8 g/100 kJ.

Concerning BW, the WHO/FAO/UNU (78) gives reference values of 0.9 g/kg BW per day from 3 to 18 years of age for boys and from 3 to 15 years of age for girls. This value decreases slightly for girls to 0.8 g/kg BW per day between 15 and 18 years of age. The protein energy percentage necessary to cover the adequate protein intake can be calculated by combining these reference values with the reference values for energy intake for age and sex. The average requirement calculated as E% is about 5.3 E% at 6 months, followed by a decline to 4.3 E% at 2 years. After that, there is a gradual increase. The recommended safe level has decreased for all ages, especially for the first 2 years (Table 5).

Expressed as E%, the protein intake increases considerably during the first 1–2 years of life when the infant gradually changes from breast milk with about 5 E% to the family diet that typically provides around 15 E% from protein. The average protein intake among children varies between 13 and 16 E% from about 1 year of age in most European countries, including the Nordic and Baltic countries (52). In the Nordic setting, quantity is more important than quality because the protein sources are usually of animal origin.

The appropriate upper limit for protein intake in infancy and childhood has not yet been firmly established, and there is interest in understanding the short- and long-term effects of different levels of protein intake during this period. Research has focused on growth, serum lipids, glucose and insulin, blood pressure, BW, body composition, and bone mineral density. While some studies suggest that higher protein intake during infancy and early childhood may increase the risk of obesity later in life (79, 80), results have been inconclusive for other long-term health parameters (81). Available data show that a protein intake between 15 and 20 E% during the first 2 years of life may increase the risk of being overweight later in life (43).

Table 5. A safe level of protein intake (average requirement + 1.96 SD) in weaned infants and children (52)

Age	Protein g/kg BW	E%	g/100 kJ
6–11 months	1.1	7–15	0.4–0.9
12–23 months	1.0	10–15	0.6–0.9
2–17 years	0.9	10–20	

There is suggestive evidence that animal protein intake, especially from dairy products, has a stronger association with growth, particularly with weight gain, than plant protein. The evidence also suggests that a higher animal protein intake in childhood was associated with earlier onset of puberty (82, 83).

Pregnant and lactating women

During pregnancy, the average protein requirement is increased to provide additional protein for deposition in maternal (blood, uterus, and breasts), fetal, and placental tissues. Extra protein is also needed to maintain the increased mass of the pregnant body. According to the EFSA report (35), pregnant women should consume 1, 9, and 28 g of protein per day during their first, second, and third trimesters, respectively, in addition to the recommended protein intake for non-pregnant women. On the other hand, the German recommendations (84) for protein intake during pregnancy are comparatively lower than those of the EFSA, as they suggest an additional 10 g per day during the second and third trimesters.

However, increased protein intake during pregnancy – due to increased energy intake – should consist of regular food rather than high-protein supplements.

The average protein requirement is also increased during lactation when the breast milk produced by a woman provides all the protein her infant needs. The EFSA (35) recommends as a safe level of additional protein for lactating women an additional 19 g protein per day during the first 6 months of lactation (exclusive breastfeeding) and 13 g protein per day after 6 months (partial breastfeeding) above the recommended intake for non-lactating women. Therefore, a lactating and pregnant woman can, in most cases, cover the protein requirements with a regular diet if energy requirements are covered.

Older adults

As people grow older, their vulnerability to chronic illnesses increases. Nevertheless, it is essential to acknowledge that the health and functional capabilities of individuals aged 65 years and above vary and are not solely dependent on their age. Such conditions might lead to periodic temporary loss of body protein through chronic inflammations, catabolic exacerbations of the disease, brief periods of bed rest, and loss of appetite. The losses must be replaced from the diet, thus representing an added need for dietary protein (85). In addition, older individuals exhibit a gradual loss of muscle mass and strength with age. This is estimated to be a daily loss of 0.5 mg N per kg BW (86) that occurs naturally and is not simply due to decreased physical activity (87).

Frailty (88) and sarcopenia (89) indicate an increased risk of developing adverse health outcomes such as the

onset of disability, morbidity, institutionalization, or mortality (90). Physical function, or physical performance, is the clinically relevant outcome of muscle mass. An SR and meta-analysis of observational studies by Coelho-Júnior et al. (91) to investigate the association of relative protein intake and physical function in older adults showed that high protein intake (≥ 1.2 g/kg/day) and protein intake ≥ 1.0 g/kg per day among older adults resulted in better lower limb physical functioning and walking speed performance, respectively, in comparison to individuals who present relatively low protein (< 0.80 g/kg/day) intake. The authors concluded that these findings provided additional evidence for increasing dietary protein recommendations for community-dwelling older adults. Other studies on protein supplementation and resistance exercise in healthy older adults or with sarcopenia or frailty support these findings (91, 92). Pooled data (93) from European and North American community-dwelling older adults showed that higher daily protein intake might reduce physical function decline not only in older adults with protein intake below the current recommended dietary allowance of 0.8 g/kg BW per day, but also in those with a protein intake that is already considered sufficient, that is, above and beyond 0.8 g/kg BW per day. This dose-dependent association was observed for all levels of physical activity, suggesting that the results on the association between protein intake and physical function do not change by physical activity level. Other studies have shown an association between protein intake and lean muscle mass. However, other studies in healthy populations with protein supplements and physical training have not shown muscle augmentation (94, 95).

Results from prospective cohort studies suggested that a safe intake of up to 1.2–1.5 g protein/kg BW per day or approximately 15–20 E% represents an optimal intake level. Concerning the age-related decrease in energy intake, a diet with protein content in the range of 10–14 E% might not sufficiently cover the need for protein in absolute amounts. It is important to note that the older adult population is diverse, and individuals aged 65–75 years are generally considered healthy and able to maintain their energy levels. The recommendation from NNR2012 (96) for food planning was 18 E%, corresponding to about 1.2 g protein/kg BW per day. This recommendation aligns with several expert groups (94, 97) which have suggested a daily amount of 1.0–1.2 g/kg BW per day for healthy older adults. The ESPEN guidelines (98) recommend that protein intake be at least 1 g/kg BW per day and up to 2 g/kg BW per day for illness, injury, or malnutrition. However, an SR by the health council of the Netherlands concluded that increased protein intake has a possibly beneficial effect on lean body mass and, when combined with physical exercise, muscle strength; likely no effect on muscle strength when

not combined with physical exercise, or on physical performance and bone health; an ambiguous effect on serum lipids; and that too few RCTs were available to allow for conclusions on the other outcomes. This SR provides insufficient evidence that increasing protein in older adults with a protein intake ≥ 0.8 g/kg BW per day elicits health benefits (99).

Upper intake levels

The WHO (34) and NASEM (74) recommend that individuals without chronic kidney disease can safely consume higher-protein diets. The upper limit for protein intake in adults in NNR2012 was 20 E%. It was suggested to consume the following upper limits of protein, assuming adequate intake of other essential nutrients: 10 E% for 0–6 months, 15 E% for 6–11 months, 17 E% for 12–23 months, and 20 E% for 2 years and above.

Based on the risk of mortality and morbidity, the SR by Pedersen et al. (57) also assessed the evidence for a potential adverse effect of a high protein intake. There was no indication of adverse effects of protein intake concerning bone health provided a sufficient calcium intake, and an included meta-analysis did not find support for the acid hypothesis that a diet high in animal protein results in an increased systemic acid load and causes osteoporosis (100). One study with elderly subjects (101) found that animal protein intake was related to an increased risk of hypertension among persons ≥ 70 years of age. The lowest tertile of total protein intake was 14 E%, and the highest was 19 E%. An increase in GFR is a physiological adaptation to increased protein intake (102). Walrand et al. (103) found that a high protein intake did not increase GFR in the elderly participants in their study from a baseline GFR lower than that of the young participants. This was probably due to reduced kidney function in older adults. Individuals with mild-to-moderate chronic kidney disease also do not show the usual protein-induced increase in GFR (104). Caution is also required due to the observation of a decline in GFR among women with mild kidney insufficiency (105) and because older adults might have undiagnosed compromised kidney function due to severely reduced GFR. Regarding microalbuminuria, one experimental study found an increase in urinary albumin after seven days of a high protein intake of 2.4 g/kg BW per day. However, a similar increase in protein intake in another short-term experimental study of healthy young men did not find an increase in 24-h urinary albumin excretion (35). Further studies are needed to settle whether this discrepancy is due to the different durations of the studies or to various methods of analysis of albumin in the urine. Friedman (106) cites an earlier 3-week study showing a reduction in proteinuria with reduced protein intake (75–43 g per day). Caution is required until this matter is settled.

Protein and physical exercise

According to a position paper from the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine from 2016 (107), co-ingestion of protein with carbohydrates during recovery results in improved net protein balance post-exercise. Ingesting approximately 20–30 g total protein, or approximately 10 g of non-indispensable amino acids during exercise or the recovery period (post-exercise), seems to increase whole-body and muscle protein synthesis and improve N-balance. Current data suggest that dietary protein intake necessary to support metabolic adaptation, repair, remodeling, and protein turnover generally ranges from 1.2 up to 2.0 g/kg per day in cases of energy restriction or elevated protein intakes as high because of injury (102). Daily protein intake goals for the general population can be met with a meal plan providing a regular distribution of moderate amounts of high-quality protein across the day and following strenuous training sessions.

To match energy expenditure, adequate energy consumption, particularly from carbohydrates, is vital so amino acids are spared for protein synthesis and are not oxidized (108). According to Morton et al. (109), after reviewing 49 studies with 1,863 participants, it was found that dietary protein supplementation can greatly improve muscle strength and size changes during prolonged resistance exercise in healthy adults. However, consuming more than ~1.6 g/kg per day of protein through supplementation does not lead to additional gains in fat-free mass induced by resistance training.

A meta-analysis by Messina et al. (110) with 9 studies and 266 participants showed no difference between the effects of supplementing with soy protein and animal protein on gains in muscle mass and strength in response to resistance exercise, which is interesting in the transition toward plant-based protein sources, especially among young people.

Although animal protein is usually considered a more potent stimulator of muscle protein synthesis than plant protein, the effect of protein sources on lean mass and muscle strength must be systematically reviewed, considering more sustainable protein sources. One meta-analysis of 16 studies where total protein intakes were generally above the recommended dietary allowance at baseline and end of intervention showed that protein source did not affect changes in absolute lean mass or muscle strength. However, the authors concluded that animal protein tends to be more beneficial for lean mass than plant protein, especially in younger adults (<50 years old) (111).

References

1. Warensjö Lemming E, Pitsi T. The Nordic Nutrition Recommendations 2022 – food consumption and nutrient intake in the adult population of the Nordic and Baltic countries. *Food Nutr Res.* 2022; 66: e8572. doi: 10.29219/fnr.v66.8572
2. Samuelsson J, Rothenberg E, Lissner L, Eiben G, Zettergren A, Skoog I. Time trends in nutrient intake and dietary patterns among five birth cohorts of 70-year-olds examined 1971–2016: results from the Gothenburg H70 birth cohort studies, Sweden. *Nutr J.* 2019; 18(1): 66. doi: 10.1186/s12937-019-0493-8
3. Karlsson M, Becker W, Cederholm TE, Byberg L. A posteriori dietary patterns in 71-year-old Swedish men and the prevalence of sarcopenia 16 years later. *Br J Nutr.* 2021; 128(5): 1–12. doi: 10.1017/S0007114521003901
4. Wang R, Fratiglioni L, Liang Y, Welmer AK, Xu W, Mangialasche F, et al. Prevalence, pharmacological treatment, and control of cardiometabolic risk factors among older people in central Stockholm: a population-based study. *PLoS One.* 2015; 10(3): e0119582. doi: 10.1371/journal.pone.0119582
5. Castro-Mejia JL, Khakimov B, Krych L, Bülow J, Bechshøft RL, Højfeldt G, et al. Physical fitness in community-dwelling older adults is linked to dietary intake, gut microbiota, and metabolomic signatures. *Aging Cell.* 2020; 19(3): e13105. doi: 10.1111/ace1.13105
6. Helgadóttir H, Thorisdóttir B, Gunnarsdóttir I, Halldorsson TI, Palsson G, Thorsdóttir I. Lower intake of saturated fatty acids is associated with improved lipid profile in a 6-year-old nationally representative population. *Nutrients.* 2022; 14(3): e671. doi: 10.3390/nu14030671
7. World Health Organization, Food and Agriculture Organization of the United Nations. Nitrogen and protein content measurement and nitrogen to protein conversion factors for dairy and soy protein-based foods: a systematic review and modelling analysis. Geneva: World Health Organization; 2019.
8. Krul E. Calculation of nitrogen-to-protein conversion factors: a review with a focus on soy protein. *J Am Oil Chem Soc.* 2019; 96: 339–64. doi: 10.1002/aocs.12196
9. Reeds PJ. Dispensable and indispensable amino acids for humans. *J Nutr.* 2000; 130(7): 1835s–40s. doi: 10.1093/jn/130.7.1835S
10. Uauy R, Greene HL, Heird W. Conditionally essential nutrients: cysteine, taurine, tyrosine, arginine, glutamine, choline, inositol and nucleotides. Nutritional needs of the preterm infant. Pawling NY: Caduceus Medical Publishers Inc.; 1993.
11. Williams JZ, Abumrad N, Barbul A. Effect of a specialized amino acid mixture on human collagen deposition. *Ann Surg.* 2002; 236(3): 369–74; discussion 374–5. doi: 10.1097/0000658-200209000-00013
12. Hendrickson NR, et al. Conditionally essential amino acid supplementation reduces postoperative complications and muscle wasting after fracture fixation: a randomized controlled trial. *J Bone Joint Surg Am.* 2022; 104(9): 759–66. doi: 10.2106/JBJS.21.01014
13. Wang B, Wu G, Zhou Z, Dai Z, Sun Y, Ji Y, et al. Glutamine and intestinal barrier function. *Amino Acids.* 2015; 47(10): 2143–54. doi: 10.1007/s00726-014-1773-4
14. Blomhoff R, Andersen R, Arnesen EK, Christensen JJ, Eneroth H, Erkkola M, et al. Nordic Nutrition Recommendations 2023: integrating environmental aspects. Nord. Copenhagen: Nordisk Ministerråd; 2023, p. 388.
15. Christensen JJ, Arnesen EK, Andersen R, Eneroth H, Erkkola M, Høyer A, et al. The Nordic Nutrition Recommendations 2022 – principles and methodologies. *Food Nutr Res.* 2020; 64: e4402. doi: 10.29219/fnr.v64.4402
16. Stipanuk MH, Caudill MA. Biochemical, physiological, and molecular aspects of human nutrition. 4 ed. New York, NY: Elsevier Publication; 2019.
17. FAO Expert Consultation. Dietary protein quality evaluation in human nutrition. FAO Food Nutritional Paper. Auckland: FAO; 2013, p. 1–66.

18. Lee WT, Weisell R, Albert J, Tomé D, Kurpad AV, Uauy R. Research approaches and methods for evaluating the protein quality of human foods proposed by an FAO expert working group in 2014. *J Nutr.* 2016; 146(5): 929–32. doi: 10.3945/jn.115.222109
19. Bandyopadhyay S, Kashyap S, Calvez J, Devi S, Azzout-Marniche D, Tomé D, et al. Evaluation of protein quality in humans and insights on stable isotope approaches to measure digestibility – a review. *Adv Nutr.* 2022; 13(4): 1131–43. doi: 10.1093/advances/nmab134
20. Herreman L, Nommensen P, Pennings B, Laus MC. Comprehensive overview of the quality of plant- And animal-sourced proteins based on the digestible indispensable amino acid score. *Food Sci Nutr.* 2020; 8(10): 5379–91. doi: 10.1002/fsn3.1809
21. Sarwar Gilani G, Wu Xiao C, Cockell KA. Impact of antinutritional factors in food proteins on the digestibility of protein and the bioavailability of amino acids and on protein quality. *Br J Nutr.* 2012; 108 Suppl 2: S315–32. doi: 10.1017/S0007114512002371
22. Joye I. Protein digestibility of cereal products. *Foods.* 2019; 8(6): e199. doi: 10.3390/foods8060199
23. Päivärinta E, Iitonen ST, Pellinen T, Lehtovirta M, Erkkola M, Pajari AM. Replacing animal-based proteins with plant-based proteins changes the composition of a whole nordic diet – a randomised clinical trial in healthy finnish adults. *Nutrients.* 2020; 12(4): e943. doi: 10.3390/nu12040943
24. Pellinen T, Päivärinta E, Isotalo J, Lehtovirta M, Iitonen ST, Korkalo L, et al. Replacing dietary animal-source proteins with plant-source proteins changes dietary intake and status of vitamins and minerals in healthy adults: a 12-week randomized controlled trial. *Eur J Nutr.* 2022; 61(3): 1391–404.
25. Iitonen ST, Päivärinta E, Pellinen T, Viitakangas H, Risteli J, Erkkola M, et al. Partial replacement of animal proteins with plant proteins for 12 weeks accelerates bone turnover among healthy adults: a randomized clinical trial. *J Nutr.* 2021; 151(1): 11–19. doi: 10.1007/s00394-021-02729-3
26. Kaartinen NE, Tapanainen H, Maukonen M, Päivärinta E, Valsta LM, Iitonen ST, et al. Partial replacement of red and processed meat with legumes: a modelling study of the impact on nutrient intakes and nutrient adequacy on the population level. *Public Health Nutr.* 2022; 7: 1–12. doi: 10.1017/S1368980022002440
27. Ten Haaf DSM, van Dongen EJI, Nuijten MAH, Eijssvogels TMH, de Groot L, Hopman MTE. Protein intake and distribution in relation to physical functioning and quality of life in community-dwelling elderly people: acknowledging the role of physical activity. *Nutrients.* 2018; 10(4): nu10040506. doi: 10.3390/nu10040506
28. Loenneke JP, Loprinzi PD, Murphy CH, Phillips SM. Per meal dose and frequency of protein consumption is associated with lean mass and muscle performance. *Clin Nutr.* 2016; 35(6): 1506–11. doi: 10.1016/j.clnu.2016.04.002
29. Waterlow JC. Whole-body protein turnover in humans – past, present, and future. *Annu Rev Nutr.* 1995; 15: 57–92. doi: 10.1146/annurev.nu.15.070195.000421
30. Millward DJ. Nutrition and sarcopenia: evidence for an interaction. *Proc Nutr Soc.* 2012; 71(4): 566–75. doi: 10.1017/S0029665112000201
31. Eckart A, Struja T, Kutz A, Baumgartner A, Baumgartner T, Zurlfluh S, et al. Relationship of nutritional status, inflammation, and serum albumin levels during acute illness: a prospective study. *Am J Med.* 2020; 133(6): 713–22.e7. doi: 10.1016/j.amjmed.2019.10.031
32. Bharadwaj S, Ginoya S, Tandon P, Gohel TD, Guirguis J, Vallabh H, et al. Malnutrition: laboratory markers vs nutritional assessment. *Gastroenterol Rep* 2016; 4(4): 272–80. doi: 10.1093/gastro/gow013
33. Zhang Z, Pereira SL, Luo M, Matheson EM. Evaluation of blood biomarkers associated with risk of malnutrition in older adults: a systematic review and meta-analysis. *Nutrients.* 2017; 9(8): e829. doi: 10.3390/nu9080829
34. Joint WHO/FAO/UNU Expert Consultation. Protein and amino acid requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation. Geneva: World Health Organization; 2007.
35. Van Elswyk ME, Weatherford CA, McNeill SH. A systematic review of renal health in healthy individuals associated with protein intake above the US recommended daily allowance in randomized controlled trials and observational studies. *Adv Nutr.* 2018; 9(4): 404–18. doi: 10.1093/advances/nmy026
36. Van Elswyk ME, Weatherford CA, McNeill SH. Evaluation of protein requirements for trained strength athletes. *J Appl Physiol* (1985). 1992; 73(5): 1986–95. doi: 10.1152/jappl.1992.73.5.1986
37. Gunnarsdóttir S, Þorgeirsdóttir H, Torfadóttir JE, Steingrimsdóttir L, Tryggvadóttir EA, Geirsdóttir ÓG, et al. Könnun á mataræði Íslendinga 2019–2021 – Helstu niðurstöður og samanburður við könnun frá 2010–2011, in Könnun á mataræði Íslendinga. Reykjavík: Embætti landlæknis og Rannsóknastofa inæringarfræði við Háskóla Íslands; 2022.
38. Moon J, Koh G. Clinical evidence and mechanisms of high-protein diet-induced weight loss. *J Obes Metab Syndr.* 2020; 29(3): 166–73. doi: 10.7570/jomes20028
39. Schwingshackl L, Hoffmann G. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J.* 2013; 12: 48. doi: 10.1186/1475-2891-12-48
40. Rolland-Cachera MF, Deheeger M, Akrouit M, Bellisle F. Influence of macronutrients on adiposity development: a follow up study of nutrition and growth from 10 months to 8 years of age. *Int J Obes Relat Metab Disord.* 1995; 19(8): 573–8.
41. Madsen AL, Larnkjær A, Mølgaard C, Michaelsen KF. IGF-I and IGFBP-3 in healthy 9 month old infants from the SKOT cohort: breastfeeding, diet, and later obesity. *Growth Horm IGF Res.* 2011; 21(4): 199–204. doi: 10.1016/j.ghir.2011.05.003
42. Larnkjær A, Ingstrup HK, Schack-Nielsen L, Hoppe C, Mølgaard C, Skovgaard IM, et al. Early programming of the IGF-I axis: negative association between IGF-I in infancy and late adolescence in a 17-year longitudinal follow-up study of healthy subjects. *Growth Horm IGF Res.* 2009; 19(1): 82–6. doi: 10.1016/j.ghir.2008.06.003
43. Hörnell A, Lagström H, Lande B, Thorsdóttir I. Protein intake from 0 to 18 years of age and its relation to health: a systematic literature review for the 5th Nordic Nutrition Recommendations. *Food Nutr Res.* 2013; 57: e21083. doi: 10.3402/fnr.v57i0.21083
44. Tian S, Xu Q, Jiang R, Han T, Sun C, Na L. Dietary protein consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Nutrients.* 2017; 9(9): 982. doi: 10.3390/nu9090982
45. Virtanen HEK, Koskinen TT, Voutilainen S, Mursu J, Tuomainen TP, Kokko P, et al. Intake of different dietary proteins and risk of type 2 diabetes in men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Br J Nutr.* 2017; 117(6): 882–93. doi: 10.1017/S0007114517000745
46. Chen Z, Franco OH, Lamballais S, Ikram MA, Schoufour JD, Muka T, et al. Associations of specific dietary protein with longitudinal insulin resistance, prediabetes and type 2 diabetes: The

- Rotterdam Study. *Clin Nutr*. 2020; 39(1): 242–9. doi: 10.1016/j.clnu.2019.01.021
47. Laffel LM, Kanapka LG, Beck R, W, Bergamo K, Clements MA, Criego A, et al. Effect of continuous glucose monitoring on glycemic control in adolescents and young adults with type 1 diabetes: a randomized clinical trial. *JAMA*. 2020; 323(23): 2388–96. doi: 10.1001/jama.2020.6940
 48. Naghshi S, Sadeghi O, Willett WC, Esmailzadeh A. Dietary intake of total, animal, and plant proteins and risk of all cause, cardiovascular, and cancer mortality: systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ*. 2020; 370: m2412. doi: 10.1136/bmj.m2412
 49. Thorpe MP, Evans EM. Dietary protein and bone health: harmonizing conflicting theories. *Nutr Rev*. 2011; 69(4): 215–30. doi: 10.1111/j.1753-4887.2011.00379.x
 50. Lamberg-Allardt C, Bärebring L, Arnesen EK, Nwaru BI, Thorisdottir B, Ramel A, et al. Animal versus plant-based protein and risk of cardiovascular disease and type 2 diabetes: a systematic review of randomized controlled trials and prospective cohort studies. *Food Nutr Res*. 2023; 67: e9003. doi: 10.29219/fnr.v67.9003
 51. Sahni S, Cupples LA, McLean RR, Tucker KL, Broe KE, Kiel DP, et al. Protective effect of high protein and calcium intake on the risk of hip fracture in the Framingham offspring cohort. *J Bone Miner Res*. 2010; 25(12): 2770–6. doi: 10.1002/jbmr.194
 52. EFSA Panel on Dietetic Products, Nutrition and Allergies. Scientific opinion on dietary reference values for protein. *EFSA J*. 2012; 10(2): 2557. doi: 10.2903/j.efsa.2012.2557
 53. Dawson-Hughes B, Harris SS. Calcium intake influences the association of protein intake with rates of bone loss in elderly men and women. *Am J Clin Nutr*. 2002; 75(4): 773–9. doi: 10.1093/ajcn/75.4.773
 54. Pedersen AN, Kondrup J, Børsheim E. Health effects of protein intake in healthy adults: a systematic literature review. *Food Nutr Res*. 2013; 57: e21245. doi: 10.3402/fnr.v57i0.21245
 55. Munger RG, Cerhan JR, Chiu BC. Prospective study of dietary protein intake and risk of hip fracture in postmenopausal women. *Am J Clin Nutr*. 1999; 69(1): 147–52. doi: 10.1093/ajcn/69.1.147
 56. Devries MC, Sithamparapillai A, Brimble KS, Banfield L, Morton RW, Phillips SM. Changes in kidney function do not differ between healthy adults consuming higher- compared with lower- or normal-protein diets: a systematic review and meta-analysis. *J Nutr*. 2018; 148(11): 1760–75.
 57. Pedersen AN, Cederholm T. Health effects of protein intake in healthy elderly populations: a systematic literature review. *Food Nutr Res*. 2014; 11(58): e23364. doi: 10.3402/fnr.v58.23364
 58. Bruci A, Tuccinardi D, Tozzi R, Balena A, Santucci S, Frontani R, et al. Very low-calorie ketogenic diet: a safe and effective tool for weight loss in patients with obesity and mild kidney failure. *Nutrients*. 2020; 12(2): e333.
 59. Acharya P, Acharya C, Thongprayoon C, Hansrivijit P, Kanduri SR, Kovvuru K, et al. Incidence and characteristics of kidney stones in patients on ketogenic diet: a systematic review and meta-analysis. *Diseases*. 2021; 9(2): 39. doi: 10.3390/diseases9020039
 60. Lai R, Bian Z, Lin H, Ren J, Zhou H, Guo H. The association between dietary protein intake and colorectal cancer risk: a meta-analysis. *World J Surg Oncol*. 2017; 15(1): 169. doi: 10.1186/s12957-017-1241-1
 61. Liao LM, Loftfield E, Ettemadi A, Graubard BI, Sinha R. Substitution of dietary protein sources in relation to colorectal cancer risk in the NIH-AARP cohort study. *Cancer Causes Control*. 2019; 30(10): 1127–35. doi: 10.1007/s10552-019-01210-1
 62. Clinton SK, Giovannucci EL, Hursting SD. The World Cancer Research Fund/American Institute for Cancer Research Third Expert Report on diet, nutrition, physical activity, and cancer: impact and future directions. *J Nutr*. 2020; 150(4): 663–71. doi: 10.1093/jn/nxz268
 63. Kong F, Geng E, Ning J, Liu Z, Wang A, Zhang S, et al. The association between dietary protein intake and esophageal cancer risk: a meta-analysis. *Biosci Rep*. 2020; 40(1): BSR20193692.
 64. Mao Y, Tie Y, Du J. Association between dietary protein intake and prostate cancer risk: evidence from a meta-analysis. *World J Surg Oncol*. 2018; 16(1): 152. doi: 10.1186/s12957-018-1452-0
 65. Wu J, Zeng R, Huang J, Li X, Zhang J, Ho J, et al. Dietary protein sources and incidence of breast cancer: a dose-response meta-analysis of prospective studies. *Nutrients*. 2016; 8(11): 730. doi: 10.3390/nu8110730
 66. Song M, Fung TT, Hu FB, Willett WC, Longo VD, Chan AT, et al. Association of animal and plant protein intake with all-cause and cause-specific mortality. *JAMA Intern Med*. 2016; 176(10): 1453–63.
 67. Schwingshackl L, Schwedhelm C, Hoffmann G, Lampousi AM, Knüppel S, Iqbal K, et al. Food groups and risk of all-cause mortality: a systematic review and meta-analysis of prospective studies. *Am J Clin Nutr*. 2017; 105(6): 1462–73. doi: 10.3945/ajcn.117.153148
 68. Qi X-X, Shen P. Associations of dietary protein intake with all-cause, cardiovascular disease, and cancer mortality: a systematic review and meta-analysis of cohort studies. *Nutr Metab Cardiovasc Dis*. 2020; 30(7): 1094–105.
 69. Nachvak S, M, Moradi S, Anjom-Shoae J, Rahmani J, Nasiri M, Maleki V, et al. Soy, soy isoflavones, and protein intake in relation to mortality from all causes, cancers, and cardiovascular diseases: a systematic review and dose-response meta-analysis of prospective cohort studies. *J Acad Nutr Diet*. 2019; 119(9): 1483–500.e17. doi: 10.1016/j.jand.2019.04.011
 70. Huang J, Liao LM, Weinstein SJ, Sinha R, Graubard BI, Albanes D, et al. Association between plant and animal protein intake and overall and cause-specific mortality. *JAMA Intern Med*. 2020; 180(9): 1173–84. doi: 10.1001/jamainternmed.2020.2790
 71. Langsetmo L, Harrison S, Jonnalagadda S, Pereira SL, Shikany JM, Farsijani S, et al. Low protein intake irrespective of source is associated with higher mortality among older community-dwelling men. *J Nutr Health Aging*. 2020; 24(8): 900–5. doi: 10.1007/s12603-020-1422-4
 72. Chen Z, Glisic M, Song M, Aliahmad HA, Zhang X, Mounmdjian AC, et al. Dietary protein intake and all-cause and cause-specific mortality: results from the Rotterdam Study and a meta-analysis of prospective cohort studies. *Eur J Epidemiol*. 2020; 35(5): 411–29. doi: 10.1007/s10654-020-00607-6
 73. Wu G. *Amino acids: biochemistry and nutrition*. 2 ed. Boca Raton, FL: CRC Press; 2021, p. 816.
 74. Trumbo P, Schlicker S, Yates A, A, Poos M. Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein and amino acids. *J Am Diet Assoc*. 2002; 102(11): 1621–30. doi: 10.1016/S0002-8223(02)90346-9
 75. Rand WM, Pellett PL, Young VR. Meta-analysis of nitrogen balance studies for estimating protein requirements in healthy adults. *Am J Clin Nutr*. 2003; 77(1): 109–27. doi: 10.1093/ajcn/77.1.109
 76. Li SS, Blanco Mejia S, Lytvyn L, Stewart SE, Vigiouliouk E, Ha V, et al. Effect of plant protein on blood lipids: a systematic review and meta-analysis of randomized controlled trials. *J Am Heart Assoc*. 2017; 6(12): e006659.

77. Koletzko B, Demmelmair H, Grote V, Totzauer M. Optimized protein intakes in term infants support physiological growth and promote long-term health. *Semin Perinatol.* 2019; 43(7): 1511-53. doi: 10.1053/j.semperi.2019.06.001
78. Dietary protein quality evaluation in human nutrition. Report of an FAQ Expert Consultation. *FAO Food Nutr Pap.* 2013; 92: 1–66.
79. Koletzko B, von Kries R, Closa R, Escribano J, Scaglioni S, Giovannini M, et al. Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr.* 2009; 89(6): 1836–45. doi: 10.3945/ajcn.2008.27091
80. Patro-Gola B, Zalewski BM, Kouwenhoven SM, Karaś J, Koletzko B, Bernard van Goudoever J, et al. Protein concentration in milk formula, growth, and later risk of obesity: a systematic review. *J Nutr.* 2016; 146(3): 551–64. doi: 10.3945/jn.115.223651
81. Voortman T, Vitezova A, Bramer WM, Ars CL, Bautista PK, Buitrago-Lopez A, et al. Effects of protein intake on blood pressure, insulin sensitivity and blood lipids in children: a systematic review. *Br J Nutr.* 2015; 113(3): 383–402. doi: 10.1017/S0007114514003699
82. Günther AL, Karaolis-Danckert N, Kroke A, Remer T, Buyken AE. Dietary protein intake throughout childhood is associated with the timing of puberty. *J Nutr.* 2010; 140(3): 565–71. doi: 10.3945/jn.109.114934
83. Moslehi N, Asghari G, Mirmiran P, Azizi F. Longitudinal association of dietary sources of animal and plant protein throughout childhood with menarche. *BMC Pediatr.* 2021; 21(1): 206.
84. German Society for Nutrition. Protein [Internet]. German Society for Nutrition; 2017. Available from: webmaster@dge.de [cited 29 October 2023].
85. Nowson C, O'Connell S. Protein requirements and recommendations for older people: a review. *Nutrients.* 2015; 7(8): 6874–99. doi: 10.3390/nu7085311
86. Esmarck B, Anersen JL, Olsen S, Richter EA, Mizuno M, Kjaer M. Timing of postexercise protein intake is important for muscle hypertrophy with resistance training in elderly humans. *J Physiol.* 2001; 535(Pt 1): 301–11.
87. Hughes VA, et al. Longitudinal changes in body composition in older men and women: role of body weight change and physical activity. *Am J Clin Nutr.* 2002; 76(2): 473–81. doi: 10.1093/ajcn/76.2.473
88. Hogan DB. Chapter 3 – models, definitions, and criteria for frailty. In: Ram JL, Conn PM, eds. *Conn's handbook of models for human aging* (Second Edition). Oxford: Academic Press; 2018, pp. 35–44.
89. Bhasin S, Travison TG, Manini TM, Patel S, Pencina KM, Fielding RA, et al. Sarcopenia definition: the position statements of the sarcopenia definition and outcomes consortium. *J Am Geriatr Soc.* 2020; 68(7): 1410–8. doi: 10.1111/jgs.16372
90. Abellan van Kan G, Rolland Y, Bergman H, Morley JE, Kritchevsky SB, Vellas B. The I.A.N.A Task Force on frailty assessment of older people in clinical practice. *J Nutr Health Aging.* 2008; 12(1): 29–37. doi: 10.1007/BF02982161
91. Coelho-Júnior HJ, Milano-Teixeira L, Rodrigues B, Bacurau R, Marzetti E, Uchida M. Relative protein intake and physical function in older adults: a systematic review and meta-analysis of observational studies. *Nutrients.* 2018; 10(9): nu10091330. doi: 10.3390/nu10091330
92. Liao CD, Tsao JY, Wu YT, Cheng CP, Chen HC, Huang YC, et al. Effects of protein supplementation combined with resistance exercise on body composition and physical function in older adults: a systematic review and meta-analysis. *Am J Clin Nutr.* 2017; 106(4): 1078–91. doi: 10.3945/ajcn.116.143594
93. Mendonça N, Hengeveld LM, Visser M, Presse N, Canhão H, Simonsick EM, et al. Low protein intake, physical activity, and physical function in European and North American community-dwelling older adults: a pooled analysis of four longitudinal aging cohorts. *Am J Clin Nutr.* 2021; 114(1): 29–41. doi: 10.1093/ajcn/nqab051
94. Finger D, Goltz FR, Umpierre D, Meyer E, Rosa LH, Schneider CD. Effects of protein supplementation in older adults undergoing resistance training: a systematic review and meta-analysis. *Sports Med.* 2015; 45(2): 245–55. doi: 10.1007/s40279-014-0269-4
95. Xu ZR, Tan ZJ, Zhang Q, Gui QF, Yang YM. Clinical effectiveness of protein and amino acid supplementation on building muscle mass in elderly people: a meta-analysis. *PLoS One.* 2014; 9(9): e109141. doi: 10.1371/journal.pone.0109141
96. Nordic Nutrition Recommendations 2012: Integrating nutrition and physical activity. 5 ed. Nord. Copenhagen: Nordisk Ministerråd; 2014, p. 627.
97. Rizzoli R, Stevenson JC, Bauer JM, van Loon LJ, Walrand S, Kanis JA, et al. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: a consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas.* 2014; 79(1): 122–32. doi: 10.1016/j.maturitas.2014.07.005
98. Volkert D, Beck AM, Cederholm T, Cruz-Jentoft A, Goisser S, Hooper L, et al. ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr.* 2019; 38(1): 10–47.
99. Hengeveld LM, de Goede J, Afman LA, Bakker SJL, Beulens JWJ, Blaak EE, et al. Health effects of increasing protein intake above the current population reference intake in older adults: a systematic review of the Health Council of the Netherlands. *Adv Nutr.* 2022; 13(4): 1083–117. doi: 10.1093/advances/nmab140
100. Fenton TR, Tough SC, Lyon AW, Eliasziw M, Hanley DA. Causal assessment of dietary acid load and bone disease: a systematic review & meta-analysis applying Hill's epidemiologic criteria for causality. *Nutr J.* 2011; 10: 41. doi: 10.1186/1475-2891-10-41
101. Altorf-van der Kuil W, Engberink MF, Brink EJ, van Baak MA, Bakker SJ, Navis G, et al. Dietary protein and blood pressure: a systematic review. *PLoS One.* 2010; 5(8): e12102. doi: 10.1371/journal.pone.0012102
102. Bankir L, Bouby N, Trinh-Trang-Tan MM, Ahloulouy M, Promeneur D. Direct and indirect cost of urea excretion. *Kidney Int.* 1996; 49(6): 1598–607. doi: 10.1038/ki.1996.232
103. Walrand S, Short KR, Bigelow ML, Sweatt AJ, Hutson SM, Nair KS. Functional impact of high protein intake on healthy elderly people. *Am J Physiol Endocrinol Metab.* 2008; 295(4): E921–8. doi: 10.1152/ajpendo.90536.2008
104. Bosch JP, Lew S, Glabman S, Lauer A. Renal hemodynamic changes in humans. Response to protein loading in normal and diseased kidneys. *Am J Med.* 1986; 81(5): 809–15.
105. Knight EL, Stampfer MJ, Hankinson SE, Spiegelman D, Curhan GC. The impact of protein intake on renal function decline in women with normal renal function or mild renal insufficiency. *Ann Intern Med.* 2003; 138(6): 460–7. doi: 10.7326/0003-4819-138-6-200303180-00009
106. Friedman AN. High-protein diets: potential effects on the kidney in renal health and disease. *Am J Kidney Dis.* 2004; 44(6): 950–62. doi: 10.1053/j.ajkd.2004.08.020
107. Thomas DT, Erdman KA, Burke LM. Position of the academy of nutrition and dietetics, dietitians of Canada, and the American college of sports medicine: nutrition and athletic

- performance. *J Acad Nutr Diet*. 2016; 116(3): 501–28. doi: 10.1016/j.jand.2015.12.006
108. Rodriguez NR, Vislocky LM, Gaine PC. Dietary protein, endurance exercise, and human skeletal-muscle protein turnover. *Curr Opin Clin Nutr Metab Care*. 2007; 10(1): 40–5. doi: 10.1097/MCO.0b013e3280115e3b
 109. Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med*. 2018; 52(6): 376–84. doi: 10.1136/bjsports-2017-097608
 110. Messina M, Lynch H, Dickinson JM, Reed KE. No difference between the effects of supplementing with soy protein versus animal protein on gains in muscle mass and strength in response to resistance exercise. *Int J Sport Nutr Exerc Metab*. 2018; 28(6): 674–85. doi: 10.1123/ijsnem.2018-0071
 111. Lim MT, Pan BJ, Toh DWK, Sutanto CN, Kim JE. Animal protein versus plant protein in supporting lean mass and muscle strength: a systematic review and meta-analysis of randomized controlled trials. *Nutrients*. 2021; 13(2): nu13020661. doi: 10.3390/nu13020661

***Ólöf Guðný Geirsdóttir**

Faculty of Food Science and Nutrition
 School of Health Science
 University of Iceland
 102 Reykjavík
 Iceland
 Email: ogg@hi.is