



UNIVERSITY OF HELSINKI



<https://helda.helsinki.fi>

Helda

---

## The Potent Phytoestrogen 8-Prenylnaringenin: A Friend or a Foe?

Pohjanvirta, Raimo

Multidisciplinary Digital Publishing Institute

2022-03-15

---

Pohjanvirta, R.; Nasri, A. The Potent Phytoestrogen 8-Prenylnaringenin: A Friend or a Foe? *Int. J. Mol. Sci.* 2022, 23, 3168.

---

<http://hdl.handle.net/10138/349262>

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



Review

# The Potent Phytoestrogen 8-Prenylnaringenin: A Friend or a Foe?

Raimo Pohjanvirta <sup>1,\*</sup> and Atefeh Nasri <sup>1,2</sup>

<sup>1</sup> Department of Food Hygiene and Environmental Health, University of Helsinki, 00790 Helsinki, Finland

<sup>2</sup> Department of Veterinary Biomedical Science, Western College of Veterinary Medicine, Saskatoon, SK S7N 5B4, Canada; nasriatefeh@gmail.com

\* Correspondence: raimo.pohjanvirta@helsinki.fi

**Abstract:** 8-prenylnaringenin (8-PN) is a prenylated flavonoid, occurring, in particular, in hop, but also in other plants. It has proven to be one of the most potent phytoestrogens in vitro known to date, and in the past 20 years, research has unveiled new effects triggered by it in biological systems. These findings have aroused the hopes, expectations, and enthusiasm of a “wonder-drug” for a host of human diseases. However, the majority of 8-PN effects require such high concentrations that they cannot be reached by normal dietary exposure, only pharmacologically; thus, adverse impacts may also emerge. Here, we provide a comprehensive and up-to-date review on this fascinating compound, with special reference to the range of beneficial and untoward health consequences that may ensue from exposure to it.

**Keywords:** phytoestrogens; natural compounds; xanthohumol; isoxanthohumol; naringenin; flavonoids; beer



**Citation:** Pohjanvirta, R.; Nasri, A. The Potent Phytoestrogen 8-Prenylnaringenin: A Friend or a Foe? *Int. J. Mol. Sci.* **2022**, *23*, 3168. <https://doi.org/10.3390/ijms23063168>

Academic Editors: Nicolas Chevalier and Charlotte Hinault-Boyer

Received: 31 January 2022

Accepted: 13 March 2022

Published: 15 March 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).


## 1. Introduction

Hop (*Humulus lupulus* L.) has been used as a preservative and flavoring agent in beer for centuries. This plant is rich in phenolic acids, flavonoids, proanthocyanidins, prenylated chalcones, and flavanones, as well as catechins with potential therapeutic applications [1]. Historically, menstrual disturbances were commonly reported among female hop pickers [2], and hop extract was reported to be efficient in reducing menopause-associated hot flashes in women. Moreover, hop baths have been used for treatment of gynecological disorders over the years [3]. Therefore, a recurring suggestion was that hop may contain compounds with powerful estrogenic activities. The estrogenicity of hops was attributed to xanthohumol (XN) without any reported scientific data [2]. Preliminary studies on estrogenic potency of hop showed contradictory results: some studies revealed high estrogenic activity [4,5], while others found no or low estrogenic activity of hop [6]. These discrepancies might have been due to the varieties of extracts and the nature of assays used [5]. In 1999, Milligan et al. reported of a novel compound, 8-prenylnaringenin (8-PN), after a successful bioassay-guided fractionation of hop extracts, and provided the first evidence of its prominent estrogenic activity [7]. In fact, this compound had been originally identified from hops as early as 1984 [8] but had been forgotten thereafter. Since Milligan et al.'s discovery, numerous studies have substantiated the high estrogenic potency of 8-PN. With accumulating evidence on the wide exposure of 8-PN to the general population, and of its exceptionally high potency in vitro, concerns have also arisen regarding the possible adverse effects to humans. This review therefore presents a comprehensive and up-to-date view on 8-PN.

## 2. Occurrence, Sources of, and Exposure to 8-PN

Throughout the years, a wide variety of methods, predominantly mass spectrometry-based, have been developed for reliable quantitative determination of 8-PN and related

prenylated flavonoids (Table 1). They have enabled the analysis of these substances in diverse matrices and revealed that the main source of human 8-PN exposure was beer drinking when hop was used during the brewing process [9]. Hops are added to beer in the form of dried hop or lipophilic extracts [10]. Hop extracts or dried hops are obtained from female hop cones that are particularly rich in prenylated chalcones (XN, desmethyl-XN) and prenylated flavanones (6-PN, 8-PN) [11]. The exact amount of each ingredient varies, because chalcones undergo isomerization during the brewing process, which results in the conversion of XN to isoxanthohumol (IX) and of desmethyl-XN to a racemic mixture 6- and 8-PNs (Figure 1). Moreover, the cyclization of XN to IX may generate two enantiomers of IX, which can lead to two enantiomers of 8-PN [12]. The concentration of XN may amount to 1% while 8-PN is more than 10 times less abundant [13].



**Figure 1.** Structural formulae of prenylated flavonoids (with the most important compounds for this review in bold). XN can isomerize to IX, and desmethyl-XN to a racemic mixture of 6-PN and 8-PN. IX, in turn, may be metabolized to a variable degree to 8-PN (see text for details). The asterisk indicates the chiral center in 8-PN.

The volume of beer consumed per capita varies substantially among countries, being highest in the Czech Republic (191.8 L in 2018) [14]. The concentration of 8-PN in beer also varies. For example, in a study reported by Stevens and Page (2004) [9], European lager was found to contain a mere 1 µg/L, whereas American porter possessed a concentration of 240 µg/L. Still, in people harboring high intestinal and/or hepatic biotransformation capacities (see 8-PN pharmacokinetics), the ultimate source of 8-PN is, in most cases, IX, because its concentration in beer exceeds that of 8-PN by 10–40-fold [9,15,16]. Assuming the consumption of 1 L of, for example, “imported stout” beer with 8-PN and IX concentrations of 69 and 2100 µg/L, respectively [15], a person endowed with efficient conversion capability (80%; [17]) would be exposed to ~1700 µg 8-PN. Possemiers et al. (2006) have

also estimated that moderate beer consumption could lead to 8-PN exposure in the range of 1–2 mg/day [17].

In an opened beer bottle, IX concentration may decrease by almost 20% in a day and over 30% in a week. For 8-PN, these reductions can be even larger (up to 50% in a week) [18]. No differences in 8-PN, IX, or XN concentrations between organically and conventionally produced hops and beer were detected [19].

Apart from hops, the occurrence of 8-PN has been reported in at least 21 ethnomedicinally applied plants [20], including citrus plants [21]. It may also occur in herb or hop teas at low concentrations (<10 µg/L); IX concentrations seem to be only slightly higher in these products (<15 µg/L) [15,16].

Clarke and coworkers [22] identified and quantified phytoestrogens in 35 dietary supplements on sale in the UK, Canada, and Italy. None of the samples contained measurable levels of 8-PN, 6-PN, or 6,8-diprenylnaringenin. In a more recent study, Schretter et al. (2020) [23] were unable to detect 8-PN in two out of the three hop dietary supplements analyzed. In contrast, the third product would provide 144 µg 8-PN and 1740 µg IX at the recommended daily dosage, indicating huge differences among these products. Coldham and Sauer (2001) reported that two dietary supplements intended for breast enhancement contained ~10 µg/g 8-PN. This would result in a calculated daily intake of 103–155 µg [24].

In the study by Schretter et al. (2009) [23], a herbal medicinal product and two batches of the herbal drug “Lupuli flowers” (in pharmacopeia quality) proved to encompass similar quantities of IX and 8-PN, and the recommended daily dose for the medicinal product would lead to the intake of ~45 µg 8-PN and ~255 µg IX. Dhoooghe et al. (2010) studied hop-derived capsules, tablets, and drops on the market in Belgium for their 8-PN content [25]. In two products, 8-PN levels were undetectable. At their recommended daily doses, the others would release from 40 to 150 µg 8-PN. Thus, 8-PN intake from hop dietary supplements or medicinal products can mostly amount to a level comparable to the exposure of beer-lovers with fairly poor IX conversion capability.

### 3. Methods Applied in 8-PN Identification and Quantification

The bulk of the methods described to date for 8-PN quantification have combined LC or HPLC separation with detection by mass spectrometry (MS), although a gas chromatography (GC)–MS method has also been published [15,19,20,26–29] (Table 1). The varieties of MS used include electron impact (EI), atmospheric pressure electrospray ionization (ESI), or atmospheric pressure chemical ionization (APCI) in either positive or negative ion modes [15,18,24,30]. Apart from MS, HPLC methods with ultraviolet (UV) or UV/diode array detector (DAD) detection have been developed [18,25,31,32]. Some modifications of these techniques include ultra-high-pressure LC–tandem MS (UHPLC–MS/MS), ultra-high-performance supercritical fluid chromatography (UHPSFC; with either UV or MS detection), and the application of secondary standards in HPLC–UV/DAD [23,25]. Antibody-based methods have also found their way to 8-PN quantification. Polyclonal antibodies were used for the design of a radioimmunoassay (RIA), and monoclonal for devising an enzyme-linked immunosorbent assay (ELISA) [33,34]. The most recent new technique described for 8-PN quantification is stable isotope dilution analysis (SIDA) [16].

In most cases, these quantitative methods have measured racemic 8-PN, but there are also a couple of enantiospecific analytic techniques among them [20,32]. Moreover, as a facile means to determine the enantiomeric purity of 8-PN, a method was devised in which diastereomeric (1*S*)-(–)-camphanic acid esters of the enantiomers are first produced to be subsequently separated by reversed-phase HPLC [35].

**Table 1.** Methods used in quantitative 8-PN analysis.

Method <sup>1</sup>	Matrix	Sensitivity		Precision		Accuracy	Related Analytes Measured	Reference
		LLOQ <sup>2</sup> (ng/mL)	LOD <sup>3</sup> (ng/mL)	CV <sup>4</sup> -Intra (%)	CV-Inter (%)	RE <sup>5</sup> (%)		
GC/MS-SIM	Beer, hop pellets	NP	5 (beer)	NP	NP	≤65 (beer)	–	[26]
LC-ESI-MS	Beer	2.4	0.8	2.0	8.0	8.8	–	[27]
LC-ESI-MS	Serum, urine <sup>6</sup>	S: 50; U: 10	NP	S: ≤10.9; U: ≤14.9	S: ≤13.7; U: ≤14.1	S: 2.6; U: 2.2	–	[20]
LC-ESI-MS/MS	Urine, beer	5 (urine)	0.03 (urine)	≤13.9 (urine)	≤12.6 (urine)	≤14.6 (urine)	X, IX	[29]
HPLC-MS/MS	Beer, hop extracts, herb teas	NP	NP	8.8 (beer)	8.2 (beer)	≤10 (beer)	XN, IX, 6-PN	[15]
UHPLC-MS/MS	Serum	1	NP	≤10.5	≤12.1	≤4	XN, IX, 6-PN	[28]
UHPLC-MS/MS	Beer, hop pellets	NP	NP	≤6	≤5	≤13	XN, IX	[19]
HPLC-APCI-MS	Dietary supplement for breast enhancement	NP	NP	6	2	3	XN, IX, 6-PN, 6,8-diPN	[24]
HPLC-APCI-MS	Serum, urine	S: 4.8; U: 1.2	S: 1.5; U: 0.4	S: 3.9; U: 6.0	S: 9.7; U: 14.4	S: 7.4; U: 10.1	XN, IX	[30]
HPLC-APCI-MS/MS	Beer, hop extracts, herb teas	NP	NP	≤7.9	≤8.2 (beer)	≤10	XN, IX, 6-PN	[15]
HPLC-APCI-MS	Beer, hop extracts	20	6	NP	NP	NP	XN, IX	[18]
HPLC-UV	Beer, hop extracts	100	30	NP	NP	NP	XN, IX	[18]
HPLC-UV	Beer, hops, hop pellets	30	10	<0.3	≤2.0	≤4.8	XN, IX	[32]
HPLC-UV/DAD	Hop cultivars and genotypes	3800	1000	NP	NP	3.2	XN, IX, 6-PN	[31]
HPLC-UV/DAD (+SS)	Hop extract and capsules	NP	860	<4.0	≤5.0	≤8.1	XN, IX, 6-PN	[25]
UHPSFC-UV	Hop dietary supplements, herbal products	100	60	0.0 <sup>7</sup>	0.1 <sup>7</sup>	≤7.7	XN, IX, 6-PN	[23]
UHPSFC-MS	Hop dietary supplements, herbal products	50	20	0.02 <sup>7</sup>	0.1 <sup>7</sup>	NP	XN, IX, 6-PN	[23]
RIA (polyclonal) <sup>8</sup>	Beer, urine	NP	0.3 (urine)	<9 (urine)	<27 (urine)	≤36 (urine)	–	[33]
ELISA (monoclonal) <sup>9</sup>	Serum, urine	17.1	4.4	S: 2.4; U: 0.7	S: 6.1; U: 7.2	S: 5.6; U: 5.1	X, IX	[34]
SIDA-LC-MS/MS	Beer, hop pellets, and tea	1.3 (beer)	0.32 (beer)	2.09 (beer)	8.2 (beer)	≤13 (beer)	XN, IX, 6-PN	[16]

Table 1. Cont.

Method <sup>1</sup>	Matrix	Sensitivity		Precision		Accuracy	Related Analytes Measured	Reference
		LLOQ <sup>2</sup> (ng/mL)	LOD <sup>3</sup> (ng/mL)	CV <sup>4</sup> -Intra (%)	CV-Inter (%)	RE <sup>5</sup> (%)		
<sup>1</sup> Method abbreviations:	APCI, atmospheric pressure chemical ionization; DAD, diode-array detection; ELISA, enzyme-linked immunosorbent assay; ESI, electrospray ionization; GC, gas chromatography; HPLC, high-performance liquid chromatography; LC, liquid chromatography; MS, mass spectrometry; RIA, radioimmunoassay; SIDA, stable isotope dilution analysis; UHPSFC, ultrahigh-performance supercritical fluid chromatography; UV, ultraviolet light							
<sup>2</sup> Lower limit of quantification								
<sup>3</sup> Limit of detection								
<sup>4</sup> Coefficient of variation								
<sup>5</sup> Relative error								
<sup>6</sup> For both enantiomers								
<sup>7</sup> Based on peak area								
<sup>8</sup> Both enantiomers detected; cross-reactivity with XN, IX and 6-PN $\leq$ 0.15%								
<sup>9</sup> Both enantiomers detected; cross-reactivity with XN and IX < 0.01%, with 6-PN < 0.03%								

#### 4. Pharmacokinetics of 8-PN

Studies with pure 8-PN on other aspects of its pharmacokinetics than biotransformation are scarce. After the ingestion of a single oral dose of 500 mg racemic 8-PN in 16 healthy young adults (8 women, 8 men), a maximum plasma concentration was attained at 1.6 h with large inter-individual (but not gender-related) differences in the height of the peak [36]. Interestingly, at an equimolar dose, 6-PN peaked later (2.3 h) and its area under the plasma concentration–time curve remained much smaller. When three groups of eight healthy postmenopausal women were treated orally with 50, 250, or 750 mg racemic 8-PN, the compound was rapidly absorbed, leading to maximum serum drug concentrations 1.0–1.5 h after administration [37]. Thereafter, drug serum concentrations decreased sharply, followed by a further increase in concentrations leading to a second peak, which occurred at 7–10 h. This was suggestive of marked enterohepatic recirculation, probably attributable to the prenyl group in 8-PN [36].

In agreement with these *in vivo* reports attesting to effective gastrointestinal (GI) absorption of 8-PN, in monolayers of the human intestinal epithelial cancer cell line, (Caco-2), 8-PN also exhibited good absorption, probably via passive diffusion [38]. In these cells, 8-PN was biotransformed into two glucuronides (predominantly 4'-O-glucuronide) and two sulfates, with metabolism reaching 57% by 4 h. Human liver microsomes generated a total of 12 metabolites of 8-PN, and biotransformation occurred on the prenyl group and the flavanone skeleton [39]. Two of the most abundant of the metabolites, (E)-8-(4''-hydroxy isopentenyl)naringenin (8-PN-OH) and (E)-8-(4''-oxoisopentenyl)naringenin (8-PN=O), exhibited estrogenic activity *in vitro*. Cytochromes P450 (CYPs), including CYP2C8 and CYP2C19, catalyzed the phase I reaction of alcohol formation in the prenyl side chain of 8-PN [19]. The alcohol derivatives of 8-PN produced in hepatocytes were subsequently conjugated to at least six glucuronides but apparently no sulfates. In contrast to the situation in enterocytes, 7-O-glucuronide isomer was found to be the most abundant conjugate (~80% of all). The enzymes responsible for glucuronidation of 8-PN were uridine 5'-diphosphoglucuronosyltransferase (UGT)1A1, UGT1A6, UGT1A8, and UGT1A9. Only 4% of 8-PN remained unmetabolized after a 4-h incubation with human hepatocytes [38].

A more recent study by Fang et al. (2019) [40] examined whether 8-PN biotransformation in human liver microsomes or in reactions catalyzed by 11 recombinant human UGTs was enantiomer-specific. Incubation of racemic 8-PN with human liver microsomes resulted in the formation of (2R)-8-PN-7-O-glucuronide and (2S)-8-PN-7-O-glucuronide in a ratio of 1.37:1.00. Except for UGT1A1, the UGTs produced more (2R)-8-PN-O-glucuronides than the corresponding (2S)-glucuronides from purified (2R)-8-PN and (2S)-8-PN. Therefore, the half-life of (2S)-8-PN is likely to be longer in humans.

8-PN may also be metabolized in the human GI tract, because the anaerobic intestinal bacterium *Eubacterium ramulus* isomerized and hydrogenated 8-PN into two chalcones, O-desmethyl-XN and O-desmethyl- $\alpha,\beta$ -dihydroxanthohumol [41]. In addition to this bacterium, numerous fungi have been shown to be able to transform 8-PN [42]. Besides an oxygenated metabolite, which was also reported as a minor derivative of 8-PN in human liver microsomes [39], glucosylated, acyl-glucosylated, and sulfated fungi-generated metabolites have been identified [42–45].

Human liver microsomes (but not Aroclor-induced rat liver microsomes [46]) are capable of converting IX to 8-PN via O-demethylation [47], catalyzed by CYP1A2 [48]. Since this reaction can also occur in the GI tract and, e.g., beer usually contains 30–40 times more IX than 8-PN [30], the exposure to 8-PN is critically dependent on IX abundance and metabolism. *In vitro* studies implied that in the human GI tract, the conversion from IX to 8-PN almost exclusively occurs in distal colon [17]. Fecal samples from female volunteers revealed that the fecal microbiota could be separated into high (8/51), moderate (11/51), and slow (32/51) IX converters, with a mean 8-PN production of 78.8, 48.5, and 6.9%, respectively [2]. Classification of humans based on urinary excretion of 8-PN yielded similar ratios [49]. The microbial bioactivation of IX was inhibited by coexposure to two other proestrogens, daidzein and secoisolariciresinol, both *in vitro* and *in vivo* (by 20.5%

and 35.4%, respectively) [50]. The intestinal bacterium responsible for the O-demethylation of IX to 8-PN was identified to be *Eubacterium limosum* [51]. After strain selection of the bacterium, a conversion efficiency of 90% was achieved [51,52].

Upon ingestion of pure 8-PN or 6-PN by young adults, the cumulative urinary excretion over 24 h for both compounds was less than 1.5% of the administered dose [36]. In the study by Rad et al. (2006) [37], described above, 4.8–8.3% of dose was excreted by 48 h in urine, almost entirely as 8-PN conjugates. In feces, slightly less than 25% of the dose administered was recovered as free 8-PN by 48 h but it may be an underestimation due to technical problems encountered. It was unclear to what extent this free 8-PN stemmed from de-conjugated metabolites, as conjugates of 8-PN seem to be hydrolyzed by human intestinal microbiota [53]. In support of these findings, in menopausal women ingesting standardized extracts of spent hops the elimination half-life for 8-PN was estimated to be over 20 h [54]. Hence, excretion of 8-PN appears to be a relatively slow process, mainly due to enterohepatic recirculation of conjugated and de-conjugated metabolites. This suggests that although oral 8-PN undergoes extensive first-pass metabolism in the GI tract and liver, it can reach effective biological concentrations due to its low elimination rate, with a possibility of accumulation in target tissues. Therefore, more studies with longer follow-up times are needed to ascertain these kinetic aspects.

Simultaneous exposure to 8-PN may influence the kinetics of pharmaceuticals, because 8-PN proved to be a potent inhibitor of the ABCG2 efflux transporter [55]. Likewise, it was reported to inhibit two other ABC transporters, P-glycoprotein (ABCB1/P-gp) and multidrug resistance-associated protein 1 (ABCC1/MRP1) [56]. Thus, 8-PN possesses the potential to interfere with the efflux of concomitantly consumed drug substrates for these transporters.

Finally, all of the human studies have examined racemic 8-PN. In a preliminary experiment in rats, the *S*-form of 8-PN appeared to be excreted in urine at a slightly faster rate than *R*-8-PN after iv-injection of racemic 8-PN [20]. As mentioned above, indirect evidence suggests that in humans the converse may be true. On the other hand, it should be noted that back conversion to a racemic mixture might occur from both enantiomers [57].

## 5. Estrogenic Activity of 8-PN

### 5.1. In Vitro Studies

8-PN is one of the most potent phytoestrogens currently known. This has been confirmed by a variety of in vitro assays for estrogenic activity (Table 2), including yeast-based screens expressing the human estrogen receptor (ER) and a reporter gene (usually luciferase,  $\beta$ -galactosidase or chloramphenicol acetyltransferase) under the control of estrogen-responsive sequences (ERE) [5,46,58,59], human cell lines responsive to estrogenic stimulation [5,7,60,61], and ER binding [5,7,61,62]. Compared with established phytoestrogens, such as coumestrol, genistein, and daidzein, the estrogenic activity of 8-PN has generally proven to be 0.6–150, 8–150, and 50–1500-fold stronger, respectively [5,7,59,62] (but see ER $\beta$ -specific data below), while concomitantly 5–250 times weaker than that of 17 $\beta$ -estradiol (E2) [5,7,46,59–64]. The estrogenic activities of IX and 6-PN have been reported to be ~1% and <1% relative to 8-PN, respectively, while XN appears to be devoid of estrogenic properties [7]. Likewise, the parent compound of 8-PN, naringenin, shows only very weak estrogenicity [65,66].

The studies cited above have primarily measured ER $\alpha$ -mediated responses. Initial studies suggested that the two enantiomers of 8-PN exhibit similar estrogenic activities in vitro and show no preferential affinity to the two forms of ERs—ER $\alpha$  and ER $\beta$  [5]. However, this view was later contested by Schaefer et al. (2003) [63], who reported that 2*S*(–)8-PN shows moderately higher ER affinity and estrogenic activity in vitro and in vivo than 2*R*(+) 8-PN. Furthermore, in their hands the affinity of 8-PN for ER $\alpha$  was >2-fold higher than for ER $\beta$  measured by in vitro competitive binding assay and the estrogenic activity at ER $\alpha$  was >3.6-fold higher than at ER $\beta$  in a transactivation analysis. In support of this, Overk et al. (2005) reported a 3-fold higher IC<sub>50</sub> for ER $\beta$  vs. ER $\alpha$  binding [61],

and Helle et al. (2014) recorded a 3-fold greater relative affinity to ER $\alpha$  vs. ER $\beta$  for 8-PN relative to E2 [66]. On the other hand, while using yeast cells expressing human ERs, Bovee et al. (2004) found quite the opposite: the affinity of 8-PN for ER $\beta$  was 3 times as high as for ER $\alpha$  [67]. In any case, even the latter study confirmed that relative to E2, 8-PN shows higher potency at ER $\alpha$  vs. ER $\beta$ , and in this regard appears to conspicuously stand out from the bulk of other phytoestrogens, such as coumestrol, genistein, and daidzein, which are all clearly more potent at ER $\beta$  [67]. In fact, genistein and daidzein may bind to ER $\beta$  with an affinity 85 or 1,4 times as great as that of 8-PN, respectively [61]. They have also been reported to activate ER $\beta$  at lower concentrations than 8-PN [68].

In a battery of tests measuring estrogenicity in estrogen-sensitive MCF-7 cells, the maximal responses generated by 8-PN were similar to those generated by E2, implying that it is a full estrogen agonist. Moreover, no evidence was found for any estrogen antagonist action by 8-PN at concentrations of up to  $10^{-6}$  M [62]. Likewise, results from a yeast-based estrogen bioassay indicated a purely agonistic action at both ER subtypes [69]. In silico modeling of ER binding further suggested that chain prenylation without an increase in molecular length, which enables 8-PN to fit into the hydrophobic pocket in the human ER, is responsible for the high agonist activity of 8-PN compared with naringenin [69]. However, conflicting data were obtained in rat N46A-B14 cells derived from raphe nuclei region of the brain (thus serotonergic in origin) and stably expressing a transgenic human ER $\beta$  along with an ERE-luciferase construct [68]. In this test system, co-treatment of 8-PN with E2 resulted in a slight but statistically significant inhibition of E2 impact, suggesting partial agonism as the mode of interaction for 8-PN with ER $\beta$ . The same research team used these rat cells also in a subsequent study aimed at comparing other phytoestrogens with E2 for their influence on the expression of selected estrogen-responsive genes [70]. In proliferative conditions, 8-PN at a 1000-fold concentration faithfully reproduced the pattern of E2-induced effects; in differentiating conditions, the responsiveness of the analyzed genes towards E2 and 8-PN treatments was almost completely lost.

In a yeast assay designed to reveal agonistic and antagonistic properties of substances on human ER $\alpha$  and androgen receptor, 8-PN again proved to be a pure ER $\alpha$  agonist. It had no activity on the androgen receptor [71]. However, this latter outcome was contested by another group also using a yeast-based androgen receptor assay: here, 8-PN demonstrated anti-androgenic activities. Yet, when assessed in the same study by prostate specific antigen production in androgen receptor positive PC3(AR) $_2$  cells, 8-PN failed to show any anti-androgenicity [72]. A subsequent study revealed the critical role of the prenyl group at C(8) for 8-PN ER subtype and action selectivity [73]. When the prenyl group was substituted with alkyl chains of varying lengths and branching patterns, the new 8-alkylnaringenins were found to span an activity spectrum ranging from full agonism all the way via partial agonism to antagonism. An intriguing example is 8-(2,2-dimethylpropyl)naringenin, which was a potent agonist of ER $\alpha$ , but pronounced antagonist of ER $\beta$ . Overall, the antagonist effect of the derivatives was more substantial on ER $\beta$  than ER $\alpha$  [73]. Taken together, the weight of evidence points to a solely agonist action for 8-PN at ER $\alpha$ . At ER $\beta$  or the androgen receptor, the issue is not equally clear and awaits further studies.

In addition to its role as a ligand-activated transcription factor, ER can also mediate rapid nongenomic actions, such as activation of various protein-kinase cascades. These are believed to involve membrane-associated ERs [74]. Intriguingly, while both E2 and 8-PN induced the association of ER $\alpha$  with c-Src tyrosine kinase in MCF-7 cells, only E2 did the same with PI3 kinase. This failure of 8-PN resulted in a rapid decline in cyclin D1 and E steady state protein levels and induction of apoptosis [75]. It should be noted that induction of these non-genomic ER actions required high, pharmacological concentrations of 8-PN ( $\geq 10$   $\mu$ M), but still the difference in the bioactive profiles of E2 and 8-PN in vitro is noteworthy in light of their somewhat distinctive effects in subchronic exposure in vivo (see below).

In a recent study employing the human endometrial adenocarcinoma cell line, which expresses both ER $\alpha$  and ER $\beta$ , Ishikawa cells, alkaline phosphatase (AP) activity was used

as a measure of estrogenic action [76]. The researchers were interested in finding out whether the prenylated compounds in beer might have functional interactions with one another or with pesticides whose residues may occur in beer. They discovered that XN and IX exhibited from additive to slightly synergistic combinatory effect with 8-PN, and the mixture reached its peak induction of AP activity (about 80% of that caused by 1 nM E2) at a dose which corresponded to 10% of the estimated average dose in beer. A slightly synergistic interaction was also recorded between the mixture of 8-PN, IX, and XN and that of the pesticides. In contrast to this outcome, 8-PN did not modify (or subtly inhibited) the estrogenic activities of two fungal xenoestrogens, zearalenone and  $\alpha$ -zearalenol in Ishikawa cells. XN antagonized the fungal products more substantially [77].

### 5.2. In Vivo Studies

In in vivo conditions, the estrogenicity of 8-PN may be less prominent than in vitro. In juvenile male medaka fish, the positive control compound ethinylestradiol at the concentration of 0.34 nM induced sex reversal as indicated by the expression of the ovary-specific gene *figla* and the estrogen-dependent gene *vitellogenin*. However, 674 nM 8-PN (i.e., ~2000-fold concentration) failed to cause either of these effects [64]. When assessed by uterine vascular responses in ovariectomized (OVX) mice after subcutaneous (sc.) administration, the estrogenic activity of 8-PN was only <1% compared with E2. It was even lower (<0.1%) based on mitotic activity of vaginal epithelia upon peroral treatment. Moreover, in contrast to E2, 8-PN failed to induce mitosis in uterine epithelium or increase uterus weight in OVX mice at oral doses of up to 16 mg/kg/day for 3 days. [5]. This may be a species-dependent failure, because in OVX Wistar rats treated for 3 days with 10 mg/kg/day of either 8-PN or genistein, 8-PN proved to cause a greater elevation in uterus weight (~2 times the control weight) than genistein (~1.4 times the control weight) [68]. Exactly the same dosing regimen for 8-PN and animal model was also applied in another study in which the aim was to compare the impacts of E2 and 8-PN in vivo [78]. E2 at 30  $\mu$ g/kg/day increased uterine wet weight as well as epithelial height in the uterus and vagina and altered expression of estrogen-responsive genes in the uterus and liver. 8-PN elicited an identical pattern of responses, but the effects were blunted except for the induction of *insulin-like growth factor binding protein-1* (IGF-1) gene expression in the liver, which was more prominent upon 8-PN than E2 treatment.

Similarly, sc. exposure of young adult OVX Lewis rats to 15 mg/kg/day 8-PN for three days induced uterine weight 1.8-fold and elevated the height of the glandular epithelium [79]. However, in identical experimental conditions, E2 at the dose of 4  $\mu$ g/kg/day increased uterine wet weight by 4-fold augmenting the height of the luminal and glandular epithelia as well as myometrial thickness. In the vagina, 8-PN failed to stimulate the epithelial thickness but increased the percentage of proliferation cell nuclear antigen (PCNA)-positive epithelial cells as quantified by immunohistochemistry, which implies that epithelial cell proliferation was promoted all the same. This research team also assessed by an identical experimental setting the effect of 8-PN on the mammary gland [66]. Here, 8-PN induced the formation of terminal end buds and stimulated expression of proliferation markers in epithelial ductal cells with the same pattern as 4  $\mu$ g/kg/day E2, albeit in a much smaller magnitude [66]. The only exception was progesterone receptor, whose gene expression was induced by E2 alone.

A time-course study was conducted in OVX Wistar rats by exposing sc. the animals to 10  $\mu$ g/kg/day of E2 or 15 mg/kg/day of 8-PN and harvesting the uteri at 7, 24, or 72 h [80]. E2 elevated uterus wet weight and modified the expression levels of estrogen-sensitive genes in this tissue. The temporal and directional patterns in these variables were highly similar after 8-PN administration. In another rat study with a single dose (amount not reported) to juvenile female animals, 8-PN proved to be 20,000 times weaker than E2 in its ability to stimulate the mass of uterus or vagina [63]. It is also relevant in this context that in a rat model of menopausal hot flashes, represented by an elevation in the tail skin temperature of estrogen-deprived (OVX) rats, a 6-day oral or sc. administration of 8-PN

was able to restore the temperature into the normal range. The effect was similar to that caused by E2 at a 100-fold lower dose and blocked by an ER antagonist [81]. Overall, it seems that in qualitative terms, 8-PN closely resembles E2 in its profile of short-term in vivo effects but possesses a lower relative potency than in in vitro conditions, presumably due to its rapid metabolism.

The estrogenic effects of 8-PN have further been assessed after a longer exposure period. In OVX rats treated with 30 mg/kg/day 8-PN for two weeks, 8-PN was reported to increase uterus weight and also suppress ovariectomy-induced bone loss [82]. Treatment of adult OVX Sprague–Dawley rats with 4 or 40 mg/kg/day IP for 21 days increased uterus weight (although less than 10 µg/rat/day E2), while 0.4 mg/kg/day was ineffective [83]. The three 8-PN treatment levels dose-dependently enhanced uterine epithelial cell height, with the effect maximally amounting to about half of the impact of E2. In a subchronic study, OVX Sprague–Dawley rats were treated with either E2 (0.17 or 0.7 mg/kg/day) or 8-PN (6.8 or 68.4 mg/kg/day) added in their soy-free chow for three months, and uterine, vaginal, and mammary gland samples were analyzed histologically [84]. Compared with untreated OVX rats, the higher 8-PN dose augmented uterine weight (by increasing endo- and myometrial thickness), but the low and high E2 treatments were 1.7 and 2.0 times, respectively, as effective in this regard. The lower 8-PN dose did not morphologically affect the uterus, while the high doses of E2 and 8-PN led to the formation of squamous metaplasia, cystic glands, and hyperplastic/hypertrophic glands (the last ones were also detected in the low E2 group). Interestingly, two changes occurred exclusively in a single group. Forty-five percent of the rats in the high E2 group exhibited pyometra, whereas 60% of the high 8-PN rats showed polypoid structures with cysts. A clear distinction between the two compounds was also seen in the vagina. Whereas all E2- and no 8-PN-exposed rats displayed keratinization, vacuolization was only recorded in 8-PN groups (clear in 90% of the high dose group, incipient in 55% of the low and 10% of the high dose groups). In the mammary gland, there was luminal formation clearly observable in all animals treated with E2 and in 40% of the high-8-PN group. Thus, after subchronic exposure in rats, the estrogenic effect profile of 8-PN appears to be largely but not completely overlapping with E2 [84].

**Table 2.** Estrogenic activity of 8-PN.

<b>1. In Vitro</b>								
Test Assay	Type of 8-PN	Variable Measured	8-PN	E2	Coumestrol	Genistein	Daidzein	Reference
Binding to ER in rat uterine cytosol	Racemic	Relative affinity	0.023	1	0.008	0.003		[7]
Stimulation of alkaline phosphatase activity in Ishikawa cells	Racemic	EC <sub>50</sub> (nM)	4	0.8	30	200	1500	[7]
ERE-reporter gene expression in yeast cells transfected with the human ER $\alpha$	Racemic	EC <sub>50</sub> (nM)	40	0.3	70	1200	2200	[7]
Human ER $\alpha$ binding in vitro	(R)-8-PN	Relative affinity	~0.01	1				[5]
Human ER $\alpha$ binding in vitro	(S)-8-PN	Relative affinity	~0.01	1				[5]
Human ER $\alpha$ binding in vitro	(R)-8-PN	Relative affinity	~0.01	1				[5]
Human ER $\alpha$ binding in vitro	(S)-8-PN	Relative affinity	~0.01	1				[5]
Inhibition of E2 binding to ER in MCF-7 cells	Racemic	Molar excess over E2	45		35	1000	>10 <sup>4</sup>	[62]
ERE-CAT induction in MCF-7 cells	Racemic	Relative potency	100	1	330	250	3300	[62]
Proliferation of MCF-7 cells after 7 days	Racemic	Relative potency	3.3	1	500	500	5000	[62]
ERE-reporter gene expression in yeast cells transfected with the human ER $\alpha$	Racemic	EC <sub>50</sub> (nM)	0.02	0.005		40		[85]
Proliferation of MCF-7 cells after 24 h	Racemic	EC <sub>50</sub> (nM)	5	0.2		830		[85]
ERE-reporter gene expression in yeast cells transfected with the human ER $\alpha$	Racemic	EC <sub>50</sub> (nM)	100	0.8	140	2000	No resp.	[67]
ERE-reporter gene expression in yeast cells transfected with the human ER $\beta$	Racemic	EC <sub>50</sub> (nM)	33	0.2	3	8	800	[67]
Human ER $\alpha$ binding in vitro	Racemic	IC <sub>50</sub> (nM)	510	20		300	17,000	[61]
Human ER $\beta$ binding in vitro	Racemic	IC <sub>50</sub> (nM)	1700	15		20	1200	[61]
Human ER $\alpha$ binding in vitro	Racemic	IC <sub>50</sub> (nM)	59	11				[66]
Human ER $\beta$ binding in vitro	Racemic	IC <sub>50</sub> (nM)	65	4.2				[66]
ERE-reporter gene expression in yeast cells transfected with the human ER $\alpha$	Racemic	EC <sub>50</sub> (nM)	130	0.8		9300		[64]
Binding to the ligand-binding domain of human ER $\alpha$ in vitro	Racemic	IC <sub>50</sub> (nM)	57	1.2		1145		[73]
Binding to the ligand-binding domain of human ER $\beta$ in vitro	Racemic	IC <sub>50</sub> (nM)	68	1.4		25		[73]
<b>2. In vivo</b>								
Assay used	Type of 8-PN	Variable measured	8-PN	E2	Coumestrol	Genistein	Daidzein	Reference
Vascular permeability in mouse uterus	Racemic	Relative potency	<0.01	1	<0.01	<0.001	No effect	[5]
Vaginal epithelial mitosis in OVX mice	Racemic	Relative potency	<0.001	1				[5]

## 6. Endocrine Roles of Estrogen and 8-PN

Secondary metabolites from plants, including 8-PN, have been considered as alternatives to the classic hormone therapy in women. Studies on the endocrine properties of 8-PN have demonstrated that this compound is a natural selective estrogen receptor modulator (SERM) because its effect spectrum is not fully identical with that of E2. As already referred to earlier, E2 induces cellular changes through nuclear and non-nuclear mechanisms, and ER exists in 2 forms, ER $\alpha$  and ER $\beta$ , which have multiple isoforms and exhibit distinct tissue expression patterns and functions [86,87]. The classical nuclear mechanism of ER's action typically occurs within hours, leading to activation or repression of target genes. However, estrogens can also induce rapid signals that act within seconds or minutes through extranuclear, membrane-associated forms of ERs as well as a G protein-coupled estrogen receptor (GPCR1; also called GPCR30) [88].

Despite the similar affinity of the two well-established SERMs tamoxifen and endoxifen to both isoforms of ERs [89], SERMs usually exhibit preference to one isoform of ER over the other one, which may explain the varying and tissue-specific responses to SERMs. As described earlier, the majority of phytoestrogens, inclusive of coumestrol, genistein, apigenin, naringenin, and kaempferol, have been reported to display higher affinity towards ER $\beta$  than to ER $\alpha$  [90–92]. 8-PN is one of the very first known ER $\alpha$ -preferring phytoestrogens. Moreover, the ratio of ER $\alpha$ /ER $\beta$  abundances in target organs influences the overall action of SERMs in that tissue. In many tissues, ER $\beta$  receptors demonstrate antiproliferative actions, while ER $\alpha$  receptors mediate the opposite effect [93]. Therefore, an increased ratio of ER $\alpha$ /ER $\beta$  correlates well with high levels of cellular proliferation [94]. ERs and GPCR1 are abundantly expressed in central and peripheral tissues, reflecting the multifunctional nature of endo- and exogenous estrogens [58]. Since 8-PN has preference for ER $\alpha$ , we will here focus on the tissue expression of ER $\alpha$  as well as the potential endocrine role of ER $\alpha$ –8-PN complex in the contexts of energy metabolism, pituitary function and bone physiology.

### 6.1. ER $\alpha$ Tissue Expression

ER $\alpha$  is mainly expressed in endometrium, ovarian stroma, bone, mammary gland, placenta, pancreas, skeletal muscle, white and brown adipose tissues, and various neuroendocrine areas of the brain [95–97]. Strikingly, ER $\alpha$  is abundantly present in the ventrolateral portion of the ventromedial nucleus, the arcuate nucleus and the paraventricular nuclei of the hypothalamus as well as the medial preoptic area, whereas ER $\beta$  is significantly less expressed in these locations [98–103]. ER $\alpha$  occurs to a variable degree in all the secretory populations of the anterior pituitary neuroendocrine cells with a higher density in lactotrophs, gonadotrophs and somatotrophs and a lower density in thyrotrophs [104]. Therefore, the activation of ER $\alpha$  by potent exogenous ER $\alpha$  agonists, such as 8-PN, may interfere with the regulation of energy metabolism, glucose homeostasis, and other endocrine milieus, as will be discussed next.

### 6.2. Energy Metabolism

#### 6.2.1. Effects of 8-PN and Related Compounds on Energy Balance

8-PN treatment in drinking water at a dose of 10 mg/L/day for 20 weeks was reported to lead to inhibited body weight gain in OVX mice when compared with the control OVX group, an effect similar to that observed in estrogen-treated OVX mice. The lower body weight gain was accompanied by food intake reduction in 8-PN-treated OXV mice [105,106]. The majority of studies of naringenin (8-PN precursor) supplementation for at least 12 weeks to obese rodent models demonstrated a reduction in body weight gain without affecting food intake [107–110]. On the other hand, naringenin stimulated the release of the anorexigenic cholecystokinin hormone from intestinal secretin tumor cell line (STC-1) in vitro (secretion of CCK 12.5-fold at 0.1 mM, 23.0-fold at 0.5 mM, and 45.9-fold at 1.0 mM compared with control) [111]. Hence, these studies show that 8-PN and its precursor influence body weight gain, but the mechanism warrants further research.

### 6.2.2. Effects of 8-PN and Related Compounds on Lipid Metabolism

Provided in drinking water, 8-PN at the dose of 10 mg/L/day for 20 weeks ameliorated plasma lipid profile in a high-fat diet-induced type 2 diabetes mouse model. The changes triggered included reduced total cholesterol and triglyceride levels and enhanced HDL concentration and HDL/LDL ratio in plasma. In the same study, 8-PN activated AMPK signaling, thereby inhibiting SREBP-1c protein expression and its downstream lipogenic enzyme targets FAS and ACC [105]. Additionally, liver and muscle take-up fatty acids mainly through the translocase CD36 [112]. Vascular endothelial growth factor receptor-1 (VEGFR-1) and its ligand VEGF-B induce the expression of fatty acid transport proteins involved in lipid transport and uptake from blood to tissue [113]. In the afore-mentioned mouse study [105], the authors also recorded downregulation of hepatic and muscle CD36, VEGFR-1, and VEGF-B by 8-PN, which may have resulted in prevention of ectopic lipid accumulation [106].

In another study, 8-PN along with XN and IX were shown to be potential ligands for human farnesoid X receptor (FXR), based on collective findings from fluorescence titration, molecular docking studies and hydrogen deuterium exchange mass spectrometry. Since FXR activates the repressive orphan nuclear receptor and small heterodimer partner, thereby inhibiting gluconeogenesis and de novo lipogenesis, the observed anti-lipogenic and glucose-lowering (see below) effects of 8-PN might occur by a mechanism involving FXR [114].

### 6.2.3. Effects of 8-PN and Related Compounds on Glucose Homeostasis and Insulin Sensitivity

Similar to E2, 8-PN has also been reported to exert an antidiabetic effect. It lowered plasma glucose levels and improved glucose handling in oral glucose and insulin tolerance tests in a type 2 diabetes mouse model. It was further shown to augment the protein abundance of the insulin-regulated signaling molecule AS160 in skeletal muscle, suggesting improvement of glucose uptake by this tissue [106]. Insulin stimulates the PI3K/AKT signaling cascade which activates AS160 downstream, triggering GLUT4 translocation to cell membrane [115].

Regarding humans, naringenin supplementation (150 mg t.i.d for 8 weeks) proved to reduce body weight and insulin resistance as well as upregulate PPAR $\alpha$  and PPAR $\gamma$  [110].

Taking together, although there is a paucity of data available regarding the metabolic and endocrine impacts of 8-PN, this phytoestrogen appears a promising therapeutic candidate in protection against diet-induced obesity and metabolic dysfunctions.

## 6.3. Pituitary Function

### 6.3.1. Effects on LH and FSH Secretion

Similar to E2 at a concentration of  $10^{-9}$  M, 8-PN at a 1000-fold higher concentration directly suppressed LH release in a primary culture of rat pituitary cells. This response was significantly antagonized by the pure ER antagonist ICI 182,780 (Fulvestrant, Faslodex) [105]. Interestingly, high repeated oral dosing of 8-PN in diet (68.4 mg/kg/day for 12 weeks) to OVX rats also suppressed serum LH. In magnitude, serum LH was damped by 8-PN as effectively as by a treatment with 0.17 mg/kg BW/day E2, suggesting that a long-term intake of 8-PN could counteract the hot flashes [116]. Noteworthy in the same study, serum FSH concentration was lowered by a high E2 dose (0.7 mg/kg/day) alone. At the pituitary level, 8-PN and the lower E2 dose reduced the mRNA abundances of  $\alpha$ - and  $\beta$ -subunits of LH and gonadotropin-releasing hormone (GnRH) receptor. ER $\beta$  mRNA was increased by 8-PN and the higher dose of E2 [116].

The lack of E2 in postmenopausal women causes changes in the release of neurotransmitters involved in the regulation of the hypothalamic GnRH pulse generator, and thereby results in overactivity of the pulse generator [117]. These affected neurotransmitters spill over into adjacent neurons involved in the regulation of temperature and heart beats, inducing hot flashes [118,119]. Therefore, LH suppression by estrogenic compounds

alleviates this discomfort. Similar to what was reported in rats [116], 8-PN did not affect circulating FSH concentrations. The single doses tested (50–750 mg) were also well tolerated in postmenopausal women [37]. In the same study, the highest of these doses was shown to significantly decrease serum LH concentration in these women. This finding supports the view that the *in vitro* and *in vivo* findings recorded in rats and rat cells on 8-PN-induced inhibition of LH secretion (see above) are also relevant to humans, which forms a promising basis for the treatment of menopausal symptoms. However, this seems to require a high dose of 8-PN. Further support to the potential of 8-PN as a future remedy to hot flashes in women provides the finding that daily intake of a hop extract standardized on 8-PN (100 or 250 µg) exerted an alleviating therapeutic effect on vasomotor and other menopausal discomforts in women [120].

### 6.3.2. Effects on Other Pituitary Hormones

E2 or 8-PN at concentrations of  $10^{-9}$  and  $10^{-6}$  M, respectively, directly induced TSH release by a primary culture of rat pituitary cells [105]. This response was significantly antagonized by the ER antagonist ICI 162,780. In the *in vivo* part of this study, a single oral dose of E2 (15.5 mg/kg BW) or 8-PN (161.4 mg/kg BW) elevated the circulating TSH level in OVX rats. Concurrently, E2 showed lowering effects on free and bound T3 while 8-PN demonstrated a declining tendency for total T3 alone. In stark contrast to the response to a single dose, a high daily dose of 8-PN (68.4 mg/kg BW/day for 12 weeks) increased circulating total T3 in OVX rats. However, neither E2 nor 8-PN at any oral doses tested affected total T4 or free T4 in OVX rats [105].

ER $\alpha$  is also expressed in growth hormone (GH)-immunoreactive cells in the pituitary, and radiolabeled estrogen can be found in these cells [121]. Circulating GH stimulates IGF-1 production in multiple tissues including the liver and bone [122], and IGF-1 acts as a downstream mediator of GH effects, exerting feedback inhibition on GH secretion [123]. 8-PN at a concentration of 0.5 µM directly suppressed GH synthesis in GH3 pituitary adenoma cells of the rat [124]. In light of this finding, it was highly surprising that a 3-month treatment with either E2 (0.7 mg/kg/day) or 8-PN (68.4 mg/kg/day) increased serum GH in OVX rats. However, this probably resulted from weakened feedback inhibition by IGF-1, whose serum concentration was decreased by both E2 and 8-PN [125]. An identical outcome was previously reported in estrogen-treated postmenopausal women [123].

## 7. Regulation of Bone Homeostasis

One of the common consequences of estrogen decline in menopause is loss of bone mass that may lead to osteoporosis. Similar to E2, 8-PN and naringenin were reported to promote osteogenic differentiation *in vitro* [126]. In order to gain further insight into the importance of the prenyl group in 8-PN for its antiosteoporotic effects, 8-PN and naringenin were compared in *in vitro* conditions [127]. 8-PN was found to have a stronger ability than naringenin to improve osteoblast differentiation and osteogenic function in cultured rat calvarial osteoblasts. Concomitantly, 8-PN was more effective in inhibiting osteoclastogenesis, inducing osteoclast apoptosis and reducing the resorptive activity of osteoclasts in rabbit bone marrow cells, thus confirming the importance of the prenyl group in the naringenin structure for the bone protective mechanism. In another *in vitro* study, 8-PN closely mimicked E2 by enhancing osteoblast activity in the MC3T3-E1 osteoblast cell line and inhibiting osteoclastic differentiation from the multinuclear macrophage RAW264.7 [128]. In this study, the effects of 8-PN could be largely abolished by the selective ER $\alpha$  antagonist MPP but not with the selective ER $\beta$  antagonist PHTPP. The magnitude of 8-PN effects was smaller than that of a 1000-fold lower concentration of E2, but larger compared with equimolar concentrations of genistein and daidzein.

One of the first *in vivo* studies on the effects of 8-PN on bone metabolism showed that in OVX rats, a 28-day *sc.* treatment with 8-PN at the highest dose tested (18 mg/kg/day) prevented OVX-induced trabecular bone loss. Furthermore, there was no difference between the two 8-PN enantiomers in their capabilities of inhibiting the OVX-simulated bone

loss [129]. Excellent results were also reported in a subsequent study in OVX rats [130]. Here, the duration of treatment (via feed) was 12 weeks. At 68.4 mg/kg/day, 8-PN improved the biomechanical properties of bone much to the same degree as 0.7 mg/kg/day E2. 8-PN also increased cancellous bone mineral density, although not as effectively as E2. Overall, the two other phytoestrogens tested, genistein and resveratrol, proved less promising treatment alternatives for osteoporosis in this animal model.

In contrast to these positive outcomes, in another study conducted in OVX rats, daily sc. treatment with 1.77 mg/kg 8-PN for 10 weeks failed to bring about any noticeable improvement in the biomechanical properties or structure of osteoporotic lumbar vertebrae and femora, either alone or in combination with whole-body vertical vibration [131].

As to humans, a cross-sectional epidemiological study of some 1700 healthy women revealed that phalangeal bone ultrasound values were greater in beer drinkers compared with subjects avoiding beer drinking and/or drinking wine [132]. This association suggests that a modifying role for 8-PN in bone structure is also possible in humans, but due to the discrepant findings in rats, more studies are needed.

## 8. Effects on Tumor Cells In Vitro

A variety of in vitro studies have probed the impact of 8-PN and related hop flavonoids on cancer cell invasion, proliferation and apoptosis. In the majority of these studies, 8-PN along with its precursors and other naringenin derivatives displayed growth inhibitory and apoptotic effects in various cancer cell lines. For example, Delmulle et al. (2006) exposed the human prostate cancer cell lines PC-3 and DU145 to 8-PN, 6-PN, XN, IX and desmethyl-XN for 2 days. They found inhibited growth of the cells, with the following order of compound potency ( $IC_{50}$  [ $\mu M$ ] for DU145 and PC-3, respectively): XN (12.3, 13.2) > 6-PN (29.1, 18.4) > 8-PN (43.1, 33.5) > IX (47.4, 45.2) > desmethyl-XN (53.8, 49.9) [133]. In these cell lines, 8-PN, IX and 6-PN were shown to induce a caspase-independent form of cell death, suggested to be autophagy [134]. 8-PN and 6-PN also dose-dependently (6.25–100  $\mu M$ ) reduced proliferation of PC-3 and UO.31 renal carcinoma cells [135]. In the human leukemic T lymphocyte cell line Jurkat, the antiproliferative and proapoptotic effects of 8-PN were associated with inhibition of voltage-gated Kv1.3 potassium channels [136].

At concentrations of 40 and 50  $\mu M$ , respectively, 8-PN and IX decreased the viability of Caco-2 cells. While the toxicity of IX was associated with a concentration-dependent increase in G2/M ratio and an increased sub-G1 cell-cycle fraction, the treatment with 8-PN was associated with an elevated G0/G1 ratio and an increased sub-G1 cell-cycle fraction [137]. In another study, female hop flavonoids suppressed cell growth and induced apoptosis in the breast cancer Sk-Br-3 cell line 24 h post-incubation with  $IC_{50}$  values of 7.1, 22.6, and 41  $\mu M$  for XN, 8-PN and IX, respectively [138]. Likewise, 8-PN suppressed the proliferation of human Burkitt lymphoma cells at concentrations > 50  $\mu M$ . The degree of growth inhibition was 17% at 50  $\mu M$  and 41% at 100  $\mu M$  [139]. In clinical oncology, histone deacetylase (HDAC) inhibitors are currently investigated as new anticancer compounds. 100  $\mu M$  8-PN and 6-PN inhibited all 11 conserved human HDACs of classes I, II, and IV. Treatment of human melanoma SK-MEL-28 cells with 8-PN or 6-PN induced hyperacetylation of the histone complex H3 within 2 h. Furthermore, these compounds imparted a prominent, dose-dependent reduction of cellular proliferation and viability of SK-MEL-28 as well as BLM melanoma cells [140].

Some studies have reported differential impacts of 8-PN on malign and normal cells. For example, 8-PN and 6-PN displayed cytotoxic activity against the HeLa cervical cancer cells (with  $IC_{50}$  values ranging from 10 to 60  $\mu M$ ), but were less toxic to normal HL-7702 cells [141]. Similarly, 8-PN exhibited a strong growth inhibitory effect on human colorectal carcinoma HCT-116 cells with an  $IC_{50}$  value of 24  $\mu g/mL$  after 48 h. However, at similar concentrations and experimental timepoints, the compound did not show any cytotoxic effect to non-cancerous colon cells (CCD-41) [142]. Based on these studies, the prenyl group seems to be crucial to the anticancer activity of flavones, since it may lead to enhanced cell membrane targeting and thus increased intracellular activity [143].

In contrast to the antiproliferative and anticancer effects of 8-PN described above, some other studies have reported potentially untoward effects in cancer cell lines. For example, 8-PN at concentrations of  $10^{-5}$  and  $10^{-6}$  M was found to stimulate E-cadherin-dependent aggregation and growth of MCF-7/6 cells in suspension [144]. E-cadherin/ $\beta$ -catenin complexes are regulated by E2 at transcriptional and/or post-translational level in mammary cancer cell lines [144]. These complexes play an important role in maintaining epithelial integrity, and their aberrant expression is associated with a wide variety of human malignancies [145]. Similarly, in a 7-day colony-forming assay in the V79 fibroblast cell line, concentrations of 8-PN up to 10  $\mu\text{g}/\text{mL}$  ( $\approx 30 \mu\text{M}$ ) promoted cell proliferation by approximately 2-fold. At concentrations of  $\geq 30 \mu\text{g}/\text{mL}$  all cells died [146]. Likewise, 8-PN induced cell growth in H295R/MCF-7 co-culture with an  $\text{EC}_{50}$  of  $9.9\text{E-}6 \text{ mg}/\text{mL}$  even in the presence of the aromatase inhibitors letrozole or tamoxifen. This finding suggests that the use of phytoestrogen supplements should be avoided during breast cancer treatment [147].

Low toxicity of 8-PN was also reported in the RAW 264.7 murine macrophage cell line, where the  $\text{LC}_{50}$  for 8-PN was  $72 \mu\text{M}$  [148]. Furthermore, 8-PN at concentrations of 1–50  $\mu\text{M}$  failed to cause cytotoxicity in human acute myeloid leukemia (HL-60) cells and tended to increase metabolically active cells when tested in MCF-7 breast cancer cells [149].

In estrogen-responsive cancer cell lines, the concentration of 8-PN appears to be decisive to the outcome as shown by Brunelli et al. (2007). At concentrations below 10  $\mu\text{M}$ , 8-PN showed estrogenic properties, increasing the growth of ER positive MCF-7 human breast cancer cells, while higher concentrations of 8-PN inhibited proliferation and induced apoptosis [75].

## 9. Other Beneficial and Adverse Effects of 8-PN

### 9.1. Cytotoxicity to Somatic Primary Cells

Regarding human somatic primary cells, no decrease in cell viability was detected in human umbilical vein endothelial cells (HUVEC) or human aortic smooth muscle cells (HASMC) at any 8-PN concentration tested (highest 20  $\mu\text{M}$ ). In fact, the converse was true of HUVEC, in which 20  $\mu\text{M}$  8-PN caused a significant increase in viable cell number. Meanwhile, 10 and 20  $\mu\text{M}$  concentrations of XN and IX led to reduced viability of HUVEC and/or HASMC [150]. In keeping with these findings, apoptosis was dose-dependently inhibited by 8-PN but promoted by XN and IX in both cell lines. On the other hand, cell proliferation displayed opposite effects only in HUVEC (increased by 8-PN, decreased by XN and IX). In HASMC, XN, and IX inhibited proliferation and 8-PN exhibited the same tendency, albeit weaker. Overall, the in vitro toxicity of 8-PN to somatic cells seems to be low and it exhibits a proliferative tendency.

### 9.2. Effects on Gonadal Cells

Similarly to E2, 8-PN significantly stimulated capacitation and the acrosome reaction in incapacitated epididymal mouse spermatozoa, compared with untreated controls. Unexpectedly, however, 8-PN proved to be some 1000 times more potent than E2, being active already at low nM concentrations. The SERM compound hydroxytamoxifen failed to interfere with these actions. In capacitated sperm, E2 had no discernible effect whereas 8-PN was again able to induce the acrosome reaction. Both E2 and 8-PN enhanced sperm fertilizing ability in vitro [151]. Another phytoestrogen, genistein, exhibited the same effect pattern and potency, and low concentrations of genistein and 8-PN were more effective in combination than individually to accelerate capacitation [152]. Moreover, sensitivity to genistein was even higher for human than mouse sperm (sensitivities to 8-PN were not tested). As precocious acrosome reaction may impede fertility in vivo, these findings are alarming and call for further studies, especially in vivo.

8-PN and its precursor, IX, were shown to disrupt androgen production in rat Leydig cells. These compounds inhibited hCG-stimulated androgen synthesis at all stages of Leydig cell development and reduced cAMP production. However, they failed to suppress androgen production activated by an exogenous cAMP analog in adult and immature stages

of Leydig cells. These results thus suggest a cAMP-dependent cellular steroidogenesis effect of 8-PN and IX [153].

8-PN at concentrations of 1 and 3  $\mu\text{M}$  inhibited expansion of porcine cumulus oocytes in vitro while 3.12  $\mu\text{M}$  E2 did not. On the other hand, whereas that concentration of E2 diminished *Cyp19* (aromatase) gene expression by ~40% in these cells, 8-PN tended to increase it ~2-fold at all concentrations tested (0.1–3  $\mu\text{M}$ ; significantly at 0.3  $\mu\text{M}$ ). Both compounds impaired spindle formation, but the morphology of the meiotic spindles after these treatments was different [154]. Some of the recorded effects on oocytes may thus be non-ER-mediated, and 8-PN exhibits high in vitro potency in exerting them.

The developing oocytes are surrounded by granulosa cells in the ovary, and these also express aromatase. An extremely potent inhibitory impact on aromatase activity by 8-PN was reported in KGN human granulosa-like tumor cells, with an  $\text{IC}_{50}$  level of only 8 nM. Concomitantly, 8-PN (3  $\mu\text{M}$ ) induced *CYP19A1* gene expression about 3-fold but did not affect CYP19A1 protein abundance at 24 h, implying that 8-PN is a catalytic inhibitor of this enzyme [155]. Of note, none of the other phytoestrogens tested exhibited aromatase inhibition.

### 9.3. Effects on Aromatase in Other Cell Types

Aromatase is a major enzyme in the biosynthesis of steroids, catalyzing a critical step for estrogen production from circulating androgens. It is expressed not only in the gonads but also in adipose tissue, vasculature, bone, brain, placenta, fetal liver, and estrogen-dependent cancers [156]. In choriocarcinoma-derived JAR cells, which express high levels of aromatase, 8-PN proved to be a highly potent inhibitor of its activity. The  $\text{IC}_{50}$  was as low as 65 nM, while for XN and IX it was 20 and 140  $\mu\text{M}$ , respectively. However, no effect was recorded on aromatase gene expression (*CYP19*) [157]. A similarly potent inhibitory impact on aromatase activity was reported in the aromatase-expressing breast cancer cell line Sk-Br-3, in which the  $\text{IC}_{50}$  for 8-PN was 80 nM [138]. Slightly higher  $\text{IC}_{50}$  values were obtained in H295R human adrenocortical carcinoma cells, human placental microsomes and human breast fibroblasts, 100, 200, and 300 nM, respectively [147].

### 9.4. Effects on Other Enzymes of Clinical Relevance

AKR1B1 or human aldose reductase mediates the first step in the reduction of glucose to sorbitol in the polyol pathway, which under hyperglycemic conditions is co-responsible for the diabetic complications (i.e., retinopathy, neuropathy, nephropathy, cataract) [158]. Consequently, aldose reductase inhibitors are at the focus of research aiming at the prevention of these complications. The AKR1B1 homologue AKR1B10, in turn, is a NADPH-dependent oxidoreductase that converts carbonyl group containing compounds to their corresponding alcohols. It is overexpressed in several cancers and precancerous lesions, thus possibly playing a crucial role in the development of cancer [159]. 8-PN proved to inhibit both enzymes with  $\text{IC}_{50}$  values of 0.81–1.87 and 0.99–3.96  $\mu\text{M}$  for AKR1B1 and AKR1B10, respectively. IX was a roughly equipotent inhibitor with 8-PN while XN was about 10-fold less potent. None of them influenced the activity of the closely related AKR member AKR1A1 [160].

In addition to AKR1B1 and AKR1B10, in human chemoresistant cancer cell lines also carbonyl reductase 1 (CBR1) is upregulated and partially responsible for the resistance. Therefore, CBR1 inhibitors to be given adjuvantly with cytostatic therapy could be therapeutically advantageous. 8-PN consistently inhibited CBR1 activity with  $\text{IC}_{50}$  values ranging from 0.4 to 11.7  $\mu\text{M}$  depending on test system and substrate. XN and IX were up to 6-fold less potent in this regard [161].

8-PN did not exhibit appreciable acetylcholinesterase inhibition, but it did inhibit plasma cholinesterase, albeit modestly ( $\text{IC}_{50}$  = 86.6  $\mu\text{M}$ ) [162].

### 9.5. Effects on Blood Vessels

8-PN inhibited growth factor-induced angiogenesis of bovine endothelial cells in vitro, with an  $IC_{50}$  between 3 and 10  $\mu$ M. In the chicken early chorioallantois membrane, it reduced vessel diameter and tended to shorten vessel length. In both assay types, 8-PN was roughly equipotent to genistein [163].

In stark contrast to these findings, 8-PN was found to stimulate formation of capillary-like structures by primary HUVEC on GFR-Matrigel<sup>®</sup>, whereas XN and IX tended to inhibit it. Supporting evidence was obtained from in vivo experiments: in a mouse Matrigel<sup>®</sup> plug assay, IX and (partly) XN blocked the vascularizing effect of VEGF but 8-PN did not. In rat wound-healing assay, 8-PN enhanced angiogenesis in the wounded skin tissue whilst XN and IX impaired it. However, further blurring the view were findings from extracellular matrix invasion, an essential step in the angiogenic process. 8-PN, XN, and IX inhibited the invasive capacity of HASMC in a fairly similar manner at all three concentrations tested (0.1, 1, and 10  $\mu$ M). For HUVEC, IX led to the same outcome while XN did not impair invasiveness, and 8-PN did it only at 0.1  $\mu$ M concentration [150].

More recently, impacts of 8-PN and XN on angiogenesis were also assessed in a mouse model of type 2 diabetes brought about by feeding on a high-fat diet. In the kidney, diabetes induced neovascularization, whereas in the cardiac left ventricle it impaired the formation of new blood vessels. 8-PN and XN counteracted these phenomena in both tissues, with XN tending to be slightly more powerful in the kidney and 8-PN in the left ventricle. Concomitantly, these compounds acted to normalize diabetes-associated changes in VEGF-A and VEGF-B levels in these tissues, with the afore-mentioned subtle potency difference discernible in VEGF-A concentrations [164]. Altogether, the data available on 8-PN's impacts on angiogenesis are discrepant, suggesting tissue-specificity and non-monotonic dose-responses. Thus, additional studies are called for.

Prostacyclin ( $PGI_2$ ) is a critical vasculo- and cardioprotective agent [165]. A 10-nM concentration of 8-PN proved to double  $PGI_2$  production by HUVEC [148]. However, at higher concentrations the effect diminished, levelling off at 10  $\mu$ M 8-PN. This increase could be abolished by the selective COX-2 inhibitor NS-398. The authors therefore analyzed the effect of 8-PN on COX-2 protein expression in HUVEC. By 48 h, 10 nM 8-PN increased it by 55%. This change, in turn, could be blocked by the non-selective ER antagonist ICI 182780. Hence, the induction of  $PGI_2$  production by 8-PN appears to be attributable to its estrogenicity.

### 9.6. Effects on Inflammation and Immune Reactions

A high, pharmacological concentration of 8-PN (30  $\mu$ M) was shown to inhibit lipopolysaccharide (LPS)-induced gene expression in RAW 264.7 murine macrophage cells. A dose-response experiment revealed that already a 10- $\mu$ M concentration was almost as effective as 30  $\mu$ M in inhibiting the release of pro-inflammatory mediators ( $TNF\alpha$ , NO,  $PGE_2$ ). The mechanism appeared to involve NF- $\kappa$ B, whose LPS-induced DNA binding was virtually abolished by 10 and 30  $\mu$ M 8-PN [148].

In support of that report, 8-PN at ~7.5 or 15  $\mu$ M concentrations inhibited IL-12 secretion by murine splenic macrophages stimulated by LPS or LPS +  $IFN\gamma$ . 8-PN was equally effective to IX but less powerful than XN [166].

Overproduction of hyaluronan may occur and be harmful in, e.g., osteoarthritis. Hyaluronan is exported from fibroblasts and chondrocytes by the ATP-binding cassette transporter multidrug resistance associated protein 5 (MRP5) [167]. Therefore, inhibitors of MRP5 might have therapeutic value. 8-PN was found to inhibit hyaluronan export from bovine chondrocytes by MRP5 with an  $IC_{50}$  of 15  $\mu$ M. IX was again equipotent to it and XN twice as effective. A high concentration of 8-PN (50  $\mu$ M) also inhibited proteoglycan loss and collagen degradation induced by interleukins [168].

When applied topically daily on rat skin wounds for 7 days, 50  $\mu$ M 8-PN increased whereas XN and IX reduced the inflammation markers *N*-acetylglucosaminidase activity and IL1 $\beta$  levels in serum. This was associated with augmented formation of gran-

ulation tissue by 8-PN while XN and IX alleviated its formation [150]. The opposite effects of these compounds in this case are noteworthy even though they occurred at pharmacological doses.

#### 9.7. Effects on Platelets and Blood Coagulation

Phytoestrogens appear to impart beneficial effects against cardiovascular disease, mainly by inhibiting platelet function [169]. Moreover, 8-PN was demonstrated to inhibit platelet aggregation induced by different agonists and platelet adhesion to collagen matrix. Although 8-PN activated the inhibitory NO/cGMP/PKG/VASP pathway in human platelets, this pathway was not involved in 8-PN-mediated inhibition of platelet function. Instead, the authors concluded the inhibitory mechanism to rely on different downstream molecules of collagen and thrombin receptor engagements, including Pyk2, Akt, and ERK1/2. Of note, this effect was not mediated by ERs [170]. However, in postmenopausal women, daily treatment for 5 days with a standardized extract from spent hops did not interfere with blood clotting [171]. This may be ascribable to dosing or counteracting mixture effects, and additional research on this topic is needed.

#### 9.8. Effects on Muscle

In primary mouse myotubes in vitro, 8-PN at physiological concentrations activated the PI3K/Akt/P70S6K1 pathway. In vivo, dietary 8-PN accelerated muscle recovery from disuse atrophy and prevented the reduction of Akt phosphorylation in male C57BL/6 mice with immobilized right limb. Since E2 also enhanced the recovery of muscle mass, it is likely that the estrogenic activity of 8-PN was involved in its positive impact [172].

In another muscle disuse model in mice, 18-day preoperative 8-PN administration in feed reduced sciatic denervation-induced muscle atrophy. Akt phosphorylation was lowered by denervation in control mice, but 8-PN intake retained the phosphorylation status on Akt in denervated muscle comparable to that seen in sham-operated muscle. Denervation also induced the expression of atrogen-1 (a skeletal muscle-specific ubiquitin ligase), but 8-PN curtailed this increase by about 50% [173].

#### 9.9. Effects on Neurons and Behavior

The generation of new neurons in the adult brain from neural stem cells (neurogenesis) offers the potential for endogenous brain repair and functional regeneration. Adult neurogenesis is partially under the control of E2, and therefore 8-PN as a potent phytoestrogen has attracted research interest in this field.

In mouse embryonic forebrain cells, 10  $\mu$ M 8-PN exhibited no activity in promoting neuronal differentiation when assessed by a promoter assay after a three-day incubation period. However, after seven-day incubation, evidence of significant promotion was obtained by immunocytochemistry [174,175]. Additional studies with related compounds implied that estrogen-like activity of prenylflavanones can be dissociated from their activity of differentiation induction in neural precursor cells [175].

Using in silico prediction of molecular properties and molecular docking, Monteiro et al. (2018) [176] assessed 46 phytochemicals for their neuroprotective activity potential by virtue of binding to certain key targets for Parkinson's and Alzheimer's diseases, concluding that 8-PN is among the best drug candidates for both. Related to this, in contrast to four other compounds extracted from *Sophora flavescens*, 8-PN did not block the conversion of L-tyrosine to L-DOPA by tyrosinase in vitro [177].

To evaluate their potential as aids in insomnia and anxiety management, 8-PN, IX, and XN were subjected to a study on their effects on GABA<sub>A</sub> receptors. Of the three compounds, 8-PN proved the most potent at displacing the non-competitive GABA<sub>A</sub> receptor blocker ethynylbicycloorthobenzoate (EBOB) from native and recombinant GABA<sub>A</sub> receptors, both in the presence and absence of GABA. For example, the IC<sub>50</sub> values for the potentiation of GABA-induced displacement of [<sup>3</sup>H]EBOB in native GABA<sub>A</sub> receptors were 7.3, 11.6, and 29.7  $\mu$ M for 8-PN, IX, and XN, respectively. There was moderate GABA<sub>A</sub> receptor subtype

selectivity discernible. The GABAergic modulatory effects of these hop prenylflavonoids appeared not to be mediated via the high-affinity benzodiazepine binding site [178].

An in vivo study in rats yielded supporting evidence for behavioral modulation by 8-PN. When rats were treated with 10 mg/kg/day racemic 8-PN for 21 days, they exhibited prolonged escape latencies from the elevated T-maze open arm, implying a panicolytic effect. The positive control substance fluoxetine acted similarly. A subsequent docking study revealed that the *R* configuration of 8-PN had greater affinity to the transporters for serotonin, noradrenaline, and dopamine than does the *S* enantiomer, suggesting that the (*R*)-8-PN is the active form [179].

#### 9.10. Effects on Barrier Tissue Integrity

8-PN was capable of both preventing TNF $\alpha$ -induced epithelial disruption in the human intestinal epithelial cell line Caco-2 and restoring barrier integrity after TNF $\alpha$ -induced dysfunction, whereas XN and IX were ineffective and 6-PN only effective in the prevention phase. Based on these findings, 8-PN has potential as a remedy for a variety of diseases in which defective intestinal epithelial barrier function plays a major role, such as celiac disease, Crohn's disease, and inflammatory bowel disease [180].

TR146 cells are epithelial cells derived from human buccal carcinoma. Exposure of these cells to 5  $\mu$ M 8-PN or tamoxifen increased E-cadherin protein abundance in them, and irradiation of the cells did not interfere with this change. The E-cadherin/catenin complex at the membrane of normal epithelial cells is instrumental in maintaining cell–cell adhesion [181]. When TR146 cell aggregates were irradiated in vitro, the aggregate volume declined due to cell shedding. However, 5  $\mu$ M 8-PN or tamoxifen totally prevented the volume reduction. In vivo, mice treated topically on tongue mucosa with 8-PN or TAM displayed a significantly delayed onset of irradiation-induced oral mucositis and ulcers in the tongue [182]. Thus, 8-PN appears to be effective in maintaining epithelial barrier integrity also in the mouth.

#### 9.11. Antimicrobial Effects

8-PN also has antimicrobial properties. It showed significant activity against methicillin-sensitive and -resistant *Staphylococcus aureus* and *Staphylococcus epidermidis* strains with MIC<sub>80</sub> (the concentration that inhibits the growth of 80% of organisms) values of 25 or 50  $\mu$ g/mL. However, it proved to be inactive against *Listeria monocytogenes* and *Salmonella typhimurium* [183]. Likewise, it failed to inhibit the growth of *Escherichia coli* [184]. On the other hand, 8-PN was equally effective against both the drug-sensitive wildtype strain and two drug-resistant strains (resistant to pentamidine or to pentamidine and melarsoprol) of the single-celled parasite *Trypanosoma brucei* with an EC<sub>50</sub> of 6–7  $\mu$ g/mL [185].

8-PN proved to possess notable antifungal activity as well. For *Trichophyton mentagrophytes*, it had the same minimum inhibitory concentration (MIC) as the well-established dermatophytosis drug griseofulvin (6.25  $\mu$ g/mL). For *Trichophyton rubrum* and *Mucor rouxi-anus* the MIC was 12.5  $\mu$ g/mL. 8-PN showed no activity against the yeast *Candida albicans* or the plant pathogen fungus *Fusarium oxysporum* [8].

#### 9.12. Effects on Oxidative Stress

The data on the influence of 8-PN on oxidative stress are more discrepant than for most other phenomena. Soon after the discovery of 8-PN it was reported that at a 25  $\mu$ M—but not at a 5  $\mu$ M—concentration, 8-PN was able to inhibit oxidation of human low-density lipoprotein (LDL) induced by copper sulfate in vitro. IX and especially XN exhibited this ability already at the 5  $\mu$ M concentration [186]. However, no antioxidant activity by 8-PN was detected in the 2,2-diphenyl-1-picrylhydrazyl test or after menadione treatment of HL-60 cell cultures (tested concentration: 10  $\mu$ M) [149].

In contrast, pre-treatment with 8-PN reduced dose-dependently oxidative stress induced by LPS (0.1  $\mu$ g/mL; 12 h) in RAW 264.7 murine macrophage cells, abolishing it fully at 10  $\mu$ M concentration [148]. At the tested concentrations of 10 and 25  $\mu$ M, 8-PN

also displayed antioxidant activity in H<sub>2</sub>O<sub>2</sub>-treated H2P2G cells and in the 2,20-azino-bis(3-ethylbenzothiazoline)-6-sulfonic acid (ABTS) radical cation assay, respectively [175]. Moreover, in a N,N-dimethyl-*p*-phenylenediamine dihydrochloride (DMPD•<sup>+</sup>) scavenging assay 8-PN showed dose-dependent antioxidant activity, albeit not prominent, while naringenin was practically ineffective and 6-PN even demonstrated slight pro-oxidant effects at high concentrations [184].

In a mouse model of dietary type 2 diabetes, oxidative stress evolved in the liver and kidney as evidenced by 7- and 5-fold, respectively, augmented intrinsic fluorescence of advanced glycation end products (AGEs) in these tissues [187]. AGEs are modified proteins and/or lipids which are induced under oxidative conditions and which themselves enhance reactive oxygen species formation [188]. 8-PN treatment (0.1% in drinking water [~130 mg/kg/day]) was capable of fully preventing the AGE increase in the liver and of reducing it by approximately 50% in the kidney. XN also showed an inhibitory effect but less prominent than 8-PN. Another index of oxidative stress, protein nitration, was likewise elevated by diabetes in the liver and kidney (cortex). In this instance, 8-PN blocked the increase in both tissues whereas XN was only effective in the liver [187].

In contrast to the ABTS cation assay outcome described above, 8-PN (5–25 μM) was not able to reduce stable colored ABTS radicals *in vitro* in a more recent study. In the *in vivo* part of the same study, 8-PN at the tested 100 μM concentration slightly diminished intracellular accumulation of reactive oxygen species in *Caenorhabditis elegans*, but the inhibition did not reach statistical significance [189]. Overall, it thus appears that 8-PN has potential antioxidant activity, but this activity only emerges at relatively high concentrations.


#### 9.13. Modulation of Other Compound Toxicities

Unexpectedly, 8-PN augmented the formation of aflatoxin B1-DNA adducts in primary human hepatocytes (statistically significant change at 10 μM). 8-PN also proved to be a potent inhibitor of the human cytochrome P450 enzymes tested, with IC<sub>50</sub> values of 1.7, 4.5 and 8.4 for CYP1A1, CYP1A2 and CYP3A4, respectively. At the gene expression level, only the abundance of *CYP1A1* mRNA was altered (doubled by 10 μM 8-PN). The mechanism by which 8-PN enhances DNA adduct formation by aflatoxin B1 remained thus elusive [190].

## 10. Conclusions and Future Research Needs

8-PN is a potent phytoestrogen with a multitude of target organs and tissues. It also influences a wide variety of cellular signaling pathways, both ER-dependent and ER-independent (some examples are shown in Figure 2). While these effects have been repeatedly and convincingly established *in vitro*, the *in vivo* data are much more meagre. Furthermore, in most cases, the effects have required such high concentrations that they cannot be reached physiologically from dietary exposure to 8-PN and its precursors. The inter-individual differences in intestinal and hepatic generation of 8-PN from IX are yet noteworthy if hop-derived dietary supplements are used.

Getting back to our original question posed in the title of this review, is 8-PN ultimately a friend or a foe? So far, researchers have focused on emphasizing the health-promoting impacts of 8-PN, which indeed constitute an impressive list. However, critical analysis of the literature also reveals several less-favorable effects (Table 3), which should be kept in mind, especially if 8-PN is intended to be applied in medical therapy in the future. The most urgent information needs currently include the health consequences of chronic exposures to high doses of 8-PN (heavy beer drinkers with high IX conversion capability) and—closely related to this—its effects in men. Since hops have mainly found therapeutic use in the treatment of menopausal health problems in women, the great preponderance of human studies on 8-PN have understandably confined to female subjects. However, given that 8-PN has endocrine disrupting potency in males, beer is more commonly consumed by men than women, and that sperm has proven an exceptionally sensitive target to 8-PN, men should no longer be neglected in 8-PN studies.



**Figure 2.** Signaling cascades mediated by 8-PN in tissues and cells (BioRender.com, license purchased). First panel: 8-PN affected steroidogenesis through cAMP-dependent pathway in primary culture of porcine Leydig cells [153]. Second panel: in mouse skeletal muscle, 8-PN stimulated the PI3K/Akt signaling pathway at AS160, which triggers GLUT4 translocation to plasma membrane [106]. Third panel: 8-PN at physiological concentrations activated the PI3K/Akt/P70S6K1 pathway in mouse myotubes and accelerated muscle recovery from disuse atrophy [172]. Fourth panel: In mice, 8-PN activated AMPK, which plays a pivotal role in lipid and glucose metabolism in muscle and liver [106]. Fifth panel: 8-PN is a potential ligand for the human farnesoid X receptor (FXR), based on collective findings from fluorescence titration, molecular docking studies and hydrogen deuterium exchange mass spectrometry [114]. Sixth panel: 8-PN directly activated the inhibitory NO/cGMP/PKG/VASP pathway in human platelets [170]. Seventh panel: 8-PN rapidly activated ERK1/2 MAP kinase in MCF-7 cells [75].

**Table 3.** Potentially beneficial and adverse effects of 8-PN.

Beneficial	Adverse
- Antidiabetic	- Affects sperm with extraordinary potency
- Counteracts the metabolic syndrome	- Impairs spindle formation in oocytes and inhibits cumulus expansion
- Relieves hot flashes	- Disrupts androgen production in Leydig cells
- Inhibits the growth of many cancer cell lines	- Promotes the growth of some cancer cell lines
- Induces osteogenesis	- Aggravates inflammatory response topically in skin
- Inhibits AKR1B1 and AKR1B10	- Inhibits transporter proteins (ABCG2, ABCB1/P-gp, ABCC1/MRP1)
- Inhibits CBR1 activity	- Alters TSH, T3, GH and IGF-1 serum levels
- Induces PGI <sub>2</sub> production	- Inhibits aromatase in various cell types
- Mitigates LPS-induced effects	- Enhances DNA adduct formation by aflatoxin
- Inhibits MRP5	
- Inhibits platelet aggregation	
- Accelerates muscle recovery from disuse atrophy	
- Maintains barrier tissue integrity	
- Has antimicrobial properties	
- Promotes neuronal differentiation	
- Has high potential for neuroprotective activity	

**Author Contributions:** Conceptualization, R.P. and A.N.; figure designs, R.P. (Figure 1), A.N. (Figure 2); table designs, R.P.; writing, R.P. and A.N. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Only published and thus publicly available data were included.

**Acknowledgments:** Open access funding provided by University of Helsinki.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Tronina, T.; Popłoński, J.; Bartmańska, A. Flavonoids as phytoestrogenic components of hops and beer. *Molecules* **2020**, *25*, 4201. [CrossRef] [PubMed]
- Verzele, M. 100 years of hop chemistry and its relevance to brewing. *J. Inst. Brew.* **1986**, *92*, 32–48. [CrossRef]
- Goetz, P. Traitement des bouffées de chaleur par insuffisance ovarienne par l'extrait de houblon (*Humulus lupulus*). *Rev. Phytothérapie Prat.* **1990**, *4*, 13–15.
- Hesse, R.; Hoffmann, B.; Karg, H.; Vogt, K. Untersuchungen über den Nachweis von Phytoöstrogenen in Futterpflanzen und Hopfen mit Hilfe eines Rezeptortests 1. *Zent. Für Veterinärmedizin Reihe A* **1981**, *28*, 442–454. [CrossRef]
- Milligan, S.; Kalita, J.; Pocock, V.; Heyerick, A.; De Cooman, L.; Rong, H.; De Keukeleire, D. Oestrogenic activity of the hop phyto-oestrogen, 8-prenylnaringenin. *Reprod. -Camb.* **2002**, *123*, 235–242. [CrossRef]
- Fenselau, C.; Talalay, P. Is oestrogenic activity present in hops? *Food Cosmet. Toxicol.* **1973**, *11*, 597–603. [CrossRef]
- Milligan, S.; Kalita, J.; Heyerick, A.; Rong, H.; De Cooman, L.; De Keukeleire, D. Identification of a potent phytoestrogen in hops (*Humulus lupulus* L.) and beer. *J. Clin. Endocrinol. Metab.* **1999**, *84*, 2249. [CrossRef]
- Mizobuchi, S.; Sato, Y. A new flavanone with antifungal activity isolated from hops. *Agric. Biol. Chem.* **1984**, *48*, 2771–2775.
- Stevens, J.F.; Page, J.E. Xanthohumol and related prenylflavonoids from hops and beer: To your good health! *Phytochemistry* **2004**, *65*, 1317–1330. [CrossRef]
- Bolton, J.L.; Dunlap, T.L.; Hajirahimkhan, A.; Mbachu, O.; Chen, S.-N.; Chadwick, L.; Nikolic, D.; van Breemen, R.B.; Pauli, G.F.; Dietz, B.M.; et al. The multiple biological targets of hops and bioactive compounds. *Chem. Res. Toxicol.* **2019**, *32*, 222–233. [CrossRef]
- Karabin, M.; Hudcova, T.; Jelinek, L.; Dostalek, P. Biotransformations and biological activities of hop flavonoids. *Biotechnol. Adv.* **2015**, *33*, 1063–1090. [CrossRef] [PubMed]
- Kodama, S.; Yamamoto, A.; Sato, A.; Suzuki, K.; Yamashita, T.; Kemmei, T.; Taga, A.; Hayakawa, K. Enantioseparation of isoxanthohumol in beer by hydroxypropyl- $\gamma$ -cyclodextrin-modified micellar electrokinetic chromatography. *J. Agric. Food Chem.* **2007**, *55*, 6547–6552. [CrossRef] [PubMed]
- Nikolic, D.; B Van Breemen, R. Analytical methods for quantitation of prenylated flavonoids from hops. *Curr. Anal. Chem.* **2013**, *9*, 71–85. [CrossRef]
- Kirin Beer University. Global Beer Consumption by Country in 2018. Available online: [https://www.kirinholdings.com/en/newsroom/release/2019/1224\\_01.html](https://www.kirinholdings.com/en/newsroom/release/2019/1224_01.html) (accessed on 30 January 2022).
- Stevens, J.F.; Taylor, A.W.; Deinzer, M.L. Quantitative analysis of xanthohumol and related prenylflavonoids in hops and beer by liquid chromatography–tandem mass spectrometry. *J. Chromatogr. A* **1999**, *832*, 97–107. [CrossRef]
- Buckett, L.; Schinko, S.; Urmann, C.; Riepl, H.; Rychlik, M. Stable Isotope Dilution Analysis of the Major Prenylated Flavonoids Found in Beer, Hop Tea, and Hops. *Front. Nutr.* **2020**, *7*, 619921. [CrossRef] [PubMed]
- Possemiers, S.; Bolca, S.; Grootaert, C.; Heyerick, A.; Decroos, K.; Dhooge, W.; De Keukeleire, D.; Rabot, S.; Verstraete, W.; Van de Wiele, T.; et al. The prenylflavonoid isoxanthohumol from hops (*Humulus lupulus* L.) is activated into the potent phytoestrogen 8-prenylnaringenin in vitro and in the human intestine. *J. Nutr.* **2006**, *136*, 1862–1867. [CrossRef]
- Česlová, L.; Holčapek, M.; Fidler, M.; Drštičková, J.; Lisa, M. Characterization of prenylflavonoids and hop bitter acids in various classes of Czech beers and hop extracts using high-performance liquid chromatography–mass spectrometry. *J. Chromatogr. A* **2009**, *1216*, 7249–7257. [CrossRef]
- Golubović, J.B.; Heath, E.; Košir, I.J.; Ogrinc, N.; Potočnik, D.; Strojnik, L.; Heath, D. Differences in the levels of the selected phytoestrogens and stable isotopes in organic vs. conventional hops and beer. *Foods* **2021**, *10*, 1839. [CrossRef]
- Martinez, S.E.; Lakowski, T.M.; Davies, N.M. Enantiospecific analysis of 8-prenylnaringenin in biological fluids by liquid-chromatography-electrospray ionization mass spectrometry: Application to preclinical pharmacokinetic investigations. *Chirality* **2014**, *26*, 419–426. [CrossRef]
- Ito, C.; Mizuno, T.; Matsuoka, M.; Kimura, Y.; Sato, K.; Kajijura, I.; Omura, M.; JU-Ichi, M.; Furukawa, H. A new flavonoid and other new components from Citrus plants. *Chem. Pharm. Bull.* **1988**, *36*, 3292–3295. [CrossRef]
- Clarke, D.B.; Bailey, V.; Lloyd, A. Determination of phytoestrogens in dietary supplements by LC-MS/MS. *Food Addit. Contam.* **2008**, *25*, 534–547. [CrossRef] [PubMed]

23. Schretter, C.; Langeder, J.; Freisinger, V.; Rollinger, J.M.; Grienke, U. Quantitative Analysis of Prenylated Constituents in Commercial Hops Samples Using Ultrahigh-Performance Supercritical Fluid Chromatography. *Planta Med.* **2020**, *86*, 1140–1147. [[CrossRef](#)]
24. Coldham, N.; Sauer, M. Identification, quantitation and biological activity of phytoestrogens in a dietary supplement for breast enhancement. *Food Chem. Toxicol.* **2001**, *39*, 1211–1224. [[CrossRef](#)]
25. Dhooghe, L.; Naessens, T.; Heyerick, A.; De Keukeleire, D.; Vlietinck, A.J.; Pieters, L.; Apers, S. Quantification of xanthohumol, isoxanthohumol, 8-prenylnaringenin, and 6-prenylnaringenin in hop extracts and derived capsules using secondary standards. *Talanta* **2010**, *83*, 448–456. [[CrossRef](#)] [[PubMed](#)]
26. Tekel', J.; De Keukeleire, D.; Rong, H.; Daeseleire, E.; Van Peteghem, C. Determination of the hop-derived phytoestrogen, 8-prenylnaringenin, in beer by gas chromatography/mass spectrometry. *J. Agric. Food Chem.* **1999**, *47*, 5059–5063. [[CrossRef](#)]
27. Maragou, N.C.; Rosenberg, E.; Thomaidis, N.S.; Koupparis, M.A. Direct determination of the estrogenic compounds 8-prenylnaringenin, zearalenone,  $\alpha$ - and  $\beta$ -zearalenol in beer by liquid chromatography–mass spectrometry. *J. Chromatogr. A* **2008**, *1202*, 47–57. [[CrossRef](#)] [[PubMed](#)]
28. Yuan, Y.; Qiu, X.; Nikolic, D.; Dahl, J.H.; van Breemen, R.B. Method development and validation for ultra-high-pressure LC/MS/MS determination of hop prenylflavonoids in human serum. *J. AOAC Int.* **2012**, *95*, 1744–1749. [[CrossRef](#)]
29. Quifer-Rada, P.; Martínez-Huélamo, M.; Jáuregui, O.; Chiva-Blanch, G.; Estruch, R.N.; Lamuela-Raventós, R.M. Analytical condition setting a crucial step in the quantification of unstable polyphenols in acidic conditions: Analyzing prenylflavonoids in biological samples by liquid chromatography–electrospray ionization triple quadrupole mass spectrometry. *Anal. Chem.* **2013**, *85*, 5547–5554. [[CrossRef](#)]
30. Wyns, C.; Bolca, S.; De Keukeleire, D.; Heyerick, A. Development of a high-throughput LC/APCI-MS method for the determination of thirteen phytoestrogens including gut microbial metabolites in human urine and serum. *J. Chromatogr. B* **2010**, *878*, 949–956. [[CrossRef](#)]
31. Prencipe, F.P.; Brighenti, V.; Rodolfi, M.; Mongelli, A.; dall'Asta, C.; Ganino, T.; Bruni, R.; Pellati, F. Development of a new high-performance liquid chromatography method with diode array and electrospray ionization-mass spectrometry detection for the metabolite fingerprinting of bioactive compounds in *Humulus lupulus* L. *J. Chromatogr. A* **2014**, *1349*, 50–59. [[CrossRef](#)]
32. Moriya, H.; Tanaka, S.; Iida, Y.; Kitagawa, S.; Aizawa, S.I.; Taga, A.; Terashima, H.; Yamamoto, A.; Kodama, S. Chiral separation of isoxanthohumol and 8-prenylnaringenin in beer, hop pellets and hops by HPLC with chiral columns. *Biomed. Chromatogr.* **2018**, *32*, e4289. [[CrossRef](#)] [[PubMed](#)]
33. Schaefer, O.; Bohlmann, R.; Schleuning, W.-D.; Schulze-Forster, K.; Hümpel, M. Development of a radioimmunoassay for the quantitative determination of 8-prenylnaringenin in biological matrices. *J. Agric. Food Chem.* **2005**, *53*, 2881–2889. [[CrossRef](#)] [[PubMed](#)]
34. Wyns, C.; Derycke, L.; Soenen, B.; Bolca, S.; Deforce, D.; Bracke, M.; Heyerick, A. Production of monoclonal antibodies against hop-derived (*Humulus lupulus* L.) prenylflavonoids and the development of immunoassays. *Talanta* **2011**, *85*, 197–205. [[CrossRef](#)] [[PubMed](#)]
35. Philbin, C.S.; Schwartz, S.J. Resolution of diastereomeric flavonoid (1S)-(–)-camphanic acid esters via reversed-phase HPLC. *Phytochemistry* **2007**, *68*, 1206–1211. [[CrossRef](#)]
36. Calvo-Castro, L.A.; Burkard, M.; Sus, N.; Scheubeck, G.; Leischner, C.; Lauer, U.M.; Bosy-Westphal, A.; Hund, V.; Busch, C.; Venturelli, S.; et al. The Oral Bioavailability of 8-Prenylnaringenin from Hops (*Humulus Lupulus* L.) in Healthy Women and Men is Significantly Higher than that of its Positional Isomer 6-Prenylnaringenin in a Randomized Crossover Trial. *Mol. Nutr. Food Res.* **2018**, *62*, 1700838. [[CrossRef](#)]
37. Rad, M.; Hümpel, M.; Schaefer, O.; Schoemaker, R.; Schleuning, W.D.; Cohen, A.; Burggraaf, J. Pharmacokinetics and systemic endocrine effects of the phyto-estrogen 8-prenylnaringenin after single oral doses to postmenopausal women. *Br. J. Clin. Pharmacol.* **2006**, *62*, 288–296. [[CrossRef](#)] [[PubMed](#)]
38. Nikolic, D.; Li, Y.; Chadwick, L.R.; van Breemen, R.B. In vitro studies of intestinal permeability and hepatic and intestinal metabolism of 8-prenylnaringenin, a potent phytoestrogen from hops (*Humulus lupulus* L.). *Pharm. Res.* **2006**, *23*, 864–872. [[CrossRef](#)] [[PubMed](#)]
39. Nikolic, D.; Li, Y.; Chadwick, L.R.; Grubjesic, S.; Schwab, P.; Metz, P.; van Breemen, R.B. Metabolism of 8-prenylnaringenin, a potent phytoestrogen from hops (*Humulus lupulus*), by human liver microsomes. *Drug Metab. Dispos.* **2004**, *32*, 272–279. [[CrossRef](#)]
40. Fang, J.-B.; Nikolić, D.; Lankin, D.C.; Simmler, C.; Chen, S.-N.; Ramos Alvarenga, R.F.; Liu, Y.; Pauli, G.F.; van Breemen, R.B. Formation of (2R)- and (2S)-8-Prenylnaringenin Glucuronides by Human UDP-Glucuronosyltransferases. *J. Agric. Food Chem.* **2019**, *67*, 11650–11656. [[CrossRef](#)] [[PubMed](#)]
41. Paraiso, I.L.; Plagmann, L.S.; Yang, L.; Zielke, R.; Gombart, A.F.; Maier, C.S.; Sikora, A.E.; Blakemore, P.R.; Stevens, J.F. Reductive Metabolism of Xanthohumol and 8-Prenylnaringenin by the Intestinal Bacterium *Eubacterium ramulus*. *Mol. Nutr. Food Res.* **2019**, *63*, 1800923. [[CrossRef](#)] [[PubMed](#)]
42. Bartmańska, A.; Tronina, T.; Huszcza, E. Biotransformation of the phytoestrogen 8-prenylnaringenin. *Z. Für Nat. C* **2010**, *65*, 603–606. [[CrossRef](#)] [[PubMed](#)]
43. Kim, H.J.; Kim, S.-H.; Kang, B.Y.; Lee, I.-S. Microbial metabolites of 8-prenylnaringenin, an estrogenic prenylflavanone. *Arch. Pharmacol. Res.* **2008**, *31*, 1241–1246. [[CrossRef](#)] [[PubMed](#)]
44. Bartmańska, A.; Tronina, T.; Huszcza, E. Transformation of 8-prenylnaringenin by *Absidia coerulea* and *Beauveria bassiana*. *Bioorganic Med. Chem. Lett.* **2012**, *22*, 6451–6453. [[CrossRef](#)]
45. Tronina, T.; Huszcza, E. Microbial sulfation of 8-prenylnaringenin. *Z. Für Nat. C* **2013**, *68*, 231–235. [[CrossRef](#)]

46. Coldham, N.; Horton, R.; Byford, M.; Sauer, M. A binary screening assay for pro-oestrogens in food: Metabolic activation using hepatic microsomes and detection with oestrogen sensitive recombinant yeast cells. *Food Addit. Contam.* **2002**, *19*, 1138–1147. [[CrossRef](#)]
47. Nikolic, D.; Li, Y.; Chadwick, L.R.; Pauli, G.F.; van Breemen, R.B. Metabolism of xanthohumol and isoxanthohumol, prenylated flavonoids from hops (*Humulus lupulus* L.), by human liver microsomes. *J. Mass Spectrom.* **2005**, *40*, 289–299. [[CrossRef](#)]
48. Guo, J.; Nikolic, D.; Chadwick, L.R.; Pauli, G.F.; van Breemen, R.B. Identification of human hepatic cytochrome P450 enzymes involved in the metabolism of 8-prenylnaringenin and isoxanthohumol from hops (*Humulus lupulus* L.). *Drug Metab. Dispos.* **2006**, *34*, 1152–1159. [[CrossRef](#)] [[PubMed](#)]
49. Bolca, S.; Possemiers, S.; Maervoet, V.; Huybrechts, I.; Heyerick, A.; Vervarcke, S.; Depypere, H.; De Keukeleire, D.; Bracke, M.; De Henauw, S.; et al. Microbial and dietary factors associated with the 8-prenylnaringenin producer phenotype: A dietary intervention trial with fifty healthy post-menopausal Caucasian women. *Br. J. Nutr.* **2007**, *98*, 950–959. [[CrossRef](#)]
50. Bolca, S.; Wyns, C.; Possemiers, S.; Depypere, H.; De Keukeleire, D.; Bracke, M.; Verstraete, W.; Heyerick, A. Cosupplementation of isoflavones, prenylflavonoids, and lignans alters human exposure to phytoestrogen-derived 17  $\beta$ -estradiol equivalents. *J. Nutr.* **2009**, *139*, 2293–2300. [[CrossRef](#)]
51. Possemiers, S.; Heyerick, A.; Robbens, V.; De Keukeleire, D.; Verstraete, W. Activation of proestrogens from hops (*Humulus lupulus* L.) by intestinal microbiota; conversion of isoxanthohumol into 8-prenylnaringenin. *J. Agric. Food Chem.* **2005**, *53*, 6281–6288. [[CrossRef](#)]
52. Possemiers, S.; Rabot, S.; Espín, J.C.; Bruneau, A.; Philippe, C.; González-Sarrías, A.; Heyerick, A.; Tomás-Barberán, F.A.; De Keukeleire, D.; Verstraete, W.; et al. Eubacterium limosum activates isoxanthohumol from hops (*Humulus lupulus* L.) into the potent phytoestrogen 8-prenylnaringenin in vitro and in rat intestine. *J. Nutr.* **2008**, *138*, 1310–1316. [[CrossRef](#)]
53. Hanske, L.; Loh, G.; Sczesny, S.; Blaut, M.; Braune, A. Recovery and metabolism of xanthohumol in germ-free and human microbiota-associated rats. *Mol. Nutr. Food Res.* **2010**, *54*, 1405–1413. [[CrossRef](#)] [[PubMed](#)]
54. Terao, J.; Mukai, R. Prenylation modulates the bioavailability and bioaccumulation of dietary flavonoids. *Arch. Biochem. Biophys.* **2014**, *559*, 12–16. [[CrossRef](#)] [[PubMed](#)]
55. Tan, K.W.; Cooney, J.; Jensen, D.; Li, Y.; Paxton, J.W.; Birch, N.P.; Scheepens, A. Hop-derived prenylflavonoids are substrates and inhibitors of the efflux transporter breast cancer resistance protein (BCRP/ABCG2). *Mol. Nutr. Food Res.* **2014**, *58*, 2099–2110. [[CrossRef](#)]
56. Wesołowska, O.; Wiśniewski, J.; Środa, K.; Krawczyński, A.; Bielawska-Pohl, A.; Paprocka, M.; Duś, D.; Michalak, K. 8-Prenylnaringenin is an inhibitor of multidrug resistance-associated transporters, P-glycoprotein and MRP1. *Eur. J. Pharmacol.* **2010**, *644*, 32–40. [[CrossRef](#)] [[PubMed](#)]
57. Caccamese, S.; Manna, L.; Scivoli, G. Chiral HPLC separation and CD spectra of the C-2 diastereomers of naringin in grapefruit during maturation. *Chirality Pharmacol. Biol. Chem. Conseq. Mol. Asymmetry* **2003**, *15*, 661–667. [[CrossRef](#)]
58. Zhang, J.-Q.; Cai, W.-Q.; Zhou, D.-S.; Su, B.-Y. Distribution and differences of estrogen receptor beta immunoreactivity in the brain of adult male and female rats. *Brain Res.* **2002**, *935*, 73–80. [[CrossRef](#)]
59. Takamura-Enya, T.; Ishihara, J.; Tahara, S.; Goto, S.; Totsuka, Y.; Sugimura, T.; Wakabayashi, K. Analysis of estrogenic activity of foodstuffs and cigarette smoke condensates using a yeast estrogen screening method. *Food Chem. Toxicol.* **2003**, *41*, 543–550. [[CrossRef](#)]
60. Zierau, O.; Gester, S.; Schwab, P.; Metz, P.; Kolba, S.; Wulf, M.; Vollmer, G. Estrogenic activity of the phytoestrogens naringenin, 6-(1,1-dimethylallyl) naringenin and 8-prenylnaringenin. *Planta Med.* **2002**, *68*, 449–451. [[CrossRef](#)]
61. Overk, C.R.; Yao, P.; Chadwick, L.R.; Nikolic, D.; Sun, Y.; Cuendet, M.A.; Deng, Y.; Hedayat, A.; Pauli, G.F.; Farnsworth, N.R.; et al. Comparison of the in vitro estrogenic activities of compounds from hops (*Humulus lupulus*) and red clover (*Trifolium pratense*). *J. Agric. Food Chem.* **2005**, *53*, 6246–6253. [[CrossRef](#)]
62. Matsumura, A.; Ghosh, A.; Pope, G.; Darbre, P. Comparative study of oestrogenic properties of eight phytoestrogens in MCF7 human breast cancer cells. *J. Steroid Biochem. Mol. Biol.* **2005**, *94*, 431–443. [[CrossRef](#)] [[PubMed](#)]
63. Schaefer, O.; Hümpel, M.; Fritze, K.-H.; Bohlmann, R.; Schleuning, W.-D. 8-Prenyl naringenin is a potent ER $\alpha$  selective phytoestrogen present in hops and beer. *J. Steroid Biochem. Mol. Biol.* **2003**, *84*, 359–360. [[CrossRef](#)]
64. Zierau, O.; Hamann, J.; Tischer, S.; Schwab, P.; Metz, P.; Vollmer, G.; Gutzeit, H.O.; Scholz, S. Naringenin-type flavonoids show different estrogenic effects in mammalian and teleost test systems. *Biochem. Biophys. Res. Commun.* **2005**, *326*, 909–916. [[CrossRef](#)] [[PubMed](#)]
65. Kretzschmar, G.; Zierau, O.; Wober, J.; Tischer, S.; Metz, P.; Vollmer, G. Prenylation has a compound specific effect on the estrogenicity of naringenin and genistein. *J. Steroid Biochem. Mol. Biol.* **2010**, *118*, 1–6. [[CrossRef](#)] [[PubMed](#)]
66. Helle, J.; Kräker, K.; Bader, M.I.; Keiler, A.M.; Zierau, O.; Vollmer, G.; Welsh, J.; Kretzschmar, G. Assessment of the proliferative capacity of the flavanones 8-prenylnaringenin, 6-(1,1-dimethylallyl) naringenin and naringenin in MCF-7 cells and the rat mammary gland. *Mol. Cell. Endocrinol.* **2014**, *392*, 125–135. [[CrossRef](#)]
67. Bovee, T.F.; Helsdingen, R.J.; Rietjens, I.M.; Keijer, J.; Hoogenboom, R.L. Rapid yeast estrogen bioassays stably expressing human estrogen receptors  $\alpha$  and  $\beta$ , and green fluorescent protein: A comparison of different compounds with both receptor types. *J. Steroid Biochem. Mol. Biol.* **2004**, *91*, 99–109. [[CrossRef](#)] [[PubMed](#)]
68. Amer, D.A.; Kretzschmar, G.; Müller, N.; Stanke, N.; Lindemann, D.; Vollmer, G. Activation of transgenic estrogen receptor-beta by selected phytoestrogens in a stably transduced rat serotonergic cell line. *J. Steroid Biochem. Mol. Biol.* **2010**, *120*, 208–217. [[CrossRef](#)]
69. van de Schans, M.G.; Ritschel, T.; Bovee, T.F.; Sanders, M.G.; de Waard, P.; Gruppen, H.; Vincken, J.P. Involvement of a hydrophobic pocket and helix 11 in determining the modes of action of prenylated flavonoids and isoflavonoids in the human estrogen receptor. *ChemBioChem* **2015**, *16*, 2668–2677. [[CrossRef](#)]
70. Amer, D.A.; Jähne, M.; Weigt, C.; Kretzschmar, G.; Vollmer, G. Effect of 17 $\beta$ -estradiol and flavonoids on the regulation of expression of newly identified oestrogen responsive genes in a rat raphe nuclei-derived cell line. *J. Cell. Physiol.* **2012**, *227*, 3434–3445. [[CrossRef](#)]

71. Bovee, T.F.; Schoonen, W.G.; Hamers, A.R.; Bento, M.J.; Peijnenburg, A.A. Screening of synthetic and plant-derived compounds for (anti) estrogenic and (anti) androgenic activities. *Anal. Bioanal. Chem.* **2008**, *390*, 1111–1119. [[CrossRef](#)]
72. Zierau, O.; Morrissey, C.; Watson, R.W.G.; Schwab, P.; Kolba, S.; Metz, P.; Vollmer, G. Antiandrogenic activity of the phytoestrogens naringenin, 6-(1, 1-dimethylallyl) naringenin and 8-prenylnaringenin. *Planta Med.* **2003**, *69*, 856–858. [[PubMed](#)]
73. Roelens, F.; Heldring, N.; Dhooze, W.; Bengtsson, M.; Comhaire, F.; Gustafsson, J.-Å.; Treuter, E.; De Keukeleire, D. Subtle side-chain modifications of the hop phytoestrogen 8-prenylnaringenin result in distinct agonist/antagonist activity profiles for estrogen receptors  $\alpha$  and  $\beta$ . *J. Med. Chem.* **2006**, *49*, 7357–7365. [[CrossRef](#)]
74. Bjornstrom, L.; Sjoberg, M. Mechanisms of estrogen receptor signaling: Convergence of genomic and nongenomic actions on target genes. *Mol. Endocrinol.* **2005**, *19*, 833–842. [[CrossRef](#)] [[PubMed](#)]
75. Brunelli, E.; Minassi, A.; Appendino, G.; Moro, L. 8-Prenylnaringenin, inhibits estrogen receptor- $\alpha$  mediated cell growth and induces apoptosis in MCF-7 breast cancer cells. *J. Steroid Biochem. Mol. Biol.* **2007**, *107*, 140–148. [[CrossRef](#)] [[PubMed](#)]
76. Aichinger, G.; Bliem, G.; Marko, D. Systemically Achievable Doses of Beer Flavonoids Induce Estrogenicity in Human Endometrial Cells and Cause Synergistic Effects With Selected Pesticides. *Front. Nutr.* **2021**, *8*, 691872. [[CrossRef](#)] [[PubMed](#)]
77. Aichinger, G.; Beisl, J.; Marko, D. The hop polyphenols xanthohumol and 8-prenyl-naringenin antagonize the estrogenic effects of fusarium mycotoxins in human endometrial cancer cells. *Front. Nutr.* **2018**, *5*, 85. [[CrossRef](#)]
78. Diel, P.; Thomae, R.B.; Caldarelli, A.; Zierau, O.; Kolba, S.; Schmidt, S.; Schwab, P.; Metz, P.; Vollmer, G. Regulation of gene expression by 8-prenylnaringenin in uterus and liver of Wistar rats. *Planta Med.* **2004**, *70*, 39–44. [[PubMed](#)]
79. Keiler, A.M.; Dörfelt, P.; Chatterjee, N.; Helle, J.; Bader, M.I.; Vollmer, G.; Kretschmar, G.; Kuhlee, F.; Thieme, D.; Zierau, O.; et al. Assessment of the effects of naringenin-type flavanones in uterus and vagina. *J. Steroid Biochem. Mol. Biol.* **2015**, *145*, 49–57. [[CrossRef](#)] [[PubMed](#)]
80. Zierau, O.; Kretschmar, G.; Möller, F.; Weigt, C.; Vollmer, G. Time dependency of uterine effects of naringenin type phytoestrogens in vivo. *Mol. Cell. Endocrinol.* **2008**, *294*, 92–99. [[CrossRef](#)] [[PubMed](#)]
81. Bowe, J.; Li, X.F.; Kinsey-Jones, J.; Heyerick, A.; Brain, S.; Milligan, S.; O'Byrne, K. The hop phytoestrogen, 8-prenylnaringenin, reverses the ovariectomy-induced rise in skin temperature in an animal model of menopausal hot flashes. *J. Endocrinol.* **2006**, *191*, 399–405. [[CrossRef](#)]
82. Miyamoto, M.; Matsushita, Y.; Kiyokawa, A.; Fukuda, C.; Sugano, M.; Akiyama, T. Prenylflavonoids: A new class of non-steroidal phytoestrogen (part 2). estrogenic effects of 18-isopentenyl naringenin on bone metabolism. *Planta Med.* **1998**, *64*, 516–519. [[CrossRef](#)] [[PubMed](#)]
83. Overk, C.R.; Guo, J.; Chadwick, L.R.; Lantvit, D.D.; Minassi, A.; Appendino, G.; Chen, S.-N.; Lankin, D.C.; Farnsworth, N.R.; Pauli, G.F.; et al. In vivo estrogenic comparisons of *Trifolium pratense* (red clover) *Humulus lupulus* (hops), and the pure compounds isoxanthohumol and 8-prenylnaringenin. *Chem. Biol. Interact.* **2008**, *176*, 30–39. [[CrossRef](#)] [[PubMed](#)]
84. Rimoldi, G.; Christoffel, J.; Wuttke, W. Morphologic changes induced by oral long-term treatment with 8-prenylnaringenin in the uterus, vagina, and mammary gland of castrated rats. *Menopause* **2006**, *13*, 669–677. [[CrossRef](#)]
85. Nasri, A.; Pohjanvirta, R. In vitro estrogenic, cytotoxic, and genotoxic profiles of the xenoestrogens 8-prenylnaringenin, genistein and tartrazine. *Environ. Sci. Pollut. Res.* **2021**, *28*, 27988–27997. [[CrossRef](#)] [[PubMed](#)]
86. Jensen, E.V. Basic guides to the mechanism of estrogen action. *Recent Progr. Horm. Res.* **1962**, *18*, 387–414.
87. Kuiper, G.; Enmark, E.; Peltö-Huikko, M.; Nilsson, S.; Gustafsson, J.-A. Cloning of a novel receptor expressed in rat prostate and ovary. *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 5925–5930. [[CrossRef](#)]
88. Gruber, C.J.; Tschugguel, W.; Schneeberger, C.; Huber, J.C. Production and actions of estrogens. *N. Engl. J. Med.* **2002**, *346*, 340–352. [[CrossRef](#)]
89. Lim, Y.C.; Li, L.; Desta, Z.; Zhao, Q.; Rae, J.M.; Flockhart, D.A.; Skaar, T.C. Endoxifen, a secondary metabolite of tamoxifen, and 4-OH-tamoxifen induce similar changes in global gene expression patterns in MCF-7 breast cancer cells. *J. Pharmacol. Exp. Ther.* **2006**, *318*, 503–512. [[CrossRef](#)]
90. Kuiper, G.G.; Lemmen, J.G.; Carlsson, B.; Corton, J.C.; Safe, S.H.; Van Der Saag, P.T.; Van Der Burg, B.; Gustafsson, J.-A.K. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor  $\beta$ . *Endocrinology* **1998**, *139*, 4252–4263. [[CrossRef](#)]
91. Ososki, A.L.; Kennelly, E.J. Phytoestrogens: A review of the present state of research. *Phytother. Res. Int. J. Devoted Pharmacol. Toxicol. Eval. Nat. Prod. Deriv.* **2003**, *17*, 845–869. [[CrossRef](#)]
92. Simons, R.; Gruppen, H.; Bovee, T.F.; Verbruggen, M.A.; Vincken, J.-P. Prenylated isoflavonoids from plants as selective estrogen receptor modulators (phytoSERMs). *Food Funct.* **2012**, *3*, 810–827. [[CrossRef](#)]
93. Koehler, K.F.; Helguero, L.A.; Haldosén, L.-A.; Warner, M.; Gustafsson, J.-A. Reflections on the discovery and significance of estrogen receptor  $\beta$ . *Endocr. Rev.* **2005**, *26*, 465–478. [[CrossRef](#)] [[PubMed](#)]
94. Jordan, V.C. Chemoprevention of breast cancer with selective oestrogen-receptor modulators. *Nat. Rev. Cancer* **2007**, *7*, 46–53. [[CrossRef](#)] [[PubMed](#)]
95. Pfaffl, M.; Lange, I.; Daxenberger, A.; Meyer, H. Tissue-specific expression pattern of estrogen receptors (ER): Quantification of ER $\alpha$  and ER $\beta$  mRNA with real-time RT-PCR. *Apmis* **2001**, *109*, S540–S550. [[CrossRef](#)]
96. Mauvais-Jarvis, F.; Clegg, D.J.; Hevener, A.L. The role of estrogens in control of energy balance and glucose homeostasis. *Endocr. Rev.* **2013**, *34*, 309–338. [[CrossRef](#)] [[PubMed](#)]
97. Hutson, D.D.; Gurralla, R.; Ogola, B.O.; Zimmerman, M.A.; Mostany, R.; Satou, R.; Lindsey, S.H. Estrogen receptor profiles across tissues from male and female *Rattus norvegicus*. *Biol. Sex Differ.* **2019**, *10*, 1–13. [[CrossRef](#)]

98. Simerly, R.; Swanson, L.; Chang, C.; Muramatsu, M. Distribution of androgen and estrogen receptor mRNA-containing cells in the rat brain: An in situ hybridization study. *J. Comp. Neurol.* **1990**, *294*, 76–95. [[CrossRef](#)] [[PubMed](#)]
99. Simonian, S.X.; Herbison, A.E. Differential expression of estrogen receptor alpha and beta immunoreactivity by oxytocin neurons of rat paraventricular nucleus. *J. Neuroendocrinol.* **1997**, *9*, 803–806. [[CrossRef](#)]
100. Voisin, D.; Simonian, S.; Herbison, A. Identification of estrogen receptor-containing neurons projecting to the rat supraoptic nucleus. *Neuroscience* **1997**, *78*, 215–228. [[CrossRef](#)]
101. Wilkinson, H.A.; Dahllund, J.; Liu, H.; Yudkovitz, J.; Cai, S.-J.; Nilsson, S.; Schaeffer, J.M.; Mitra, S.W. Identification and characterization of a functionally distinct form of human estrogen receptor  $\beta$ . *Endocrinology* **2002**, *143*, 1558–1561. [[CrossRef](#)]
102. Shima, N.; Yamaguchi, Y.; Yuri, K. Distribution of estrogen receptor  $\beta$  mRNA-containing cells in ovariectomized and estrogen-treated female rat brain. *Anat. Sci. Int.* **2003**, *78*, 85–97. [[CrossRef](#)]
103. Merchenthaler, I.; Lane, M.V.; Numan, S.; Dellovade, T.L. Distribution of estrogen receptor  $\alpha$  and  $\beta$  in the mouse central nervous system: In vivo autoradiographic and immunocytochemical analyses. *J. Comp. Neurol.* **2004**, *473*, 270–291. [[CrossRef](#)]
104. Manoranjan, B.; Salehi, F.; Scheithauer, B.; Rotondo, F.; Kovacs, K.; Cusimano, M. Estrogen receptors  $\alpha$  and  $\beta$  immunohistochemical expression: Clinicopathological correlations in pituitary adenomas. *Anticancer. Res.* **2010**, *30*, 2897–2904. [[PubMed](#)]
105. Christoffel, J. Endocrine potential of Genistein, Resveratrol and 8-Prenylnaringenin in Gonadal and Thyroid Axes and Related Organs. Ph.D. Thesis, Faculty of Mathematics and Natural Sciences, Christian-Albrecht University of Kiel, Kiel, Germany, 2004.
106. Costa, R.; Rodrigues, I.; Guardão, L.; Rocha-Rodrigues, S.; Silva, C.; Magalhães, J.; Ferreira-de-Almeida, M.; Negrão, R.; Soares, R. Xanthohumol and 8-prenylnaringenin ameliorate diabetic-related metabolic dysfunctions in mice. *J. Nutr. Biochem.* **2017**, *45*, 39–47. [[CrossRef](#)] [[PubMed](#)]
107. Mulvihill, E.E.; Allister, E.M.; Sutherland, B.G.; Telford, D.E.; Sawyez, C.G.; Edwards, J.Y.; Markle, J.M.; Hegele, R.A.; Huff, M.W. Naringenin prevents dyslipidemia, apolipoprotein B overproduction, and hyperinsulinemia in LDL receptor-null mice with diet-induced insulin resistance. *Diabetes* **2009**, *58*, 2198–2210. [[CrossRef](#)]
108. Assini, J.M.; Mulvihill, E.E.; Sutherland, B.G.; Telford, D.E.; Sawyez, C.G.; Felder, S.L.; Chhoker, S.; Edwards, J.Y.; Gros, R.; Huff, M.W.; et al. Naringenin prevents cholesterol-induced systemic inflammation, metabolic dysregulation, and atherosclerosis in Ldlr<sup>-/-</sup> mice [S]. *J. Lipid Res.* **2013**, *54*, 711–724. [[CrossRef](#)] [[PubMed](#)]
109. Assini, J.M.; Mulvihill, E.E.; Burke, A.C.; Sutherland, B.G.; Telford, D.E.; Chhoker, S.S.; Sawyez, C.G.; Drangova, M.; Adams, A.C.; Kharitonov, A.; et al. Naringenin prevents obesity, hepatic steatosis, and glucose intolerance in male mice independent of fibroblast growth factor 21. *Endocrinology* **2015**, *156*, 2087–2102. [[CrossRef](#)]
110. Murugesan, N.; Woodard, K.; Ramaraju, R.; Greenway, F.L.; Coulter, A.A.; Rebello, C.J. Naringenin increases insulin sensitivity and metabolic rate: A case study. *J. Med. Food* **2020**, *23*, 343–348. [[CrossRef](#)]
111. Park, M.; Kim, K.; Lee, Y.M.; Rhyu, M.R.; Kim, H.Y. Naringenin stimulates cholecystokinin secretion in STC-1 cells. *Nutr. Res. Pract.* **2014**, *8*, 146–150. [[CrossRef](#)]
112. Miquilena-Colina, M.E.; Lima-Cabello, E.; Sánchez-Campos, S.; García-Mediavilla, M.V.; Fernández-Bermejo, M.; Lozano-Rodríguez, T.; Vargas-Castrillón, J.; Buqué, X.; Ochoa, B.; Aspichueta, P.; et al. Hepatic fatty acid translocase CD36 upregulation is associated with insulin resistance, hyperinsulinaemia and increased steatosis in non-alcoholic steatohepatitis and chronic hepatitis C. *Gut* **2011**, *60*, 1394–1402. [[CrossRef](#)]
113. Hagberg, C.E.; Falkevall, A.; Wang, X.; Larsson, E.; Huusko, J.; Nilsson, I.; van Meeteren, L.A.; Samen, E.; Lu, L.; Vanwildemeersch, M.; et al. Vascular endothelial growth factor B controls endothelial fatty acid uptake. *Nature* **2010**, *464*, 917–921. [[CrossRef](#)] [[PubMed](#)]
114. Yang, L.; Broderick, D.; Campbell, Y.; Gombart, A.F.; Stevens, J.F.; Jiang, Y.; Hsu, V.L.; Bisson, W.H.; Maier, C.S. Conformational modulation of the farnesoid X receptor by prenylflavonoids: Insights from hydrogen deuterium exchange mass spectrometry (HDX-MS), fluorescence titration and molecular docking studies. *Biochim. Biophys. Acta (BBA)-Proteins Proteom.* **2016**, *1864*, 1667–1677. [[CrossRef](#)]
115. Klip, A. The many ways to regulate glucose transporter 4. *Appl. Physiol. Nutr. Metab.* **2009**, *34*, 481–487. [[CrossRef](#)]
116. Christoffel, J.; Rimoldi, G.; Wuttke, W. Effects of 8-prenylnaringenin on the hypothalamo-pituitary-uterine axis in rats after 3-month treatment. *J. Endocrinol.* **2006**, *188*, 397–405. [[CrossRef](#)] [[PubMed](#)]
117. Jarry, H.; Sprenger, M.; Wuttke, W. Rates of release of GABA and catecholamines in the mediobasal hypothalamus of ovariectomized and ovariectomized estrogen-treated rats: Correlation with blood prolactin levels. *Neuroendocrinology* **1986**, *44*, 422–428. [[CrossRef](#)]
118. Tataryn, I.V.; Meldrum, D.R.; Lu, K.; Fruraar, A.M.; Judd, H.L. LH, FSH and skin temperature during the menopausal hot flash. *J. Clin. Endocrinol. Metab.* **1979**, *49*, 152–154. [[CrossRef](#)] [[PubMed](#)]
119. Gambone, J.; Meldrum, D.; Laufer, L.; Chang, R.; Lu, J.; Judd, H. Further delineation of hypothalamic dysfunction responsible for menopausal hot flashes. *J. Clin. Endocrinol. Metab.* **1984**, *59*, 1097–1102. [[CrossRef](#)]
120. Heyerick, A.; Vervarcke, S.; Depypere, H.; Bracke, M.; De Keukeleire, D. A first prospective, randomized, double-blind, placebo-controlled study on the use of a standardized hop extract to alleviate menopausal discomforts. *Maturitas* **2006**, *54*, 164–175. [[CrossRef](#)]
121. Stefaneanu, L.; Kovacs, K.; Horvath, E.; Lloyd, R.; Buchfelder, M.; Fahlbusch, R.; Smyth, H. In situ hybridization study of estrogen receptor messenger ribonucleic acid in human adenohypophysial cells and pituitary adenomas. *J. Clin. Endocrinol. Metab.* **1994**, *78*, 83–88.

122. Szczepankiewicz, D.; Skrzypski, M.; Pruszyńska-Oszmalek, E.; Zimmermann, D.; Andralojc, K.; Kaczmarek, P.; Wojciechowicz, T.; Sassek, M.; Nowak, K. Importance of ghrelin in hypothalamus—Pituitary axis on growth hormone release during normal pregnancy in the rat. *J. Physiol. Pharmacol.* **2010**, *61*, 443.
123. Ho, K.K.; O'Sullivan, A.J.; Weissberger, A.J.; Kelly, J.J. Sex steroid regulation of growth hormone secretion and action. *Horm. Res. Paediatr.* **1996**, *45*, 67–73. [[CrossRef](#)] [[PubMed](#)]
124. Rupa, R.; Voellger, B.; Kirches, E.; Mawrin, C.; Firsching, R. 8-Prenylnaringenin decreases hormone expression in GH3 pituitary adenoma cells of the rat. In Proceedings of the 15th European Congress of Endocrinology, Copenhagen, Denmark, 27 April–1 May 2013; Volume 32, p. 839.
125. Bottner, M.; Christoffel, J.; Wuttke, W. Effects of long-term treatment with 8-prenylnaringenin and oral estradiol on the GH-IGF-1 axis and lipid metabolism in rats. *J. Endocrinol.* **2008**, *198*, 395–402. [[CrossRef](#)] [[PubMed](#)]
126. Ming, L.; Ge, B.; Wang, M.; Chen, K. Comparison between 8-prenylnaringenin and naringenin concerning their activities on promotion of rat bone marrow stromal cells' osteogenic differentiation in vitro. *Cell Prolif.* **2012**, *45*, 508–515. [[CrossRef](#)]
127. Ming, L.-G.; Lv, X.; Ma, X.-N.; Ge, B.-F.; Zhen, P.; Song, P.; Zhou, J.; Ma, H.-P.; Xian, C.J.; Chen, K.-M.; et al. The prenyl group contributes to activities of phytoestrogen 8-prenylnaringenin in enhancing bone formation and inhibiting bone resorption in vitro. *Endocrinology* **2013**, *154*, 1202–1214. [[CrossRef](#)] [[PubMed](#)]
128. Luo, D.; Kang, L.; Ma, Y.; Chen, H.; Kuang, H.; Huang, Q.; He, M.; Peng, W. Effects and mechanisms of 8-prenylnaringenin on osteoblast MC 3T3-E1 and osteoclast-like cells RAW 264.7. *Food Sci. Nutr.* **2014**, *2*, 341–350. [[CrossRef](#)] [[PubMed](#)]
129. Hümpel, M.; Isaksson, P.; Schaefer, O.; Kaufmann, U.; Ciana, P.; Maggi, A.; Schleuning, W.-D. Tissue specificity of 8-prenylnaringenin: Protection from ovariectomy induced bone loss with minimal trophic effects on the uterus. *J. Steroid Biochem. Mol. Biol.* **2005**, *97*, 299–305. [[CrossRef](#)] [[PubMed](#)]
130. Sehmisch, S.; Hammer, F.; Christoffel, J.; Seidlova-Wuttke, D.; Tezval, M.; Wuttke, W.; Stuermer, K.M.; Stuermer, E.K. Comparison of the phytohormones genistein, resveratrol and 8-prenylnaringenin as agents for preventing osteoporosis. *Planta Med.* **2008**, *74*, 794–801. [[CrossRef](#)]
131. Hoffmann, D.B.; Griesel, M.H.; Brockhusen, B.; Tezval, M.; Komrakova, M.; Menger, B.; Wassmann, M.; Stuermer, K.M.; Sehmisch, S. Effects of 8-prenylnaringenin and whole-body vibration therapy on a rat model of osteopenia. *J. Nutr. Metab.* **2016**, *2016*, 6893137. [[CrossRef](#)]
132. Pedrera-Zamorano, J.D.; Lavado-Garcia, J.M.; Roncero-Martin, R.; Calderon-Garcia, J.F.; Rodriguez-Dominguez, T.; Canal-Macias, M.L. Effect of beer drinking on ultrasound bone mass in women. *Nutrition* **2009**, *25*, 1057–1063. [[CrossRef](#)]
133. Delmulle, L.; Bellahcene, A.; Dhooge, W.; Comhaire, F.; Roelens, F.; Huvaere, K.; Heyerick, A.; Castronovo, V.; De Keukeleire, D. Anti-proliferative properties of prenylated flavonoids from hops (*Humulus lupulus* L.) in human prostate cancer cell lines. *Phytomedicine* **2006**, *13*, 732–734. [[CrossRef](#)]
134. Delmulle, L.; Berghe, T.V.; Keukeleire, D.D.; Vandenabeele, P. Treatment of PC-3 and DU145 prostate cancer cells by prenylflavonoids from hop (*Humulus lupulus* L.) induces a caspase-independent form of cell death. *Phytother. Res. Int. J. Devoted Pharmacol. Toxicol. Eval. Nat. Prod. Deriv.* **2008**, *22*, 197–203. [[CrossRef](#)] [[PubMed](#)]
135. Busch, C.; Noor, S.; Leischner, C.; Burkard, M.; Lauer, U.M.; Venturelli, S. Anti-proliferative activity of hop-derived prenylflavonoids against human cancer cell lines. *Wien. Med. Wochenschr.* **2015**, *165*, 258–261. [[CrossRef](#)] [[PubMed](#)]
136. Gąsiorowska, J.; Teisseyre, A.; Uryga, A.; Michalak, K. The influence of 8-prenylnaringenin on the activity of voltage-gated Kv1.3 potassium channels in human Jurkat T cells. *Cell. Mol. Biol. Lett.* **2012**, *17*, 559–570. [[CrossRef](#)]
137. Wang, Y.; Chen, Y.; Wang, J.; Chen, J.; Aggarwal, B.; Pang, X.; Liu, M. Xanthohumol, a prenylated chalcone derived from hops, suppresses cancer cell invasion through inhibiting the expression of CXCR4 chemokine receptor. *Curr. Mol. Med.* **2012**, *12*, 153–162. [[CrossRef](#)]
138. Monteiro, R.; Faria, A.; Azevedo, I.; Calhau, C. Modulation of breast cancer cell survival by aromatase inhibiting hop (*Humulus lupulus* L.) flavonoids. *J. Steroid Biochem. Mol. Biol.* **2007**, *105*, 124–130. [[CrossRef](#)]
139. Diller, R.A.; Riepl, H.M.; Rose, O.; Frias, C.; Henze, G.; Prokop, A. Ability of prenylflavanones present in hops to induce apoptosis in a human Burkitt lymphoma cell line. *Planta Med.* **2007**, *73*, 755–761. [[CrossRef](#)] [[PubMed](#)]
140. Venturelli, S.; Niessner, H.; Sinnberg, T.; Berger, A.; Burkard, M.; Urmann, C.; Donaubaue, K.; Böcker, A.; Leischner, C.; Riepl, H.; et al. 6- and 8-Prenylnaringenin, novel natural histone deacetylase inhibitors found in hops, exert antitumor activity on melanoma cells. *Cell. Physiol. Biochem.* **2018**, *51*, 543–556. [[CrossRef](#)]
141. Zhang, Y.; Zhou, D.; Liu, W.; Li, C.; Hao, L.; Zhang, G.; Deng, S.; Yang, R.; Qin, J.; Li, J.; et al. Cytotoxic Activity and Related Mechanisms of Prenylflavonoids Isolated from *Mallotus conspurcatus* Croizat. *Chem. Biodivers.* **2019**, *16*, e1800465. [[CrossRef](#)]
142. Koosha, S.; Mohamed, Z.; Sinniah, A.; Ibrahim, Z.A.; Seyedan, A.; Alshawsh, M.A. Antiproliferative and apoptotic activities of 8-prenylnaringenin against human colon cancer cells. *Life Sci.* **2019**, *232*, 116633. [[CrossRef](#)]
143. Stompor, M.; Uram, Ł.; Podgórski, R. In vitro effect of 8-prenylnaringenin and naringenin on fibroblasts and glioblastoma cells—cellular accumulation and cytotoxicity. *Molecules* **2017**, *22*, 1092. [[CrossRef](#)]
144. Rong, H.; Boterberg, T.; Maubach, J.; Stove, C.; Depypere, H.; Van Slambrouck, S.; Serreyn, R.; De Keukeleire, D.; Mareel, M.; Bracke, M.; et al. 8-Prenylnaringenin, the phytoestrogen in hops and beer, upregulates the function of the E-cadherin/catenin complex in human mammary carcinoma cells. *Eur. J. Cell Biol.* **2001**, *80*, 580–585. [[CrossRef](#)] [[PubMed](#)]
145. Singhai, R.; Patil, V.W.; Jaiswal, S.R.; Patil, S.D.; Tayade, M.B.; Patil, A.V. E-Cadherin as a diagnostic biomarker in breast cancer. *N. Am. J. Med. Sci.* **2011**, *3*, 227. [[CrossRef](#)]

146. Effenberger, K.E.; Johnsen, S.A.; Monroe, D.G.; Spelsberg, T.C.; Westendorf, J.J. Regulation of osteoblastic phenotype and gene expression by hop-derived phytoestrogens. *J. Steroid Biochem. Mol. Biol.* **2005**, *96*, 387–399. [[CrossRef](#)] [[PubMed](#)]
147. van Duursen, M.B.; Smeets, E.E.; Rijk, J.C.; Nijmeijer, S.M.; van den Berg, M. Phytoestrogens in menopausal supplements induce ER-dependent cell proliferation and overcome breast cancer treatment in an in vitro breast cancer model. *Toxicol. Appl. Pharmacol.* **2013**, *269*, 132–140. [[CrossRef](#)] [[PubMed](#)]
148. Paoletti, T.; Fallarini, S.; Gugliesi, F.; Minassi, A.; Appendino, G.; Lombardi, G. Anti-inflammatory and vascularprotective properties of 8-prenylapigenin. *Eur. J. Pharmacol.* **2009**, *620*, 120–130. [[CrossRef](#)]
149. Tokalov, S.V.; Henker, Y.; Schwab, P.; Metz, P.; Gutzeit, H.O. Toxicity and cell cycle effects of synthetic 8-prenylnaringenin and derivatives in human cells. *Pharmacology* **2004**, *71*, 46–56. [[CrossRef](#)]
150. Negrao, R.; Costa, R.; Duarte, D.; Taveira Gomes, T.; Mendanha, M.; Moura, L.; Vasques, L.; Azevedo, I.; Soares, R. Angiogenesis and inflammation signaling are targets of beer polyphenols on vascular cells. *J. Cell. Biochem.* **2010**, *111*, 1270–1279. [[CrossRef](#)]
151. Adeoya-Osiguwa, S.; Markoulaki, S.; Pocock, V.; Milligan, S.; Fraser, L. 17 $\beta$ -Estradiol and environmental estrogens significantly affect mammalian sperm function. *Hum. Reprod.* **2003**, *18*, 100–107. [[CrossRef](#)]
152. Fraser, L.R.; Beyret, E.; Milligan, S.R.; Adeoya-Osiguwa, S.A. Effects of estrogenic xenobiotics on human and mouse spermatozoa. *Hum. Reprod.* **2006**, *21*, 1184–1193. [[CrossRef](#)]
153. Izzo, G.; Söder, O.; Svechnikov, K. The prenylflavonoid phytoestrogens 8-prenylnaringenin and isoxanthohumol differentially suppress steroidogenesis in rat Leydig cells in ontogenesis. *J. Appl. Toxicol.* **2011**, *31*, 589–594. [[CrossRef](#)]
154. Solak, K.A.; Santos, R.R.; van den Berg, M.; Blaauboer, B.J.; Roelen, B.A.; van Duursen, M.B. Naringenin (NAR) and 8-prenylnaringenin (8-PN) reduce the developmental competence of porcine oocytes in vitro. *Reprod. Toxicol.* **2014**, *49*, 1–11. [[CrossRef](#)] [[PubMed](#)]
155. Solak, K.A.; Wijnolts, F.M.; Nijmeijer, S.M.; Blaauboer, B.J.; van den Berg, M.; van Duursen, M.B. Excessive levels of diverse phytoestrogens can modulate steroidogenesis and cell migration of KGN human granulosa-derived tumor cells. *Toxicol. Rep.* **2014**, *1*, 360–372. [[CrossRef](#)] [[PubMed](#)]
156. Simpson, E.R.; Clyne, C.; Rubin, G.; Boon, W.C.; Robertson, K.; Britt, K.; Speed, C.; Jones, M. Aromatase—A brief overview. *Annu. Rev. Physiol.* **2002**, *64*, 93–127. [[CrossRef](#)]
157. Monteiro, R.; Becker, H.; Azevedo, I.; Calhau, C. Effect of hop (*Humulus lupulus* L.) flavonoids on aromatase (estrogen synthase) activity. *J. Agric. Food Chem.* **2006**, *54*, 2938–2943. [[CrossRef](#)] [[PubMed](#)]
158. Corso, A.; Cappiello, M.; Mura, U. From a dull enzyme to something else: Facts and perspectives regarding aldose reductase. *Curr. Med. Chem.* **2008**, *15*, 1452–1461. [[CrossRef](#)]
159. Endo, S.; Matsunaga, T.; Nishinaka, T. The Role of AKR1B10 in Physiology and Pathophysiology. *Metabolites* **2021**, *11*, 332. [[CrossRef](#)]
160. Seliger, J.M.; Misuri, L.; Maser, E.; Hintzpeter, J. The hop-derived compounds xanthohumol, isoxanthohumol and 8-prenylnaringenin are tight-binding inhibitors of human aldo-keto reductases 1B1 and 1B10. *J. Enzym. Inhib. Med. Chem.* **2018**, *33*, 607–614. [[CrossRef](#)] [[PubMed](#)]
161. Seliger, J.M.; Martin, H.-J.; Maser, E.; Hintzpeter, J. Potent inhibition of human carbonyl reductase 1 (CBR1) by the prenylated chalconoid xanthohumol and its related prenylflavonoids isoxanthohumol and 8-prenylnaringenin. *Chem. Biol. Interact.* **2019**, *305*, 156–162. [[CrossRef](#)]
162. Orhan, I.E.; Jedrejek, D.; Senol, F.S.; Salmas, R.E.; Durdagi, S.; Kowalska, I.; Pecio, L.; Oleszek, W. Molecular modeling and in vitro approaches towards cholinesterase inhibitory effect of some natural xanthohumol, naringenin, and acyl phloroglucinol derivatives. *Phytomedicine* **2018**, *42*, 25–33. [[CrossRef](#)]
163. Pepper, M.S.; Hazel, S.J.; Hümpel, M.; Schleuning, W.D. 8-prenylnaringenin, a novel phytoestrogen, inhibits angiogenesis in vitro and in vivo. *J. Cell. Physiol.* **2004**, *199*, 98–107. [[CrossRef](#)]
164. Costa, R.; Rodrigues, I.; Guardao, L.; Lima, J.Q.; Sousa, E.; Soares, R.; Negrao, R. Modulation of VEGF signaling in a mouse model of diabetes by xanthohumol and 8-prenylnaringenin: Unveiling the angiogenic paradox and metabolism interplay. *Mol. Nutr. Food Res.* **2017**, *61*, 1600488. [[CrossRef](#)] [[PubMed](#)]
165. Fetalvero, K.M.; Martin, K.A.; Hwa, J. Cardioprotective prostacyclin signaling in vascular smooth muscle. *Prostaglandins Other Lipid Mediat.* **2007**, *82*, 109–118. [[CrossRef](#)]
166. Cho, Y.-C.; You, S.-K.; Kim, H.J.; Cho, C.-W.; Lee, I.-S.; Kang, B.Y. Xanthohumol inhibits IL-12 production and reduces chronic allergic contact dermatitis. *Int. Immunopharmacol.* **2010**, *10*, 556–561. [[CrossRef](#)] [[PubMed](#)]
167. Deiters, B.; Prehm, P. Inhibition of hyaluronan export reduces collagen degradation in interleukin-1 treated cartilage. *Arthritis Res. Ther.* **2008**, *10*, 1–9. [[CrossRef](#)]
168. Stracke, D.; Schulz, T.; Prehm, P. Inhibitors of hyaluronan export from hops prevent osteoarthritic reactions. *Mol. Nutr. Food Res.* **2011**, *55*, 485–494. [[CrossRef](#)]
169. Dixon, R.A. Phytoestrogens. *Annu. Rev. Plant Biol.* **2004**, *55*, 225–261. [[CrossRef](#)]
170. Di Vito, C.; Bertoni, A.; Nalin, M.; Sampietro, S.; Zanfa, M.; Sinigaglia, F. The phytoestrogen 8-prenylnaringenin inhibits agonist-dependent activation of human platelets. *Biochim. Biophys. Acta (BBA)-Gen. Subj.* **2012**, *1820*, 1724–1733. [[CrossRef](#)]
171. van Breemen, R.B.; Yuan, Y.; Banuvar, S.; Shulman, L.P.; Qiu, X.; Alvarenga, R.F.R.; Chen, S.N.; Dietz, B.M.; Bolton, J.L.; Pauli, G.F.; et al. Pharmacokinetics of prenylated hop phenols in women following oral administration of a standardized extract of hops. *Mol. Nutr. Food Res.* **2014**, *58*, 1962–1969. [[CrossRef](#)]

172. Mukai, R.; Horikawa, H.; Lin, P.-Y.; Tsukumo, N.; Nikawa, T.; Kawamura, T.; Nemoto, H.; Terao, J. 8-Prenylaringenin promotes recovery from immobilization-induced disuse muscle atrophy through activation of the Akt phosphorylation pathway in mice. *Am. J. Physiol. -Regul. Integr. Comp. Physiol.* **2016**, *311*, R1022–R1031. [[CrossRef](#)] [[PubMed](#)]
173. Mukai, R.; Horikawa, H.; Fujikura, Y.; Kawamura, T.; Nemoto, H.; Nikawa, T.; Terao, J. Prevention of disuse muscle atrophy by dietary ingestion of 8-prenylaringenin in denervated mice. *PLoS ONE* **2012**, *7*, e45048. [[CrossRef](#)]
174. Oberbauer, E.; Urmann, C.; Steffenhagen, C.; Bieler, L.; Brunner, D.; Furtner, T.; Humpel, C.; Bäumer, B.; Bandtlow, C.; Couillard-Despres, S.; et al. Chroman-like cyclic prenylflavonoids promote neuronal differentiation and neurite outgrowth and are neuroprotective. *J. Nutr. Biochem.* **2013**, *24*, 1953–1962. [[CrossRef](#)]
175. Urmann, C.; Oberbauer, E.; Couillard-Després, S.; Aigner, L.; Riepl, H. Neurodifferentiating potential of 8-prenylaringenin and related compounds in neural precursor cells and correlation with estrogen-like activity. *Planta Med.* **2015**, *81*, 305–311. [[CrossRef](#)] [[PubMed](#)]
176. Monteiro, A.F.M.; Viana, J.D.O.; Nayariseri, A.; Zondegomba, E.N.; Mendonça Junior, F.J.B.; Scotti, M.T.; Scotti, L. Computational studies applied to flavonoids against Alzheimer's and Parkinson's diseases. *Oxidative Med. Cell. Longev.* **2018**, *2018*, 912765. [[CrossRef](#)]
177. Kim, J.H.; Cho, I.S.; So, Y.K.; Kim, H.-H.; Kim, Y.H. Kushenol A and 8-prenylkaempferol, tyrosinase inhibitors, derived from *Sophora flavescens*. *J. Enzym. Inhib. Med. Chem.* **2018**, *33*, 1048–1054. [[CrossRef](#)]
178. Benkherouf, A.Y.; Soini, S.L.; Stompor, M.; Uusi-Oukari, M. Positive allosteric modulation of native and recombinant GABAA receptors by hops prenylflavonoids. *Eur. J. Pharmacol.* **2019**, *852*, 34–41. [[CrossRef](#)] [[PubMed](#)]
179. Bagatin, M.C.; Tozatti, C.S.S.; Abiko, L.A.; dos Santos Yamazaki, D.A.; Silva, P.R.A.; Perego, L.M.; Audi, E.A.; Seixas, F.A.V.; Basso, E.A.; de Freitas Gauze, G.; et al. Molecular Docking and Panicolytic Effect of 8-Prenylaringenin in the Elevated T-Maze. *Chem. Pharm. Bull.* **2014**, *62*, 1231–1237. [[CrossRef](#)]
180. Luescher, S.; Urmann, C.; Butterweck, V. Effect of hops derived prenylated phenols on TNF- $\alpha$  induced barrier dysfunction in intestinal epithelial cells. *J. Nat. Prod.* **2017**, *80*, 925–931. [[CrossRef](#)] [[PubMed](#)]
181. Baum, B.; Georgiou, M. Dynamics of adherens junctions in epithelial establishment, maintenance, and remodeling. *J. Cell Biol.* **2011**, *192*, 907–917. [[CrossRef](#)] [[PubMed](#)]
182. De Ryck, T.; Van Impe, A.; Vanhoecke, B.W.; Heyerick, A.; Vakaet, L.; De Neve, W.; Müller, D.; Schmidt, M.; Dörr, W.; Bracke, M.E.; et al. 8-prenylaringenin and tamoxifen inhibit the shedding of irradiated epithelial cells and increase the latency period of radiation-induced oral mucositis. *Strahlenther. Und Onkol.* **2015**, *191*, 429–436. [[CrossRef](#)] [[PubMed](#)]
183. Bartmańska, A.; Walecka-Zacharska, E.; Tronina, T.; Popłoński, J.; Sordon, S.; Brzezowska, E.; Bania, J.; Huszcza, E. Antimicrobial properties of spent hops extracts, flavonoids isolated therefrom, and their derivatives. *Molecules* **2018**, *23*, 2059. [[CrossRef](#)] [[PubMed](#)]
184. Ng, K.R.; Lyu, X.; Mark, R.; Chen, W.N. Antimicrobial and antioxidant activities of phenolic metabolites from flavonoid-producing yeast: Potential as natural food preservatives. *Food Chem.* **2019**, *270*, 123–129. [[CrossRef](#)] [[PubMed](#)]
185. Omar, R.M.; Igoli, J.; Gray, A.I.; Ebiloma, G.U.; Clements, C.; Fearnley, J.; Edrada Ebel, R.A.; Zhang, T.; De Koning, H.P.; Watson, D.G.; et al. Chemical characterisation of Nigerian red propolis and its biological activity against *Trypanosoma brucei*. *Phytochem. Anal.* **2016**, *27*, 107–115. [[CrossRef](#)] [[PubMed](#)]
186. Miranda, C.L.; Stevens, J.F.; Ivanov, V.; McCall, M.; Frei, B.; Deinzer, M.L.; Buhler, D.R. Antioxidant and prooxidant actions of prenylated and nonprenylated chalcones and flavanones in vitro. *J. Agric. Food Chem.* **2000**, *48*, 3876–3884. [[CrossRef](#)] [[PubMed](#)]
187. Luís, C.; Costa, R.; Rodrigues, I.; Castela, Â.; Coelho, P.; Guerreiro, S.; Gomes, J.; Reis, C.; Soares, R. Xanthohumol and 8-prenylaringenin reduce type 2 diabetes-associated oxidative stress by downregulating galectin-3. *Porto Biomed. J.* **2019**, *4*, e23. [[CrossRef](#)] [[PubMed](#)]
188. Nowotny, K.; Jung, T.; Höhn, A.; Weber, D.; Grune, T. Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules* **2015**, *5*, 194–222. [[CrossRef](#)]
189. Koch, K.; Schulz, G.; Döring, W.; Büchter, C.; Havermann, S.; Mutiso, P.C.; Passreiter, C.; Wätjen, W. Abyssinone V, a prenylated flavonoid isolated from the stem bark of *Erythrina melanacantha* increases oxidative stress and decreases stress resistance in *Caenorhabditis elegans*. *J. Pharm. Pharmacol.* **2019**, *71*, 1007–1016. [[CrossRef](#)] [[PubMed](#)]
190. Gross-Steinmeyer, K.; Stapleton, P.L.; Tracy, J.H.; Bammler, T.K.; Strom, S.C.; Buhler, D.R.; Eaton, D.L. Modulation of aflatoxin B1-mediated genotoxicity in primary cultures of human hepatocytes by diindolylmethane, curcumin, and xanthohumols. *Toxicol. Sci.* **2009**, *112*, 303–310. [[CrossRef](#)]