



UNIVERSITY OF HELSINKI

<https://helda.helsinki.fi>

Clinical Studies on Drug-Drug Interactions Involving Metabolism and Transport : Methodology, Pitfalls, and Interpretation

Tornio, Aleks; Filppula, Anne M.; Niemi, Mikko; Backman, Janne T.

2019-06

Wiley Blackwell

<http://hdl.handle.net/10138/302836>

Tornio, A, Filppula, A M, Niemi, M & Backman, J T 2019, 'Clinical Studies on Drug-Drug Interactions Involving Metabolism and Transport : Methodology, Pitfalls, and Interpretation', *Clinical Pharmacology and Therapeutics*, vol. 105, no. 6, pp. 1345-1361.
<https://doi.org/10.1002/cpt.1435>

Downloaded from Helda, University of Helsinki institutional repository. <https://helda.helsinki.fi>
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.
Please cite the original version.

Clinical Studies on Drug-Drug Interactions Involving Metabolism and Transport: Methodology, Pitfalls and Interpretation

Aleksi Tornio^{1,2}, Anne M Filppula^{1,2}, Mikko Niemi^{1,2}, Janne T Backman^{1,2}

¹Individualized Drug Therapy Research Program, Faculty of Medicine, University of Helsinki,
Helsinki, Finland

²Department of Clinical Pharmacology, University of Helsinki and Helsinki University Hospital,
Helsinki, Finland

Corresponding author:

Professor Janne T. Backman, MD

Department of Clinical Pharmacology

University of Helsinki and Helsinki University Hospital, PO Box 705, FI-00029 HUS, Finland

Fax: +358-9-471 74039

Tel. +358-50-428 0997

E-mail: janne.backman@helsinki.fi

Conflict of interest: The authors declared no competing interests for this work.

Funding: This work has been financially supported by grants from the Academy of Finland [Grant decision 278123, 2014], European Research Council (ERC) [Grant agreement 725249], Sigrid Juselius Foundation and by State funding for university-level health research, Finland.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/cpt.1435

This article is protected by copyright. All rights reserved.

Keywords: Drug-drug interaction, drug-metabolizing enzymes, drug transporters, pharmacogenetics, pharmacokinetics, physiologically-based pharmacokinetic modeling, study design

ABSTRACT

Many drug-drug interactions (DDIs) are based on alterations of the plasma concentrations of a victim drug due to another drug causing inhibition and/or induction of the metabolism or transporter-mediated disposition of the victim drug. In the worst case, such interactions cause more than tenfold increases or decreases in victim drug exposure, with potentially life-threatening consequences. There has been tremendous progress in the predictability and modeling of DDIs. Accordingly, the combination of modeling approaches and clinical studies is the current mainstay in evaluation of the pharmacokinetic DDI risks of drugs. In this article, we focus on the methodology of clinical studies on DDIs involving drug metabolism or transport. We specifically present considerations related to general DDI study designs, recommended enzyme and transporter index substrates and inhibitors, pharmacogenetic perspectives, index drug cocktails, endogenous substrates, limited sampling strategies, physiologically-based pharmacokinetic modeling, complex DDIs, methodological pitfalls, and interpretation of DDI information.

INTRODUCTION

Unintentional and mismanaged drug-drug interactions (DDIs) are a common reason for preventable adverse events.⁽¹⁾ As the population is aging and polypharmacotherapy is becoming progressively more common, there is an increased likelihood of DDIs that can inadvertently lead to exaggeration of adverse effects or – in some cases – loss of drug efficacy. As these kind of events cannot be prevented without recognizing the need to adjust medications according to DDI risks, there is a need for carefully planned preclinical and clinical DDI studies during drug development, and typically also after marketing approval, as well as for modeling studies, databases and clinical decision support systems that can be easily implemented and used to improve clinical decision making.

In the past, extreme safety concerns caused by DDIs have led to multiple market withdrawals, such as those of mibefradil, terfenadine, cisapride and cerivastatin in the late 1990's and early 2000's. Due to such unfortunate incidents and the rapid accumulation of scientific knowledge that has improved the understanding of DDI mechanisms and awareness of DDI risks, regulatory agencies have frequently updated their guidances on drug interaction studies. For example, the last clinical drug interaction studies guidance by the U.S. Food and Drug Administration (FDA) was published in 2017 and that by the European Medicines Agency (EMA) is currently being revised.(2, 3) Even though these guidelines are directed for studies performed for drugs under development, their concepts can be applied to drugs on the market as well.

The above developments have led to marked advances in the conduct of DDI studies during drug development. As a result, the number of drug withdrawals due to DDIs has dramatically decreased, and detailed knowledge on mechanisms, clinical relevance and management of DDIs mediated by inhibition or induction of cytochrome P450 enzymes (CYPs), some other enzymes and key transporters are in most cases available already at the time of marketing approval. For example, among the 34 drugs approved by the FDA in 2017, five had been identified as sensitive substrates of CYP3A or organic anion transporting polypeptide (OATP) 1B1, and three had been considered as strong inhibitors of CYP3A, OATP1B1, or BCRP, while no strong inducers had been identified.(4)

A major proportion of harmful drug interactions is based on alterations of the plasma concentrations of the victim drug due to the perpetrator drug causing a change in the metabolism or transporter-mediated disposition of the victim drug. While inhibition of drug metabolism or transporter-dependent elimination leads in most cases to elevated concentrations of the victim drug, induction increases metabolic elimination, decreasing the concentrations of the victim. In the worst case, such interactions can lead to several hundred-fold variations in drug exposure.(5, 6) During the past decade, several review articles have been published focusing on various specific aspects related to clinical DDI studies.(7-10) In this article, we present an overview of the basic methodology of clinical DDI studies that can be used when investigating a specific drug as a victim or perpetrator of pharmacokinetic DDIs mediated by inhibition or induction of drug-metabolizing enzymes and/or

transporters, with an attempt to pinpoint specific considerations that we have found important on the basis of our own experience in clinical DDI studies. While much of the methodology described in regulatory guidance is focused on studies carried out during drug development, we extend the review beyond the regulatory guidance, highlighting certain specific questions related to complex DDIs, pharmacogenetics, methodological pitfalls, and interpretation of DDI information.

DESIGN OF CLINICAL DDI STUDIES

General considerations of clinical DDI study design

With regard to pharmacokinetic DDIs, the study hypotheses and objectives essentially define the most suitable study design. Based on prior information, an evaluation of interaction risk is usually carried out continuously during drug development (**Figure 1**). When the drug enters clinical development, this evaluation is typically based on *in vitro*, animal, and *in silico* data, but the evaluation is rapidly strengthened by accumulating clinical data. If the drug is in later stages of clinical development or already on the market, DDI studies may be triggered by retrospective analyses of population data, case reports or additional *in vitro* data. Consequently, the drug of interest can be evaluated either as a substrate (victim drug), or as an inhibitor or inducer (perpetrator drug) of drug-metabolizing enzymes or transporters. For example, a retrospective epidemiologic analysis of the risk of rhabdomyolysis in cerivastatin users with concomitant clopidogrel published in 2012 (11) triggered an investigation that resulted in identification of clopidogrel as an inhibitor of CYP2C8.(12)

There are multiple design issues and requirements that need to be carefully considered already when preparing a clinical DDI study protocol (**Table 1**). The golden standard for clinical DDI studies is a prospective cross-over study, usually performed in healthy volunteers. The cross-over design reduces the effect of interindividual variability because individuals act as their own controls. Furthermore, inter-individual variability is reduced in healthy subjects, who have normal kidney and liver function, are not using concomitant medication and are devoid of other interfering factors (e.g.,

smoking). In a two-way pharmacokinetic cross-over DDI study in healthy volunteers, sample sizes as low as 10-12 individuals are often sufficient to demonstrate clinically significant interaction, although a 2-3 times larger sample may be needed to demonstrate lack of DDI. In a cross-over study, the wash-out period between study phases should be long enough to allow the drugs, metabolites and their effects to be completely eliminated before the next phase, even when their elimination is impaired by strong inhibitors. When drug toxicity or adverse effects are an issue, DDI studies can be performed in patient populations, but they typically require a larger sample size due to greater variability caused by various individual factors, particularly if a parallel group design is used.

If inhibition or induction caused by a drug is investigated, a probe or index substrate for a specific pharmacokinetic pathway is typically used as the victim drug (**Table 2**). Particularly in drug development, such studies are first focused on the enzyme towards which the investigated drug is expected to cause the strongest inhibition on the basis of *in vitro* inhibition data. If the DDI potential of a drug as an inhibitor is being studied, it should be administered preferably with multiple doses reaching and maintaining concentrations approximating steady-state at the highest clinically used doses, so that the study could reveal the worst-case scenario (**Figure 2**). In addition, our recommendation is to use dose staggering, rather than simultaneous administration, to allow maximum inhibitor concentrations at the site of inhibition (for example, an interval of one hour between the inhibitor and victim drug administration) to reveal the full extent of interaction, particularly if the absorption of the inhibitor can be delayed or if the victim is rapidly eliminated and has extensive first-pass metabolism. On the other hand, to evaluate induction effects, a sufficient multiple dosing period to reach near full induction should be used. However, when both induction and inhibition are possible, a delay between perpetrator and victim drug administration of e.g., 12 hours can be considered to ensure full induction effect of the perpetrator. For example, the prototypical CYP inducer rifampin (13) inhibits OATP1B1/1B3 immediately after dosing *in vivo* (14), but this effect is no longer present after 12 hours.(15)

When the drug is investigated as a victim, probe or index inhibitors can be utilized to reveal the contributions of specific pathways to the drug's pharmacokinetics, i.e., to document the clinical relevance of the key enzymes and transporters involved (**Table 3**). In this situation, a low single dose of the victim drug is usually sufficient. If a strong interaction is expected, even subtherapeutic victim drug doses may be necessary for ethical reasons, to ensure a large enough safety margin in drug exposure. Such a dosing scheme is adequately informative provided that the results can be extrapolated to steady-state conditions at clinically relevant doses, i.e., the victim drug displays linear dose- and time-dependent pharmacokinetics. On the other hand, when the effect of a strong enzyme/transporter inducer is investigated, a relatively high victim drug dose may be necessary to allow quantification of its concentrations. Monitoring of pharmacodynamic effects may be required for safety and can be useful when evaluating the clinical significance of the DDI. Pharmacodynamic endpoints are particularly useful in cases where the victim drug has active metabolites, where organ specific disposition of the drug is anticipated to be altered, e.g., entry to central nervous system via blood-brain barrier, or where concomitant administration of the perpetrator and victim is anticipated to occur in clinical practice. If subjective pharmacodynamic measurements are used, double-blinding of the treatment phases is crucial.

Properties of suitable index substrates of drug-metabolizing enzymes

The most important property of an index substrate for DDI studies is sensitivity for the specific pharmacokinetic pathway investigated (**Table 2**). In the context of drug-metabolizing enzymes, the fraction metabolized by the enzyme should optimally be >80%, so that an over 5-fold increase in the area under plasma concentration-time curve (AUC) can occur when the enzyme is completely inhibited (**Figure 2**). Furthermore, other disposition mechanisms of the index substrate should be known in detail, especially in cases where the perpetrator has the potential to cause interactions via other mechanisms. The properties of the substrate should also be suitable for thorough characterization of its pharmacokinetic parameters in a clinical study. Most importantly, the

elimination half-life ($t_{1/2}$) of the probe should be sufficiently short to avoid unnecessarily long sample collection (needed to cover 80-90% of its total AUC) and wash-out period. From this point of view, e.g., efavirenz, warfarin and desipramine are far from ideal index substrates. Particularly if it is important to quantify a transient effect on enzyme activity, an index drug with extensive first-pass metabolism and short $t_{1/2}$ should be preferred. An optimal index substrate exhibits linear pharmacokinetics and has itself no effects on the pharmacokinetics of other drugs. Tolerability and lack of toxicity are also key issues, and the safety margin should be large enough to cover changes of several fold in exposure. Other desirable qualities for an index substrate include ease of quantification and availability of metabolite standards, as well as commercial availability at suitable doses. If equally good substrates for the specific purposes are available, the choice should be made on the basis of the clinical relevance of the potential combinations.

There are several sensitive and selective index substrates for some CYP enzymes, e.g., CYP1A2, CYP2D6 and CYP3A, whereas sensitive index substrates for some CYPs (CYP2A6, CYP2B6, CYP2C9 and CYP2J2) are currently not available (**Table 2**). In some cases, sensitive index substrates are available, but they are also substrates for a transporter protein. For example, more than 80% of repaglinide and simvastatin are metabolized by CYP2C8 and CYP3A, respectively, making them sensitive index substrates of these enzymes, but both are also fairly sensitive substrates of OATP1B1/1B3.⁽¹⁶⁾ The usefulness of both these drugs as index substrates has been documented extensively, and their nonselectivity can thus be taken into account when interpreting the results of the study. For non-CYP enzymes, knowledge is still evolving, and useful probe substrates are known only for few of them (**Table 4**). Lastly, if a sufficiently sensitive index substrate is not otherwise available or feasible, an alternative can be to use a specific reaction, i.e., a metabolic ratio as an index of enzyme activity (see also Limited sampling strategies below).

Properties of suitable index inhibitors and inducers of drug-metabolizing enzymes

In addition to safety, the two most important characteristics of index inhibitors are strength of inhibition and selectivity. The FDA and EMA guidances define a strong index inhibitor as a drug that increases the AUC of a sensitive index substrate of a given metabolic pathway ≥ 5 -fold,(2, 3) which equals an at least 80% inhibition of clearance. Strong inhibition typically requires a repeated dosing of sufficient inhibitor doses for about 5 inhibitor half-lives before the victim drug is administered, followed by inhibitor dosing until victim AUC is fully characterized (**Tables 1 and 3, Figure 2**). Strong inhibition increases the sensitivity to detect interaction effects, especially in cases where the victim drug is not solely dependent on the single elimination pathway. An optimal index inhibitor is selective, i.e., only inhibits one enzyme, so that direct mechanistic conclusions can be drawn from the result with reasonable confidence. On the other hand, if the index inhibitor inhibits several enzymes and/or transporters, the observed effects on victim drug pharmacokinetics cannot be attributed to a single pathway without further studies. Lastly, a long $t_{1/2}$ of the inhibitor makes it impractical in terms of the required wash-out period. For example, the $t_{1/2}$ of 4-6 days of parent fluoxetine and 4-16 days of its metabolite norfluoxetine make fluoxetine a suboptimal index inhibitor of CYP2D6 in cross-over studies. There are fairly good inhibitors for most CYP enzymes, but adequately documented strong and selective inhibitors for CYP2A6, CYP2B6 and CYP2C9 are missing (**Table 3**). For instance, high doses of fluconazole result in strong inhibition of CYP2C9, but they also cause a strong inhibition of CYP3A4 and CYP2C19. Likewise, suitable strong inhibitors for non-CYP enzymes are mainly lacking; for example, the strongest identified UGT-mediated DDIs are limited to a 2-3-fold increase in AUC (**Table 4**). Therefore, also studies with moderate, weak and nonselective inhibitors are needed in particular if concomitant clinical use is likely to occur frequently.

Unlike inhibition, induction of drug-metabolizing enzymes and transporters is usually highly nonselective, with few exceptions, such as induction of CYP1A2 by polycyclic aromatic hydrocarbon (PAH) compounds.(17) Of note, the inducibility of CYP2D6 by known drugs is low or negligible, even though its activity can be increased by certain conditions, such as pregnancy.

Rifampin is the prototypical inducer that is most commonly used in clinical DDI studies because of its safety compared to other strong inducers, such as carbamazepine and phenytoin, and the strength of its inducing effect.(13) Like other compounds causing pregnane X receptor (PXR) mediated induction, rifampin causes a particularly strong induction of intestinal and hepatic CYP3A, but also induces many other drug-metabolizing enzymes and certain transporters. Thus, it can be used to test the sensitivity of a drug to strong enzyme induction. It should be noted that the CYP3A inducing effect of rifampin is so strong that it can dramatically increase the fraction of the drug metabolized by CYP3A, so that even a drug that is normally mainly eliminated via noninducible or weakly inducible pathways becomes dependent on CYP3A4.

Specific considerations for transporter DDI studies

Current evidence indicates that of the numerous transporters, at least breast cancer resistance protein (BCRP), OATP1B1, and P-glycoprotein (P-gp) mediate clinically relevant DDIs. In addition, multidrug and toxin extrusion protein (MATE) 1, MATE2-K, organic anion transporter (OAT) 1, OAT3, OATP1B3, OATP2B1, organic cation transporter (OCT) 1, and OCT2 can be involved in DDIs.(18) Accordingly, drug transporters are routinely investigated already during preclinical drug development, providing a wealth of information regarding the properties of an investigational drug as a substrate and inhibitor of different transporters.(19) With regard to clinical DDI studies, the FDA guidance instructs that the need of clinical transporter DDI studies of transporter substrates identified in *in vitro* studies should be considered on the basis of the drug's site of action, route of elimination, likelihood of concomitant use, and safety issues.(2) For example, in case of P-gp and BCRP-mediated DDIs, most clinically relevant DDIs (aliskiren, dabigatran, digoxin and fexofenadine as P-gp substrates, and rosuvastatin as BCRP substrate) are based on inhibition of the absorption limiting effects of these transporters in the small intestine or in some cases biliary or renal excretion.(20-22) In addition, based on limited evidence, P-gp-mediated DDIs may also occur at blood-tissue barriers, such as the blood-brain-barrier, warranting DDI studies if the site of pharmacological or toxic effects

of a P-gp substrate drug is behind such a barrier. Moreover, for drugs targeting the liver or eliminated via the liver, current guidance states that DDI studies focusing on OATP1B1 or OATP1B3 can be warranted, and for drugs eliminated via renal excretion, studies on OAT1 or 3, OCT2, or MATE-mediated DDIs can be needed.⁽²⁾ Recent studies support the roles of OCT1 and OATP2B1 in mediating hepatic and intestinal DDIs, indicating that they also need to be considered.⁽¹⁹⁾ Furthermore, if the drug is suspected to be a transporter inhibitor on the basis of *in vitro* data, clinical DDI studies could be conducted with likely concomitant substrate drugs that could be subject to a clinically relevant DDI, regardless of the investigational drug's own route of elimination.

DDI studies focusing on transporters have some challenges compared to drug-metabolizing enzymes. Firstly, the lack of specific index substrates and inhibitors makes extrapolation of the results of DDI studies to other drugs challenging (**Table 5, Table S3**). For example, dabigatran etexilate could be used as a fairly specific P-gp substrate because its intestinal absorption is limited by P-gp, but dabigatran elimination is not dependent on renal P-gp and its sensitivity to P-gp inhibition at clinically used doses is only modest, so that even the strongest P-gp inhibitors have increased its AUC less than 3-fold.⁽²³⁾ Similarly, recognized substrates of BCRP (e.g., rosuvastatin), OAT1/3 (e.g., adefovir, ganciclovir and famotidine), and MATE or OCT2 transporters (e.g., dofetilide, metformin) have been at most moderately sensitive to inhibition of the respective transporters.⁽²⁴⁾ In case of OATP1B1 and 1B3, there are some moderately sensitive substrates, such as simvastatin (acid) and pitavastatin, but it is unclear if their AUC could be increased more than 5-fold by inhibition of a single transporter only.

Well-established transporter inhibitors are typically either nonselective or cause only limited extent of inhibition. Although probenecid can be used as a relatively selective inhibitor of OATs 1 and 3, P-gp inhibitors, such as clarithromycin, itraconazole, quinidine and verapamil, also have inhibitory effects on CYP enzymes, and cyclosporine acts as a multi-transporter inhibitor of BCRP, P-gp, and OATPs, and also inhibits CYP3A4. The OATP inhibitor rifampin also acts as a strong inducer of drug-metabolizing enzymes. In case of rifampin, the inducing effect can be largely

avoided by using only a single rifampin dose, but this is primarily a relevant dosing scheme only in mechanistic DDI studies. In addition to the general lack of selectivity of transporter inhibitors, the strength of inhibition of many inhibitors may be limited, as it is unclear if any of the inhibitors have a sufficiently strong inhibitory effect on a single transporter to cause a more than 5-fold increase in the AUC of a victim drug.

Secondly, modulation of transporter activity can result in changes in tissue distribution depending on the localization of the transporter, without significant changes in plasma concentrations. Therefore, solely determining the plasma exposure of the victim drug might not be a valid endpoint for a transporter DDI study. In such cases, it can be useful to measure the pharmacodynamic effect of the victim drug to find out if the perpetrator caused any clinically relevant changes in drug concentration at the site of action. A good example is metformin, which is not metabolized but whose absorption from the intestine, hepatic disposition, and renal elimination are subject to active transport. A DDI study design for metformin has been proposed, where the endpoints consist of systemic pharmacokinetics, renal clearance, and antihyperglycemic effect.(25) Another possibility to study tissue distribution of drugs is provided by nuclear imaging techniques, such as PET and SPECT, where radiolabeled drug of interest is administered and non-invasively measured.(26)

Pharmacogenetics

Our recommendation is to routinely collect DNA samples from participants of DDI studies and to genotype for well-characterized genetic variants of enzymes and transporters involved in pharmacokinetics of investigated drugs. For example, in studies with index inhibitors or substrates of enzymes and transporters that exhibit common genetic variation, genotype information can be important in interpretation of the data, as genotype can markedly influence the extent of the DDI.(27) Genotype information may also be valuable in detecting causes for pharmacokinetic variability or outliers. In some cases, pharmacogenetic studies can also complement or replace DDI studies. They

Accepted Article

can be particularly useful, if no selective or strong inhibitor is available, which is common with drug transporters and non-CYP enzymes. If the enzyme/transporter exhibits considerable genetic variability resulting in altered activity, especially loss-of-function, individuals with different genotypes can be studied in a genotype panel study, to obtain clear mechanistic understanding of the importance of the specific pathway for the pharmacokinetics of the investigational drug. This type of study is only possible in cases where functionally relevant variants are known with suitable population frequencies, allowing recruitment of a sufficient number of subjects. For example, a common single nucleotide variant in *SLCO1B1* (c.521T>C, rs4149056), encoding the hepatic uptake transporter OATP1B1, results in decreased transport activity.(16) Thus, pharmacokinetics in heterozygous and homozygous variant allele carriers can be compared to that in reference allele carriers, to inform about the relevance of OATP1B1 in the disposition of the investigational drug, and its sensitivity to inhibition of OATP1B1. Of drug transporters and drug-metabolizing enzymes, at least *ABCG2*, *CES1*, *CYP2C9*, *CYP2C19*, *CYP2D6*, *CYP3A4*, *CYP3A5*, *DPYD*, *NAT2*, *SLC22A1*, *SLCO1B1*, *TPMT*, *UGT1A1*, *UGT1A3*, *UGT2B10*, and *UGT2B17* show significant genetic variability.(28-30) Variants in these genes could be used to define subject groups in a genotype panel study and genotyping for these genes should be considered if they are likely to be involved in the pharmacokinetics of the victim or perpetrator of a DDI study.

Index drug cocktails and microdosing

In the cocktail approach, more than one index substrate is administered simultaneously to measure activities of several drug-metabolizing enzymes or transporters. In the context of DDI studies, this allows a drug to be studied as a perpetrator for several pharmacokinetic pathways in the same clinical study. A cocktail needs to be validated so that its components do not interact with each other, usually by administering each substrate separately and as a part of the cocktail.(31) Recent advances in this area have occurred in the utilization of very low doses of the substrate drugs (i.e., microdosing) and in development of drug cocktails to assess drug transporter function.(32) The use of microdosing is

beneficial in terms of safety and reduced likelihood of interactions between substrates. A potential problem is that it may be difficult to extrapolate the findings to clinically relevant doses if the substrate displays nonlinear pharmacokinetics.

Several phenotyping cocktails including index substrates for different CYP enzymes have been developed over the last twenty years.(31) Most of the proposed cocktails include index drugs for up to six CYP enzymes that have most commonly been CYP1A2, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and/or CYP3A4. The individual substrates are usually relatively selective for a single enzyme so that mechanistic conclusions can be drawn from the results. One of the challenges related to a few of the earlier cocktails is that some of the suggested index substrates are no longer commercially available in some countries. Two of the more recent cocktails called the Basel and Geneva cocktails, including substrates for CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2D6, and CYP3A4 (Geneva cocktail also including fexofenadine as an index for P-gp), have been specifically validated for DDI studies.(33-35)

One of the biggest challenges in development of drug transporter cocktails is lack of selective index substrates. The first published cocktail approach for transporters included digoxin for P-gp, furosemide for OAT1 and OAT3, metformin for OCT2, MATE1 and MATE2-K, and rosuvastatin for OATP1B1, OATP1B3 and BCRP.(36) The initial study showed mutual interaction resulting in higher rosuvastatin peak plasma concentration (C_{max}) and AUC when administered as a part of the cocktail than individually. In further studies, the doses of furosemide and metformin were reduced, and a cocktail consisting of 0.25 mg digoxin, 1 mg furosemide, 10 mg metformin, and 10 mg rosuvastatin was shown to be devoid of mutual interactions (37, 38). The sensitivity of the cocktail as well the potential for mutual interactions in a situation where the exposure to one of the components is increased due to an inhibitor being studied, remains to be determined. Recently, a transporter cocktail utilizing microdosing was evaluated with known inhibitors in DDI studies (39). The microdose cocktail consisted of 10 μ g midazolam (CYP3A4), 375 μ g dabigatran etexilate (P-gp), 10 μ g pitavastatin (OATP1B), 25 μ g rosuvastatin (BRCP, OATP, and P-gp), and 50 μ g atorvastatin (OATP,

BCRP, P-gp, and CYP3A4). In the DDI studies, the cocktail produced similar results as the index drugs had produced in previous studies, and only dabigatran etexilate showed marked differences to clinical doses with an approximately 2-fold lower dose-normalized oral exposure and higher magnitude of interactions when microdosing was used.

Endogenous substrates

Instead of using index substrate drugs to assess the perpetrator's impact on a specific pharmacokinetic pathway, an endogenous compound as a measure of activity of this pathway could be a useful alternative. The current status of endogenous biomarkers for DDI studies, their limitations, as well as strategies to investigate novel biomarkers for this purpose has been reviewed in detail recently.(24, 40-42) The main advantage of this approach is that it does not require the additional intervention of administering the index substrate. This could be especially useful during early clinical drug development, or when the perpetrator cannot be administered to healthy volunteers due to safety concerns. The endogenous biomarker needs to be validated to fit this purpose, it should be selective for the enzyme or transporter under investigation, and it should not be altered by disease states or dietary factors. Well validated endogenous biomarkers that could function as an alternative to formal DDI studies are, however, still lacking. For example, 4 β -hydroxycholesterol, 6 β -Hydroxycortisol and 6 β -Hydroxycortisone could be used as sensitive biomarkers of CYP3A activity, unconjugated bilirubin as a nonselective marker of UGT1A1 and coproporphyrin as a marker of OATP1B1/1B3, but they are only suitable in initial characterization of DDI risks.(41)

Limited sampling strategies and population pharmacokinetic modeling

In a stand-alone DDI study, the pharmacokinetic profile of the victim drug should be fully characterized with a suitably frequent and long sampling schedule, so that the extrapolated part of the

total AUC is less than 20%.(2) In contrast to studies in healthy volunteers, rich sampling strategy is usually not feasible in large patient populations. However, population pharmacokinetic analyses can be useful to assess the clinical impact of concomitant medications on the pharmacokinetics of the drug of interest despite sparse or limited sampling. The strength of such studies is that DDIs can be investigated in the clinically relevant patient group using typical doses on long-term basis and that the effects of frequently coprescribed drugs can be studied irrespective of prior knowledge on relevant DDI mechanisms. To make the best use of population pharmacokinetic data, the study protocol, including sampling strategy, should be carefully designed before commencing data collection. In particular, accurate documentation of the dosage and timing of drug administration, as well as sampling times, is essential. This is especially important for drugs with variable absorption rate and short $t_{1/2}$. Ideally, also perpetrator drug exposure should be documented by using available samples. The methodology and design of DDI studies using population pharmacokinetic analyses is discussed in a recent commentary by International Society of Pharmacometrics (ISOP) Working Group.(43) For example, a simulation case study with methotrexate and trastuzumab has shown that with careful design, population pharmacokinetic studies can be a viable alternative to DDI studies.(44)

Another instance where limited sampling can be used is with index substrate drugs whose validated metrics of enzyme activity is based on only few or single sample time points, typically by calculating a metabolic ratio from parent drug and its metabolite, whose formation is selective for the enzyme in question. The interpretation of such indices should be done cautiously particularly if the fate of the metabolite is subject to changes by perpetrators. In an ideal case of a fully specific index ratio, a decrease of 80% in a metabolic ratio indicates a maximal increase of 5-fold in the AUC of a sensitive substrate drug. Furthermore, limited sampling can be subject to large variability due to e.g., delayed drug absorption, especially if the metric is based on parent drug concentration only. For example, limited sampling strategies for midazolam and warfarin have shown poor correlation with full pharmacokinetic profiling.(45, 46) On the other hand, determination of a metabolic ratio from a single time-point can be a more useful metric that accurately reflects the systemic clearance of the drug. A good example of such a metric is caffeine, which is selectively

metabolized by CYP1A2 to paraxanthine. The paraxanthine/caffeine ratio from a single plasma sample, taken 4-6 hours after caffeine intake, can be used to assess CYP1A2 activity.(47) Such single time-point metabolic ratios have been tested in index drug cocktails as well.(34, 48) In cocktail approaches, however, a single sampling time represents a compromise between the different substrates and might not be optimal for each index substrate.

One approach to limit blood sample volumes and to make blood sampling more feasible is to use capillary blood microsamples and dried blood as the sample matrix.(34) The main advantage of this approach is its low invasiveness and the flexibility it potentially offers, such as applicability in low resource environments or self-sampling by the study participants. However, the measurement of whole blood concentrations in this matrix needs to be validated against samples collected by venipuncture, and whole blood concentrations of drugs often differ from those in plasma/serum. Use of dried blood samples can be very useful in large-scale studies, such as phase 3 trials or in studies in vulnerable populations, such as children, typically in conjunction with limited sampling strategies, but its usefulness for formal DDI studies in healthy volunteers is limited.

Physiologically-based pharmacokinetic modeling

In the past two decades, major advances have been made in the modeling methods used for prediction of DDIs on the basis of *in vitro* CYP inhibition data.(49, 50) In particular, physiologically-based pharmacokinetic (PBPK) modeling, a dynamic approach incorporating changes in drug concentrations over time, has evolved and is becoming increasingly used even for regulatory decisions.(8, 51, 52) PBPK models simulate concentration-time profiles of drugs in plasma and/or an organ of interest, and allow for simultaneous modeling of multiple drug disposition processes, providing a range of opportunities.(53, 54) The reliability of PBPK models increases with accumulating clinical data and information on the disposition/elimination pathways of the drugs and their metabolites. In the ideal case, the

model can be cross-verified using data from multiple clinical studies concerning both the perpetrator and the victim drug.

PBPK DDI modeling is encouraged by regulatory agencies,(2, 3) and the pharmaceutical industry is increasingly using PBPK modeling to predict DDIs and related dosing decisions as replacement for clinical studies.(8) Examples of submissions where clinical DDI studies have been waived on the basis of PBPK DDI simulations include drugs such as cobimetinib, lesinurad and olaparib.(55) In regulatory submissions, PBPK DDI modeling has been applied to 1) predict DDIs with perpetrator/victim drugs other than those used in clinical studies, 2) predict DDIs in various subpopulations, e.g., specific patient populations or populations with genetically different CYP activities, 3) predict the effects of altered dosing regimens/repeated dosing of perpetrator/victim drug, 4) predict the effects of pH modulation on drug solubility/absorption, and 5) predict the effects of a perpetrator drug on single drug enzymes/transporters.(55) In addition, PBPK DDI modeling has been applied to support the design of clinical DDI studies, and mechanistically explain clinically observed DDIs.

There are many examples of successful PBPK DDI models, in particular in cases where the interaction is based on simple reversible CYP inhibition. In contrast, modeling of the effects of combined time-dependent inhibition and reversible inhibition, or time-dependent inhibition and induction are not as established.(56, 57) Moreover, modeling of DDIs affecting non-CYP drug-metabolizing enzymes and drug transporters may be difficult due to lack of selective *in vivo* data, uncertainties concerning the expression levels and turnover characteristics of transporters and enzymes, as well as scaling of drug concentration levels at the relevant cellular or subcellular sites.(55, 57, 58) A good example highlighting the challenges with modeling of transporter-DDIs, is related to OATP1B1-mediated influx of drugs to hepatocytes. The measured *in vitro* OATP1B1 K_i values of clarithromycin, clopidogrel acyl- β -D glucuronide, cyclosporine and rifampin do not seem to be low enough to recover DDIs observed in clinical studies, which has forced modelers to adjust OATP1B1 K_i values in PBPK models against measured values.(59-61) A possible explanation to this

disconnect from physiological principles is time-dependent inhibition of OATP1B1 by these inhibitors and interplay between drug transporters and drug-metabolizing enzymes.(62) It is obvious that the biochemical and physiological processes related to transporter-mediated DDIs are not fully understood, which poses one of the biggest challenges to development of PBPK modelling and *in vitro* – *in vivo* predictions in general.

Overall, PBPK modeling is a useful tool for DDI research to be used in concert with clinical DDI studies. However, the current limitations of PBPK modeling in complex situations, where multiple mechanisms and DDIs are involved, should be taken into account. On the whole, there has been a lack of consistency in model development and quality assessment practices, demonstrating a need for best-practice guidelines, as well as for harmonization of guidelines between regulatory agencies.(49, 55) At present, however, PBPK models can be highly valuable in interpretation and extrapolation of clinical DDI studies to different clinical situations, as well as in planning and design of further clinical studies.

COMPLEX DDIs

Although DDIs based on a single well-defined pharmacokinetic mechanism can already be predicted and extrapolated to different clinical situations with great confidence, it has become increasingly recognized that pharmacokinetic interaction mechanisms are often complex.(50, 63, 64) Complex DDIs can be difficult to anticipate and control, and can pose a big challenge during early drug development. From the drug development perspective, as well as from the clinical point of view, some of the most problematic complex mechanisms are those involving autoinhibition or induction of the drug's own metabolism, the possibility of several simultaneous mechanisms, inhibition or induction by major drug metabolites and time-dependent inhibition.(64) While comprehensive preclinical data and modeling approaches

fundamental for understanding such DDIs, clinical DDI studies are eventually indispensable to understand their clinical relevance.

For example, on the basis of modern preclinical research methods, autoinduction and autoinhibition can be suspected early, but only the first clinical pharmacokinetic studies reveal if they are clinically relevant. In the worst case, autoinduction can be so strong that adequate drug exposure cannot be obtained despite extreme doses and the drug development program fails. On the other hand, both autoinduction and autoinhibition can result in dose- and time-dependent changes in the relative importance of different disposition pathways and thereby alter the sensitivity of the drug to inhibitory perpetrators. For example, the tyrosine kinase inhibitor imatinib is a mechanism-based inhibitor of CYP3A4 (65) that can increase the plasma concentrations of CYP3A4 substrates, such as simvastatin. As CYP3A4 plays an important role in the metabolism of imatinib, the CYP3A4 inhibitory effect can lead to a complicated increase in the significance of the alternative elimination pathway (CYP2C8), and an increased risk of increased imatinib levels due to CYP2C8 inhibition during multiple imatinib dosing.(66)

An even more complex drug is the kinase inhibitor midostaurin (**Figure 3**), which is a substrate, time-dependent inhibitor and inducer of CYP3A4.(67-69) Based on mechanistic information, it is conceivable that midostaurin could cause a temporary net (auto)inhibition of CYP3A4, which gradually changes to net (auto)induction. On the basis of a short study with midazolam as a victim drug, the effect of midostaurin on CYP3A4 could be interpreted as negligible.(67) However, a DDI study carried out at two early time points may not have revealed a temporary inhibition in the beginning of treatment and leaves unclear if midostaurin acts as an inducer decreasing concentrations of CYP3A4 substrates in long term treatment. In addition, the sensitivity of single-dose midostaurin to CYP3A4

inhibition may not have revealed its full sensitivity to CYP3A4 inhibition at steady-state, i.e., after autoinduction has increased the role of CYP3A4 in its metabolism. Deciphering this kind of dose- and time-dependent complexity requires clinical DDI studies where the drug is the perpetrator, as well as studies where the drug is the victim. Moreover, clinical DDI studies at different time points during the course of treatment are needed, as PBPK models are not reliable in this area, particularly because there is paucity of clinical data on the dose- and time dependency of DDIs involving drugs with nonlinear pharmacokinetics, and because enzyme-specificity of induction is not thoroughly understood.

Examples of DDIs involving multiple mechanisms simultaneously, i.e., DDIs where a multi-pathway substrate is given simultaneously with an inhibitor and/or inducer of the same pathways, include the gemfibrozil-repaglinide (OATP1B1/1B3 and CYP2C8 inhibition), cyclosporine-rosuvastatin (OATP1B and BCRP inhibition), sofosbuvir/velpatasvir/voxilaprevir-rosuvastatin (inhibition of OATP1B1/1B3 and BCRP) and many ritonavir-victim drug (P-gp and CYP3A4 inhibition and CYP induction) interactions.(70-73) The key to building an understanding of this kind of DDI risks is to first learn with enzyme and transporter selective preclinical and clinical studies, which specific enzymes and transporters are involved in the drug's metabolism and disposition and which are inhibited or induced by the drug. Thereafter, relevant DDIs with potential for multiple mechanisms simultaneously can be studied clinically using standard methods. If additional mechanistic understanding is needed, studies with multiple perpetrators or different inhibitor dosing or timing experiments may be useful.

What comes to other types of complex DDIs, those caused by inhibitory or inducing major metabolites and mechanism-based inhibition typically require carefully performed *in vitro* studies to initially identify the risks. The contribution of metabolites as

perpetrators has only been realized during the past decade. It has, for example been estimated that the clinically observed CYP2C9 inhibition by amiodarone, CYP2D6 inhibition by bupropion and CYP3A4 inhibition by sertraline could only be explained accurately when considering their inhibitory metabolites.(74) Of note, even glucuronide metabolites can act as mechanism-based inhibitors of CYP enzymes, capable of causing strong DDIs.(75) Such DDIs can usually be studied using standard clinical DDI study methodology, although in the case of mechanism-based inhibition and in the case of metabolites that have a very long $t_{1/2}$, the time-dependency of the DDI risk may warrant long-term administration or other more focused studies. In addition, the growing number of fixed-dose combination products among newly marketed drugs (especially antiviral agents) represents a unique challenge of understanding the contribution of each drug component to the overall clinical effect when the combination is evaluated as a perpetrator

From a clinical point of view, involvement of multiple perpetrator drugs simultaneously, and the influence of genetic variants or disease states on the risk of DDIs (76) often necessitate specific DDI studies. For example, concomitant treatment with a CYP2C8 inhibitor (e.g., gemfibrozil) and a CYP3A4 inhibitor (e.g., itraconazole) can lead to an extremely strong effect on the pharmacokinetics of drugs that are substrates for both these enzymes, such as repaglinide and loperamide.(71, 77) In case of genetic variants, poor metabolizer individuals are typically insensitive to inhibition of the genetically deficient enzyme, such as inhibition of CYP2C19 by omeprazole or inhibition of CYP2D6 by paroxetine,(78, 79) but they may be particularly susceptible to DDIs mediated by inhibition of an alternative pathway, such as CYP3A4. Genetic variants leading to increased activity usually have an opposite effect. On the other hand, genetic variants can also influence the concentrations of the perpetrator drug, such as the CYP2C19 substrate voriconazole, thereby modifying the extent of inhibition/induction.(80) Moreover, for example, fluvoxamine

inhibited the oral clearance of theophylline by 62% in healthy subjects, but only by 12% in patients with severe cirrhosis, indicating that patients with severe cirrhosis are not susceptible to this DDI.(81) The conduct of clinical studies on these types of complex DDIs is based on the characteristics of the victim drug, i.e., which enzymes and transporters are relevant to the drug. Such studies typically require a larger clinical study, which includes an adequate number of subjects in each genotype or disease group, or a multi-phase crossover study, where different perpetrators and perpetrator combinations can be compared. The study designs are relatively straightforward, but the recruitment of a sufficient number of individuals with a rare genotype or rare disease may be challenging.

PITFALLS AND INTERPRETATION OF DDI STUDIES

Pitfalls

There are multiple potential pitfalls in the design and conduct DDI studies that can lead to inaccurate results, ethical problems, false alerts or in the worst case, failure to identify relevant DDI risks (**Table 1**). Particularly deficiencies in preclinical data or lack of understanding of the importance of specific pharmacokinetic pathways have led to failures to identify clinically hazardous DDIs. The past cases that have led to market withdrawals, such as those of cerivastatin, cisapride and terfenadine as victims and sorivudine as a perpetrator, have been caused mainly by failure to understand basic mechanistic issues, together with dose-dependent toxicity of the victim drugs.(82) In the following, we are not presenting an exhaustive review of all possible pitfalls, but instead describe three case examples of pitfalls that can lead to inappropriate conclusions of the DDI risk.

In late 1990's, the importance of CYP2C8 in drug metabolism was not understood. Eventually, the gemfibrozil-cerivastatin interaction revealed the potential significance of the enzyme, after a series of *in vitro* and clinical studies that were needed to understand the mechanisms of this DDI.(83) Initially, cerivastatin was brought to the market with arguments of a favorable interaction potential due to metabolism by both CYP3A4 and CYP2C8. However, it was shortly thereafter

withdrawn from the market in 2001 owing to multiple cases of rhabdomyolysis. In about half of the cases, the patients had been using gemfibrozil and cerivastatin concomitantly, and sadly, many of the cases had a fatal outcome. Soon, gemfibrozil was shown to increase the AUC of cerivastatin 5-6-fold and to greatly reduce the formation of its CYP2C8-dependent M-23 metabolite, whereas strong CYP3A4 inhibition by itraconazole had only a minor effect on cerivastatin exposure.(84, 85) At first, gemfibrozil was identified as a rather modest and nonselective inhibitor of CYP2C8 *in vitro*, and it was only later discovered that the main metabolite of gemfibrozil, gemfibrozil 1-O- β -glucuronide is a strong mechanism-based inhibitor of CYP2C8.(86) This case concretely illustrated that it is difficult to anticipate DDIs without systematic studies and thorough understanding of the DDI characteristics of both the victim and perpetrator drug.

The COX-2 selective nonsteroidal anti-inflammatory drug (NSAID) rofecoxib was initially considered to be a modest inhibitor of CYP1A2. This information in the labeling was mainly based on a DDI study using theophylline as a CYP1A2 index substrate, where rofecoxib 25 mg once daily increased the AUC of theophylline only 1.5-fold.(87) However, a DDI study with a similar design using tizanidine as a probe substrate resulted in a 13.6-fold increase in its AUC, indicating strong CYP1A2 inhibition by rofecoxib.(88) The reason for this apparent discrepancy is theophylline's lack of sensitivity as an *in vivo* index substrate of CYP1A2, partly due to its high bioavailability and lack of first-pass metabolism. At the time, *in vitro* data regarding the effect of rofecoxib on CYP1A2 was also lacking, and it was only later shown that rofecoxib is a potent, mechanism-based inhibitor of CYP1A2.(89)

As demonstrated in **Figure 2**, the inhibitor dose and timing of the substrate in relation to inhibitor administration can greatly affect the magnitude of the observed interaction. An example of this is provided by examining drug interaction studies involving the intestinal CYP3A inhibitor grapefruit juice and the CYP3A substrate lovastatin. In the study by Rogers et al., grapefruit juice was administered for four days in the morning and 40 mg lovastatin was administered in the evening of day 3.(90) Grapefruit juice increased the AUC of lovastatin 1.9-fold and that of lovastatin acid 1.6-fold only. It is notable that the exact dosing interval between the perpetrator and victim is not given in

the paper. The results are in contrast with an earlier study by Kantola et al., where grapefruit juice increased the AUC values of lovastatin and lovastatin acid by 15-fold and 5-fold, respectively. In this study, grapefruit juice was administered thrice daily for two days, and lovastatin was administered simultaneously with grapefruit juice on day 3, followed by additional grapefruit juice doses 0.5 and 1.5 hours after lovastatin (91). Even though this amount of grapefruit juice intake is not likely to occur frequently in real life, the results represent a potential worst-case scenario of the grapefruit juice-lovastatin interaction. This example also makes evident how important it is to consider the study design both when planning and interpreting the results of a DDI study.

During the past two decades, DDI guidelines have been frequently updated by the regulatory authorities, and most serious pitfalls can now be avoided by following the guidelines. However, knowledge on non-CYP enzyme-mediated metabolism and transporters still continues to emerge, and is going to further improve DDI study methodology and the predictability of DDIs. Yet, it is still difficult to evaluate the quantitative roles of all relevant disposition pathways of a drug based on *in vitro* data. Consequently, a potentially dangerous pitfall is to use PBPK modeling in situations where its reliability without additional clinical data is questionable, e.g., in the case of complex DDIs as outlined above.

Interpretation and extrapolation of findings

It is usually fairly straightforward to interpret the findings of standard DDI studies with index inhibitors of index substrates, and to classify the tested drug's sensitivity to inhibition of a specific pathway or its strength as an inhibitor or inducer of a transporter or enzyme. Key issues to consider when interpreting DDI data are the strength and selectivity of the index perpetrator at the given dose and the sensitivity of the index substrate, as well as the general design of the study including the dose- and time relationships (**Figure 2**). For example, if the index drug is not sensitive enough, even full inhibition of a single metabolic pathway does not produce an over 5-fold AUC change that would classify the perpetrator drug as a strong inhibitor.

When it comes to interpretation of complex situations and extrapolation of the findings to other scenarios, PBPK modeling can be highly valuable. Yet, if multiple mechanisms are involved in the DDI, such as involvement of both transporters and metabolism, or simultaneous induction and inhibition, extrapolation of the findings to other dosing scenarios or different drugs should be done with caution. In cases where a single mechanism, e.g., CYP3A4 inhibition, is crucial, extrapolation of the findings to other scenarios can be made with more confidence. It should be emphasized, however, that a very sensitive but nonselective index substrate due to combined roles of metabolism and transport, can be superior to a poorly sensitive index substrate, when the purpose is to detect a DDI risk. On the other hand, when the clinical significance of a DDI is evaluated, also complex issues, such as multiple simultaneous mechanisms, time- and dose-dependency of transporter/enzyme inhibition and induction, as well as time- and dose-dependent pharmacokinetics of the victim drug, often need to be considered in interpretation and extrapolation of the findings. As highlighted by the midostaurin and imatinib cases above, the sensitivity of a victim drug to DDIs may be altered by time-dependent changes in its own pharmacokinetics. Moreover, the magnitude of DDIs occurring during the absorption and first-pass metabolism of the victim may depend on the dosing interval between the perpetrator and victim, which should be considered in the interpretation, e.g., using PBPK modeling.

CONCLUSION

Evaluation of the DDI potential of drugs under development and on the market is a crucial issue for drug safety. Advances in *in vitro* methodology and modeling methods have greatly advanced our understanding of molecular mechanisms of DDIs and the ways to predict and interpret them. Clinical DDI studies remain, however, an integral part of the process of evaluating the DDI risks. Moreover, clinical DDI studies have revealed previously unknown DDIs and shed light on their mechanisms, which have then been confirmed by e.g., *in vitro* studies through reverse translation. As clinical DDI studies may not cover every single permutation of various parameters affecting the outcome, mechanistically focused clinical studies in concert with *in silico* models have great synergy in providing comprehensive understanding of DDIs and their clinical relevance. In conclusion, there is

no optimal way to study DDIs, but the design of every study needs to be based on careful evaluation of the available data, to ensure safety and maximal usefulness of the study.

REFERENCES

- (1) Dechanont, S., Maphanta, S., Butthum, B. & Kongkaew, C. Hospital admissions/visits associated with drug-drug interactions: a systematic review and meta-analysis. *Pharmacoepidemiol Drug Saf* **23**, 489-97 (2014).
- (2) US Food and Drug Administration. Center for Drug Evaluation and Research. *Clinical Drug Interaction Studies — Study Design, Data Analysis, and Clinical Implications Guidance for Industry*. <<https://www.fda.gov/downloads/drugs/guidances/ucm292362.pdf>> (2017). Accessed 15 February 2019.
- (3) European Medicines Agency. *Guideline on the investigation of drug interactions*. <https://www.ema.europa.eu/documents/scientific-guideline/guideline-investigation-drug-interactions_en.pdf> (2012). Accessed 15 February 2019.
- (4) Yu, J., Petrie, I.D., Levy, R.H. & Ragueneau-Majlessi, I. Mechanisms and Clinical Significance of Pharmacokinetic-Based Drug-Drug Interactions with Drugs Approved by the U.S. Food and Drug Administration in 2017. *Drug Metab Dispos* **47**, 135-44 (2019).
- (5) Backman, J.T., Kivistö, K.T., Olkkola, K.T. & Neuvonen, P.J. The area under the plasma concentration-time curve for oral midazolam is 400-fold larger during treatment with itraconazole than with rifampicin. *Eur J Clin Pharmacol* **54**, 53-8 (1998).
- (6) de Jong, J. *et al.* Effect of CYP3A perpetrators on ibrutinib exposure in healthy participants. *Pharmacol Res Perspect* **3**, e00156 (2015).
- (7) Fuhr, U., Hsin, C.H., Li, X., Jabrane, W. & Sörgel, F. Assessment of Pharmacokinetic Drug-Drug Interactions in Humans: In Vivo Probe Substrates for Drug Metabolism and Drug Transport Revisited. *Annu Rev Pharmacol Toxicol* **59**, 507-36 (2019).
- (8) Grimstein, M. *et al.* Physiologically Based Pharmacokinetic Modeling in Regulatory Science: An Update From the U.S. Food and Drug Administration's Office of Clinical Pharmacology. *J Pharm Sci* **108**, 21-5 (2019).
- (9) Prueksaritanont, T. *et al.* Drug-drug interaction studies: regulatory guidance and an industry perspective. *AAPS J* **15**, 629-45 (2013).
- (10) Rekić, D. *et al.* Clinical Drug-Drug Interaction Evaluations to Inform Drug Use and Enable Drug Access. *J Pharm Sci* **106**, 2214-8 (2017).
- (11) Floyd, J.S. *et al.* A screening study of drug-drug interactions in cerivastatin users: an adverse effect of clopidogrel. *Clin Pharmacol Ther* **91**, 896-904 (2012).
- (12) Tornio, A. *et al.* Glucuronidation converts clopidogrel to a strong time-dependent inhibitor of CYP2C8: a phase II metabolite as a perpetrator of drug-drug interactions. *Clin Pharmacol Ther* **96**, 498-507 (2014).
- (13) Niemi, M., Backman, J.T., Fromm, M.F., Neuvonen, P.J. & Kivistö, K.T. Pharmacokinetic interactions with rifampicin : clinical relevance. *Clin Pharmacokinet* **42**, 819-50 (2003).

- (14) Backman, J.T., Luurila, H., Neuvonen, M. & Neuvonen, P.J. Rifampin markedly decreases and gemfibrozil increases the plasma concentrations of atorvastatin and its metabolites. *Clin Pharmacol Ther* **78**, 154-67 (2005).
- (15) Lau, Y.Y., Huang, Y., Frassetto, L. & Benet, L.Z. effect of OATP1B transporter inhibition on the pharmacokinetics of atorvastatin in healthy volunteers. *Clin Pharmacol Ther* **81**, 194-204 (2007).
- (16) Niemi, M., Pasanen, M.K. & Neuvonen, P.J. Organic anion transporting polypeptide 1B1: a genetically polymorphic transporter of major importance for hepatic drug uptake. *Pharmacol Rev* **63**, 157-81 (2011).
- (17) Hukkanen, J. Induction of cytochrome P450 enzymes: a view on human in vivo findings. *Expert Rev Clin Pharmacol* **5**, 569-85 (2012).
- (18) Zamek-Gliszczynski, M.J. *et al.* Transporters in Drug Development: 2018 ITC Recommendations for Transporters of Emerging Clinical Importance. *Clin Pharmacol Ther* **104**, 890-9 (2018).
- (19) International Transporter Consortium *et al.* Membrane transporters in drug development. *Nat Rev Drug Discov* **9**, 215-36 (2010).
- (20) Elsby, R., Martin, P., Surry, D., Sharma, P. & Fenner, K. Solitary Inhibition of the Breast Cancer Resistance Protein Efflux Transporter Results in a Clinically Significant Drug-Drug Interaction with Rosuvastatin by Causing up to a 2-Fold Increase in Statin Exposure. *Drug Metab Dispos* **44**, 398-408 (2016).
- (21) Lund, M., Petersen, T.S. & Dalhoff, K.P. Clinical Implications of P-Glycoprotein Modulation in Drug-Drug Interactions. *Drugs* **77**, 859-83 (2017).
- (22) Jalava, K.M., Partanen, J. & Neuvonen, P.J. Itraconazole decreases renal clearance of digoxin. *Ther Drug Monit* **19**, 609-13 (1997).
- (23) US Food and Drug Administration. Center for Drug Evaluation and Research. *Clinical Pharmacology Biopharmaceutics Review(s): Pradaxa*. <https://www.accessdata.fda.gov/drugsatfda_docs/nda/2010/022512Orig1s000ClinPharmR_Corrected%203.11.2011.pdf> (2010). Accessed 19 February 2019.
- (24) Chu, X. *et al.* Clinical Probes and Endogenous Biomarkers as Substrates for Transporter Drug-Drug Interaction Evaluation: Perspectives From the International Transporter Consortium. *Clin Pharmacol Ther* **104**, 836-64 (2018).
- (25) Zamek-Gliszczynski, M.J. *et al.* ITC Commentary on Metformin Clinical Drug-Drug Interaction Study Design That Enables an Efficacy- and Safety-Based Dose Adjustment Decision. *Clin Pharmacol Ther* **104**, 781-4 (2018).
- (26) Guo, Y. *et al.* Advancing Predictions of Tissue and Intracellular Drug Concentrations Using In Vitro, Imaging and Physiologically Based Pharmacokinetic Modeling Approaches. *Clin Pharmacol Ther* **104**, 865-89 (2018).
- (27) Kajosaari, L.I., Niemi, M., Neuvonen, M., Laitila, J., Neuvonen, P.J. & Backman, J.T. Cyclosporine markedly raises the plasma concentrations of repaglinide. *Clin Pharmacol Ther* **78**, 388-99 (2005).
- (28) Tornio, A. & Backman, J.T. Cytochrome P450 in Pharmacogenetics: An Update. *Adv Pharmacol* **83**, 3-32 (2018).
- (29) Yee, S.W. *et al.* Influence of Transporter Polymorphisms on Drug Disposition and Response: A Perspective From the International Transporter Consortium. *Clin Pharmacol Ther* **104**, 803-17 (2018).

- Accepted Article
- (30) Bank, P.C.D. *et al.* Comparison of the Guidelines of the Clinical Pharmacogenetics Implementation Consortium and the Dutch Pharmacogenetics Working Group. *Clin Pharmacol Ther* **103**, 599-618 (2018).
 - (31) Fuhr, U., Jetter, A. & Kirchheiner, J. Appropriate phenotyping procedures for drug metabolizing enzymes and transporters in humans and their simultaneous use in the "cocktail" approach. *Clin Pharmacol Ther* **81**, 270-83 (2007).
 - (32) Mikus, G. Probes and Cocktails for Drug-Drug Interaction Evaluation: The Future Is Microdosing? *Clin Pharmacol Ther*, E-pub ahead of print, DOI: 10.1002/cpt.1350 (2019).
 - (33) Bosilkovska, M. *et al.* Geneva cocktail for cytochrome p450 and P-glycoprotein activity assessment using dried blood spots. *Clin Pharmacol Ther* **96**, 349-59 (2014).
 - (34) Bosilkovska, M. *et al.* Evaluation of Mutual Drug-Drug Interaction within Geneva Cocktail for Cytochrome P450 Phenotyping using Innovative Dried Blood Sampling Method. *Basic Clin Pharmacol Toxicol* **119**, 284-90 (2016).
 - (35) Derungs, A., Donzelli, M., Berger, B., Noppen, C., Krahenbuhl, S. & Haschke, M. Effects of Cytochrome P450 Inhibition and Induction on the Phenotyping Metrics of the Basel Cocktail: A Randomized Crossover Study. *Clin Pharmacokinet* **55**, 79-91 (2016).
 - (36) Stopfer, P. *et al.* Pharmacokinetic Evaluation of a Drug Transporter Cocktail Consisting of Digoxin, Furosemide, Metformin, and Rosuvastatin. *Clin Pharmacol Ther* **100**, 259-67 (2016).
 - (37) Stopfer, P. *et al.* Effects of Metformin and Furosemide on Rosuvastatin Pharmacokinetics in Healthy Volunteers: Implications for Their Use as Probe Drugs in a Transporter Cocktail. *Eur J Drug Metab Pharmacokinet* **43**, 69-80 (2018).
 - (38) Stopfer, P. *et al.* Optimization of a drug transporter probe cocktail: potential screening tool for transporter-mediated drug-drug interactions. *Br J Clin Pharmacol* **84**, 1941-9 (2018).
 - (39) Prueksaritanont, T. *et al.* Validation of a microdose probe drug cocktail for clinical drug interaction assessments for drug transporters and CYP3A. *Clin Pharmacol Ther* **101**, 519-30 (2017).
 - (40) Chu, X., Chan, G.H. & Evers, R. Identification of Endogenous Biomarkers to Predict the Propensity of Drug Candidates to Cause Hepatic or Renal Transporter-Mediated Drug-Drug Interactions. *J Pharm Sci* **106**, 2357-67 (2017).
 - (41) Mariappan, T.T., Shen, H. & Marathe, P. Endogenous Biomarkers to Assess Drug-Drug Interactions by Drug Transporters and Enzymes. *Curr Drug Metab* **18**, 757-68 (2017).
 - (42) Müller, F., Sharma, A., König, J. & Fromm, M.F. Biomarkers for In Vivo Assessment of Transporter Function. *Pharmacol Rev* **70**, 246-77 (2018).
 - (43) Bonate, P.L. *et al.* Methods and strategies for assessing uncontrolled drug-drug interactions in population pharmacokinetic analyses: results from the International Society of Pharmacometrics (ISOP) Working Group. *J Pharmacokinet Pharmacodyn* **43**, 123-35 (2016).
 - (44) Wang, D.D. *et al.* The Utility of a Population Approach in Drug-Drug Interaction Assessments: A Simulation Evaluation. *J Clin Pharmacol* **57**, 1268-78 (2017).
 - (45) Masters, J.C., Harano, D.M., Greenberg, H.E., Tsunoda, S.M., Jang, I.J. & Ma, J.D. Limited sampling strategy of partial area under the concentration-time curves to estimate midazolam systemic clearance for cytochrome P450 3A phenotyping. *Ther Drug Monit* **37**, 84-9 (2015).

- (46) Chang, A.T. *et al.* S-Warfarin Limited Sampling Models to Estimate Area Under the Concentration Versus Time Curve for Cytochrome P450 2C9 Baseline Activity and After Induction. *Ther Drug Monit* **38**, 383-7 (2016).
- (47) Fuhr, U. & Rost, K.L. Simple and reliable CYP1A2 phenotyping by the paraxanthine/caffeine ratio in plasma and in saliva. *Pharmacogenetics* **4**, 109-16 (1994).
- (48) de Andrés, F., Terán, S., Bovera, M., Fariñas, H., Terán, E. & Llerena, A. Multiplex Phenotyping for Systems Medicine: A One-Point Optimized Practical Sampling Strategy for Simultaneous Estimation of CYP1A2, CYP2C9, CYP2C19, and CYP2D6 Activities Using a Cocktail Approach. *OMICS* **20**, 88-96 (2016).
- (49) Sager, J.E., Yu, J., Ragueneau-Majlessi, I. & Isoherranen, N. Physiologically Based Pharmacokinetic (PBPK) Modeling and Simulation Approaches: A Systematic Review of Published Models, Applications, and Model Verification. *Drug Metab Dispos* **43**, 1823-37 (2015).
- (50) Yoshida, K. *et al.* In Vitro-In Vivo Extrapolation of Metabolism- and Transporter-Mediated Drug-Drug Interactions-Overview of Basic Prediction Methods. *J Pharm Sci* **106**, 2209-13 (2017).
- (51) European Medicines Agency. *Guideline on the qualification and reporting of physiologically based pharmacokinetic (PBPK) modelling and simulation.* <http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2016/07/WC500211315.pdf> (2016). Accessed 15 February 2019.
- (52) US Food and Drug Administration. Center for Drug Evaluation and Research. *Physiologically Based Pharmacokinetic Analyses — Format and Content Guidance for Industry.* <<https://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM531207.pdf>> (2018). Accessed 15 February 2019.
- (53) Yeo, K.R., Jamei, M. & Rostami-Hodjegan, A. Predicting drug-drug interactions: application of physiologically based pharmacokinetic models under a systems biology approach. *Expert Rev Clin Pharmacol* **6**, 143-57 (2013).
- (54) Huang, S.M. & Rowland, M. The role of physiologically based pharmacokinetic modeling in regulatory review. *Clin Pharmacol Ther* **91**, 542-9 (2012).
- (55) Shebley, M. *et al.* Physiologically Based Pharmacokinetic Model Qualification and Reporting Procedures for Regulatory Submissions: A Consortium Perspective. *Clin Pharmacol Ther* **104**, 88-110 (2018).
- (56) Jones, H.M. *et al.* Physiologically based pharmacokinetic modeling in drug discovery and development: a pharmaceutical industry perspective. *Clin Pharmacol Ther* **97**, 247-62 (2015).
- (57) Wagner, C. *et al.* Application of Physiologically Based Pharmacokinetic (PBPK) Modeling to Support Dose Selection: Report of an FDA Public Workshop on PBPK. *CPT Pharmacometrics Syst Pharmacol* **4**, 226-30 (2015).
- (58) Pan, Y. *et al.* The Application of Physiologically Based Pharmacokinetic Modeling to Predict the Role of Drug Transporters: Scientific and Regulatory Perspectives. *J Clin Pharmacol* **56 Suppl 7**, S122-31 (2016).
- (59) Shebley, M., Fu, W., Badri, P., Bow, D. & Fischer, V. Physiologically Based Pharmacokinetic Modeling Suggests Limited Drug-Drug Interaction Between Clopidogrel and Dasabuvir. *Clin Pharmacol Ther* **102**, 679-87 (2017).

- Accepted Article
- (60) Varma, M.V., Lai, Y., Feng, B., Litchfield, J., Goosen, T.C. & Bergman, A. Physiologically based modeling of pravastatin transporter-mediated hepatobiliary disposition and drug-drug interactions. *Pharm Res* **29**, 2860-73 (2012).
- (61) Yoshikado, T. *et al.* Quantitative Analyses of Hepatic OATP-Mediated Interactions Between Statins and Inhibitors Using PBPK Modeling With a Parameter Optimization Method. *Clin Pharmacol Ther* **100**, 513-23 (2016).
- (62) Shitara, Y. & Sugiyama, Y. Preincubation-dependent and long-lasting inhibition of organic anion transporting polypeptide (OATP) and its impact on drug-drug interactions. *Pharmacol Ther* **177**, 67-80 (2017).
- (63) VandenBrink, B.M. & Isoherranen, N. The role of metabolites in predicting drug-drug interactions: focus on irreversible cytochrome P450 inhibition. *Curr Opin Drug Discov Devel* **13**, 66-77 (2010).
- (64) Varma, M.V., Pang, K.S., Isoherranen, N. & Zhao, P. Dealing with the complex drug-drug interactions: towards mechanistic models. *Biopharm Drug Dispos* **36**, 71-92 (2015).
- (65) Filppula, A.M., Laitila, J., Neuvonen, P.J. & Backman, J.T. Potent mechanism-based inhibition of CYP3A4 by imatinib explains its liability to interact with CYP3A4 substrates. *Br J Pharmacol* **165**, 2787-98 (2012).
- (66) Filppula, A.M., Neuvonen, M., Laitila, J., Neuvonen, P.J. & Backman, J.T. Autoinhibition of CYP3A4 leads to important role of CYP2C8 in imatinib metabolism: variability in CYP2C8 activity may alter plasma concentrations and response. *Drug Metab Dispos* **41**, 50-9 (2013).
- (67) Dutreix, C., Munarini, F., Lorenzo, S., Roesel, J. & Wang, Y. Investigation into CYP3A4-mediated drug-drug interactions on midostaurin in healthy volunteers. *Cancer Chemother Pharmacol* **72**, 1223-34 (2013).
- (68) Wang, Y., Yin, O.Q., Graf, P., Kisicki, J.C. & Schran, H. Dose- and time-dependent pharmacokinetics of midostaurin in patients with diabetes mellitus. *J Clin Pharmacol* **48**, 763-75 (2008).
- (69) Yin, O.Q., Wang, Y. & Schran, H. A mechanism-based population pharmacokinetic model for characterizing time-dependent pharmacokinetics of midostaurin and its metabolites in human subjects. *Clin Pharmacokinet* **47**, 807-16 (2008).
- (70) Simonson, S.G. *et al.* Rosuvastatin pharmacokinetics in heart transplant recipients administered an antirejection regimen including cyclosporine. *Clin Pharmacol Ther* **76**, 167-77 (2004).
- (71) Niemi, M., Backman, J.T., Neuvonen, M. & Neuvonen, P.J. Effects of gemfibrozil, itraconazole, and their combination on the pharmacokinetics and pharmacodynamics of repaglinide: potentially hazardous interaction between gemfibrozil and repaglinide. *Diabetologia* **46**, 347-51 (2003).
- (72) US Food and Drug Administration. Center for Drug Evaluation and Research. *Clinical Pharmacology Biopharmaceutics Review(s): Vosevi*. <https://www.accessdata.fda.gov/drugsatfda_docs/nda/2017/209195Orig1s000ClinPharmR.pdf> (2017). Accessed 18 February 2019.
- (73) Foisy, M.M., Yakiwchuk, E.M. & Hughes, C.A. Induction effects of ritonavir: implications for drug interactions. *Ann Pharmacother* **42**, 1048-59 (2008).
- (74) Yeung, C.K., Fujioka, Y., Hachad, H., Levy, R.H. & Isoherranen, N. Are circulating metabolites important in drug-drug interactions?: Quantitative analysis of risk prediction and inhibitory potency. *Clin Pharmacol Ther* **89**, 105-13 (2011).

- (75) Backman, J.T., Filppula, A.M., Niemi, M. & Neuvonen, P.J. Role of Cytochrome P450 2C8 in Drug Metabolism and Interactions. *Pharmacol Rev* **68**, 168-241 (2016).
- (76) Storelli, F., Samer, C., Reny, J.L., Desmeules, J. & Daali, Y. Complex Drug-Drug-Gene-Disease Interactions Involving Cytochromes P450: Systematic Review of Published Case Reports and Clinical Perspectives. *Clin Pharmacokinet* **57**, 1267-93 (2018).
- (77) Niemi, M., Tornio, A., Pasanen, M.K., Fredrikson, H., Neuvonen, P.J. & Backman, J.T. Itraconazole, gemfibrozil and their combination markedly raise the plasma concentrations of loperamide. *Eur J Clin Pharmacol* **62**, 463-72 (2006).
- (78) Brøsen, K., Hansen, J.G., Nielsen, K.K., Sindrup, S.H. & Gram, L.F. Inhibition by paroxetine of desipramine metabolism in extensive but not in poor metabolizers of sparteine. *Eur J Clin Pharmacol* **44**, 349-55 (1993).
- (79) Yu, K.S. *et al.* Effect of omeprazole on the pharmacokinetics of moclobemide according to the genetic polymorphism of CYP2C19. *Clin Pharmacol Ther* **69**, 266-73 (2001).
- (80) Weiss, J. *et al.* CYP2C19 genotype is a major factor contributing to the highly variable pharmacokinetics of voriconazole. *J Clin Pharmacol* **49**, 196-204 (2009).
- (81) Orlando, R., Padriani, R., Perazzi, M., De Martin, S., Piccoli, P. & Palatini, P. Liver dysfunction markedly decreases the inhibition of cytochrome P450 1A2-mediated theophylline metabolism by fluvoxamine. *Clin Pharmacol Ther* **79**, 489-99 (2006).
- (82) Onakpoya, I.J., Heneghan, C.J. & Aronson, J.K. Delays in the post-marketing withdrawal of drugs to which deaths have been attributed: a systematic investigation and analysis. *BMC Med* **13**, 26 (2015).
- (83) Tornio, A., Neuvonen, P.J., Niemi, M. & Backman, J.T. Role of gemfibrozil as an inhibitor of CYP2C8 and membrane transporters. *Expert Opin Drug Metab Toxicol* **13**, 83-95 (2017).
- (84) Backman, J.T., Kyrklund, C., Neuvonen, M. & Neuvonen, P.J. Gemfibrozil greatly increases plasma concentrations of cerivastatin. *Clin Pharmacol Ther* **72**, 685-91 (2002).
- (85) Kantola, T., Kivistö, K.T. & Neuvonen, P.J. Effect of itraconazole on cerivastatin pharmacokinetics. *Eur J Clin Pharmacol* **54**, 851-5 (1999).
- (86) Ogilvie, B.W. *et al.* Glucuronidation converts gemfibrozil to a potent, metabolism-dependent inhibitor of CYP2C8: implications for drug-drug interactions. *Drug Metab Dispos* **34**, 191-7 (2006).
- (87) Bachmann, K. *et al.* An evaluation of the dose-dependent inhibition of CYP1A2 by rofecoxib using theophylline as a CYP1A2 probe. *J Clin Pharmacol* **43**, 1082-90 (2003).
- (88) Backman, J.T., Karjalainen, M.J., Neuvonen, M., Laitila, J. & Neuvonen, P.J. Rofecoxib is a potent inhibitor of cytochrome P450 1A2: studies with tizanidine and caffeine in healthy subjects. *Br J Clin Pharmacol* **62**, 345-57 (2006).
- (89) Karjalainen, M.J., Neuvonen, P.J. & Backman, J.T. Rofecoxib is a potent, metabolism-dependent inhibitor of CYP1A2: implications for in vitro prediction of drug interactions. *Drug Metab Dispos* **34**, 2091-6 (2006).
- (90) Rogers, J.D. *et al.* Grapefruit juice has minimal effects on plasma concentrations of lovastatin-derived 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Clin Pharmacol Ther* **66**, 358-66 (1999).
- (91) Kantola, T., Kivistö, K.T. & Neuvonen, P.J. Grapefruit juice greatly increases serum concentrations of lovastatin and lovastatin acid. *Clin Pharmacol Ther* **63**, 397-402 (1998).

- Accepted Article
- (92) Suzuki, E. *et al.* Identification of valproic acid glucuronide hydrolase as a key enzyme for the interaction of valproic acid with carbapenem antibiotics. *Drug Metab Dispos* **38**, 1538-44 (2010).
- (93) Myllylä, V.V., Sotaniemi, K.A., Illi, A., Suominen, K. & Keränen, T. Effect of entacapone, a COMT inhibitor, on the pharmacokinetics of levodopa and on cardiovascular responses in patients with Parkinson's disease. *Eur J Clin Pharmacol* **45**, 419-23 (1993).
- (94) Diasio, R.B. Sorivudine and 5-fluorouracil; a clinically significant drug-drug interaction due to inhibition of dihydropyrimidine dehydrogenase. *Br J Clin Pharmacol* **46**, 1-4 (1998).
- (95) Grana, E., Barbieri, A., Zonta, F. & Lucchelli, A. Potentiating actions of tranylcypromine and moclobemide on the sympathomimetic activity of dopamine. *Pharmacol Res* **24**, 337-43 (1991).
- (96) Cleary, J.M., Rosen, L.S., Yoshida, K., Rasco, D., Shapiro, G.I. & Sun, W. A phase 1 study of the pharmacokinetics of nucleoside analog trifluridine and thymidine phosphorylase inhibitor tipiracil (components of TAS-102) vs trifluridine alone. *Invest New Drugs* **35**, 189-97 (2017).
- (97) Corona, G. *et al.* Lopinavir-ritonavir dramatically affects the pharmacokinetics of irinotecan in HIV patients with Kaposi's sarcoma. *Clin Pharmacol Ther* **83**, 601-6 (2008).
- (98) Kasichayanula, S., Liu, X., Griffen, S.C., Lacreata, F.P. & Boulton, D.W. Effects of rifampin and mefenamic acid on the pharmacokinetics and pharmacodynamics of dapagliflozin. *Diabetes Obes Metab* **15**, 280-3 (2013).
- (99) Gidal, B.E., Sheth, R., Parnell, J., Maloney, K. & Sale, M. Evaluation of VPA dose and concentration effects on lamotrigine pharmacokinetics: implications for conversion to lamotrigine monotherapy. *Epilepsy Res* **57**, 85-93 (2003).
- (100) Zimm, S., Collins, J.M., O'Neill, D., Chabner, B.A. & Poplack, D.G. Inhibition of first-pass metabolism in cancer chemotherapy: interaction of 6-mercaptopurine and allopurinol. *Clin Pharmacol Ther* **34**, 810-7 (1983).

FIGURE LEGENDS

Figure 1. Investigation of DDIs. (a) Signals of a potential DDI may arise from a variety of sources. (b) Signals need careful assessment before further action is taken. (c) The golden standard for clinical DDI studies is a prospective cross-over study with index drugs, usually performed in healthy volunteers. (d) The interpretation of the obtained results determines the (e) implications of the study.

Figure 2. Design of DDI studies. (a) Effect of varying inhibitor dose (scenarios A-C) on the plasma concentrations of a substrate drug. A sufficient dose of the perpetrator drug is necessary to reach strong inhibition, and detect and accurately quantify a potential DDI. (b) Effect of a CYP inhibitor on the plasma concentrations of a substrate with varying fractions metabolized by the affected CYP ($f_{m,CYP}$). In combination with a CYP inhibitor, a substrate drug with a low $f_{m,CYP}$ will give a smaller DDI than a sensitive substrate drug with a high $f_{m,CYP}$. (c) Effect of dose staggering (scenarios A-D) on DDI magnitude. Strongest DDI can be detected when the victim is administered shortly after the inhibitor. (d) Effect of repeated dosing of perpetrator whose metabolite inhibits the CYP enzyme (scenarios A-C) on DDI magnitude. To study the worst-case scenario, the perpetrator should be dosed to steady-state and victim drug given at the time of the peak concentrations of the perpetrator. Bid, twice daily; od, once daily; sd, single dose.

Figure 3. Pharmacokinetics and DDIs of midostaurin. (a) Midostaurin exhibits time-dependent pharmacokinetics, with increasing trough plasma concentrations during the first 3-6 days of therapy, followed by a 60-70% decline in concentrations until a steady state is reached.(68, 69) Autoinduction of CYP3A4 by midostaurin and its two major metabolites is thought to be involved in the time-dependent pharmacokinetics of midostaurin.(69) (b) Midostaurin is a sensitive CYP3A4 substrate, and ketoconazole has increased its single dose AUC by more than tenfold.(67) (c) Although both midostaurin and its two major metabolites are inducers and mechanism-based inhibitors of CYP3A4 in vitro, the drug has had no effect on midazolam pharmacokinetics in healthy subjects, following either a single midostaurin dose (100 mg) or two days after the last dose of multiple midostaurin doses (50 mg twice daily for three days).(67) Clinical data shown are from (67, 68). Bid, twice daily; d, day; od, once daily; sd, single dose; tid, three times daily.

SUPPLEMENTARY MATERIAL

Table S1. Characteristics of possible index substrates of cytochrome P450 enzymes. The table contains all FDA and EMA recommended index substrates and some selected alternatives based on potential advantages in sensitivity, selectivity and/or relative safety.

Table S2. Characteristics of possible index inhibitors of cytochrome P450 enzymes. The table contains all FDA and EMA recommended index inhibitors and some selected alternatives based on potential advantages in sensitivity, selectivity and/or relative safety.

Table S3. Characteristics of transporter substrates and inhibitors.

Table 1. List of recommended considerations and requirements of interventional DDI studies and examples of potential pitfalls in DDI studies and their interpretation.

Typical requirements of a DDI study	Potential pitfalls of DDI studies
General design issues	
Healthy volunteers if no safety concerns	Risky drugs given to healthy subjects
Patients if safety concerns or clinical focus	Bias and confounding in observational patient studies
Crossover design	Parallel group design may produce bias
Sufficient wash-out to eliminate carry-over effects	Insufficient wash-out (e.g., a slowly eliminated metabolite still present)
Placebo control and blinding (e.g., in case of pharmacodynamic end points)	
Dietary control/restrictions as necessary	
Appropriate sample collection, storage and analytical method to cover > 80-90% of the AUC of the victim and metabolites	Inaccurate or insensitive analytical method, degradation of analytes during storage or analysis
Monitoring of perpetrator pharmacokinetics (compliance, quantification of exposure, presence after washout)	Perpetrator exposure not documented
Pharmacodynamic assessment	Pharmacodynamic assessments neglected
DNA samples	
Biomarker samples in selected cases	
Necessary prior knowledge considered in design	Deficiencies in preclinical and early clinical data (e.g., in mass-balance studies)
Safety issues	
Strict exclusion criteria (e.g., contraindications,	Careless exclusion criteria leading to risk of

pregnancy)	adverse effects
History, clinical examination, laboratory tests, and genotyping as necessary	
Safety monitoring and sufficient follow-up	Insufficient follow-up (residual drug effects)
Precautions and interventions to avoid adverse effects, even in the worst-case DDI scenario	Rescue interventions not prearranged
Blood sampling should generally not exceed the volume of blood donation	
Perpetrator (inhibitor/inducer)	
Selectivity and strength of index inhibitor (> 5-fold increase in AUC possible)	Suboptimal strength Nonselective inhibitor
Clinically relevant (high) dose depending on tolerability	Too low dose, leading to weak inhibition
Dosing to reach and maintain steady-state, including time-dependent inhibition and induction	Clinically atypical dosing Duration of dosing too short/long to document maximal DDI
Victim substrate	
Sensitivity of index substrate	Lack of sensitivity (particularly if not considered in interpretation)
Documented selectivity of index substrate	Lack of selectivity (particularly if not considered in interpretation)
High first-pass and short half-life preferable	Long half-life victim with a short-acting or "presystemic" inhibitor
Monitoring a specific metabolic ratio can be useful	
Dose, expecting the worst-case scenario	Too high dose
Staggered dosing	Victim drug administered too soon (or too long) after perpetrator to document maximal DDI

Table 2. Characteristics of possible index substrates of cytochrome P450 enzymes. The table contains all FDA and EMA recommended index substrates and some selected alternatives based on potential advantages in sensitivity, selectivity and/or relative safety.

Enzyme	Substrate	Sensitivity ^a	Other relevant enzymes/transporters	F	t _{1/2} (h)	Remarks	E/F ^b
CYP1A2	Agomelatine	++++		0.05	1-2	- Limited availability in some countries	
	Caffeine*	++++	NAT, XO	1.00	3-7		E/F
	Melatonin	++++	CYP1A1 ^c	0.03	0.9		
	Theophylline	++	CYP3A4, CYP2E1	0.96	9	- Adverse effects at high concentrations	E
	Tizanidine*	++++		0.34	2.5	- Adverse effects at high concentrations	F
CYP2B6	Bupropion*	+	11β-HSD1 ^c	0.9	11	- Hydroxybupropion to bupropion AUC ratio is a selective and more sensitive marker of CYP2B6 than bupropion AUC	E
	(S)-Ketamine	++	CYP3A4	0.08	2-6	- Parenteral formulation can be given orally	
	Efavirenz	n/a (++)	CYP2A6	n/a	52-76	- Limited DDI data	E
CYP2C8	Amodiaquine	n/a		n/a	5	- Limited DDI data - Limited availability - Metabolite t _{1/2} >100 h	E
	Daprodustat	++++		n/a	1	- Limited DDI data - Limited availability, currently in phase 3	
	Dasabuvir*	++++	CYP3A4	0.70	5.5-6		
	Repaglinide*	+++	CYP3A4, OATP1B1	0.56	0.8	- Adverse effects at high concentrations	E/F
CYP2C9	(S)-Warfarin	++ (+++)		0.93 (racemic)	21-43	- Bleeding risk	E/F
	Flurbiprofen	++		0.92	5.5	- 4'-hydroxyflurbiprofen/flurbiprofen ratio is a sensitive marker of CYP2C9 - Limited availability of appropriate formulation	
	Fluvastatin*	+ (++)	OATP1B1	0.29	0.7		
	Tolbutamide*	++ (+++)		0.8-0.9	5.9	- Limited availability - Adverse effects at high concentrations	E/F
CYP2C19	Lansoprazole	++	CYP3A4	0.81	0.9	- Delayed absorption	F

		(+++)					
	Omeprazole*	+++ (++++)	CYP3A4	0.53	0.7	- Delayed absorption - Inhibitor of CYP2C19	E/F
	Pantoprazole*	n/a (+++)		0.77	1.0	- Delayed absorption - Breath test established	
	Rabeprazole	++ (+++)	CYP3A4	0.52	1.5	- Delayed absorption	
CYP2D6	Desipramine	+++ (+++)	CYP3A4	0.38	28	- Limited availability in some countries	E/F
	Dextromethorphan*	++++ (++++)	CYP3A4	n/a	3.4	- Dextrophan/dextromethorphan ratio used as the index	F
	Metoprolol	++ (+++)	CYP3A4	0.38	3.2		E
	Nebivolol	+++ (++++)	CYP2C19	n/a	11		F
	Tolterodine	++++ (++++)	CYP3A4	0.26 (NMs)	2.3 (NMs)		
CYP3A4	Bupirone	++++		0.04	2.4	- Sensitive to intestinal CYP3A4 inhibition	
	Midazolam*	++++		0.44	1.9	- i.v. formulation available to assess hepatic CYP3A4 - Adverse effects at high concentrations	E/F
	Sildenafil	++++	CYP2C9	0.38	2.4		
	Simvastatin (lactone)*	++++		<0.05	2-3	- Simvastatin acid substrate of OATP1B1 - Sensitive to intestinal CYP3A4 inhibition	
	Triazolam	++++		0.44	2.9	- Adverse effects at high concentrations	F

The data are compiled primarily from the UW Metabolism and Transport Drug Interaction Database (Copyright University of Washington 1999-2019. Accessed: January-February 2019), secondarily from drug labels.

^a Indication of sensitivity is based on data from a trial with an EMA/FDA-recommended index inhibitor of the affected CYP. In some cases, a second sensitivity value is given within brackets, based on pharmacogenetic data. Classification: +++++ a maximal >10-fold increase in AUC, +++ a maximal 5-10-fold increase, ++ a maximal 2-5-fold increase, + a maximal 1.25-2-fold increase. Values for maximal fold increases in AUC are given in **Table S1**.

^b Indicates if the drug is recommended as a clinical CYP probe substrate by EMA (E) and/or FDA (F) (2-3)

^c Based on *in vitro* data only.

Abbreviations: F, bioavailability; 11 β -HSD-1, 11 β -hydroxysteroid dehydrogenase type 1; NAT, N-acetyl transferase; NMs, normal metabolizers; n/a, not available; XO, xanthine oxidase.

* = Our recommended substrate

Table 3. Characteristics of possible index inhibitors of cytochrome P450 enzymes. The table contains all FDA and EMA recommended index inhibitors and some selected alternatives based on potential advantages in sensitivity, selectivity and/or relative safety.

Enzyme	Inhibitor	Dose used in DDI trials ^a	Strength ^b	Also inhibits	t _{1/2} (h)	Remarks	E/F ^c
CYP1A2	Ciprofloxacin*	500 mg bid	+++	CYP3A4	3.3	- Some safety concerns	
	Enoxacin	400 mg bid	+++		3.3	- Some safety concerns - Limited availability	E
	Fluvoxamine*	25 mg bid- 100 mg od	++++	CYP2C19, CYP2D6, CYP3A4	15	- Strong CYP2C19 inhibitor	F
CYP2B6	Ticlopidine*	250 mg bid	+	CYP2C19	98	- Strong CYP2B6 inhibitor based on substrate metabolic ratio - Strong CYP2C19 inhibitor - MBI - Limited availability - Bleeding risk	E
CYP2C8	Clopidogrel	75 mg od (first dose 300 mg)	+++	CYP2B6, CYP2C19	6	- Moderate CYP2C8 inhibitor with 75 mg daily - MBI (glucuronide) - Bleeding risk	F
	Gemfibrozil*	600 mg bid	++++	OATP1B1, OAT3	1.1	- MBI (glucuronide)	E/F
CYP2C9	Fluconazole*	200 mg od	++	CYP2C19, CYP3A4	32	- Strong CYP2C19 inhibitor	E/F
CYP2C19	Fluconazole*	200 mg od	+++	CYP2C9, CYP3A4	32		
	Fluoxetine	60 mg od	++	CYP2D6	53	- MBI - Strong CYP2D6 inhibitor - t _{1/2} of norfluoxetine 4-16 days	
	Fluvoxamine*	25 mg bid-100 mg od	+++	CYP1A2, CYP2D6, CYP3A4	15	- Strong CYP1A2 inhibitor	F
	Omeprazole*	20-40 mg od	++		0.7	- Strength based on non-sensitive probe substrates - MBI	E
CYP2D6	Bupropion	150-300 mg od	+++		11		
	Fluoxetine	60 mg od	++++	CYP2C19	53	- Strong CYP2C19 inhibitor - t _{1/2} of norfluoxetine 4-16 days	E/F
	Mirabegron	50 mg od	++		50		F
	Paroxetine*	20 mg od	++++		17	- MBI	E/F
	Quinidine	100 mg od, 100-600 mg sd	++++	P-gp	6.2	- Prolongs QT-interval	E
	Terbinafine	250 mg od	+++	CYP1A2	200		

CYP3A4	Clarithromycin	250-500 mg bid	+++	CYP2C19, P-gp	3.3	- MBI	E/F
	Cobicistat*	150 mg od	++++	CYP2D6, P-gp	5.2	- MBI	
	Erythromycin	500 mg tid	+++	P-gp ^d	1.6	- MBI	F
	Fluconazole	200 mg od	++	CYP2C19, CYP2C9	32	- Strong CYP2C19 inhibitor	F
	Itraconazole*	100-200 mg od	++++	CYP2J2 ^d , P-gp	21		E/F
	Ketoconazole	200 mg bid-400 mg od	++++	CYP2C19, P-gp	3.3	- Limited availability	E
	Posaconazole	300-400 mg bid	+++		31		
	Ritonavir*	100-200 mg bid	++++	P-gp	4	- MBI - Induces CYP enzymes	E
	Verapamil	80 mg tid-240 mg od	++	P-gp	4	- MBI	F

The data are compiled primarily from the UW Metabolism and Transport Drug Interaction Database (Copyright University of Washington 1999-2019. Accessed: January-February 2019), secondarily from drug labels.

^a Typical dose used in clinical DDI trials. Cocktail studies not included.

^b Indication of strength is based on a trial with an index substrate of the affected CYP. Cocktail studies not included. Classification: ++++ a maximal >10-fold increase in index substrate AUC, +++ a maximal 5-10-fold increase, ++ a maximal 2-5-fold increase, + a maximal 1.25-2-fold increase. Values for maximal fold increases in AUC are given in **Table S2**.

^c Indicates if the drug is recommended as a CYP probe inhibitor by EMA (E) and/or FDA (F) (2-3)

^d Erythromycin: based on *in vitro* data only. Itraconazole: based on interaction with astemizole.

Abbreviations: bid, twice daily; MBI, mechanism-based inhibition; od, once daily; sd, single dose; tid, three times daily

* = Our recommended inhibitor

Table 4. Examples of drug-drug interactions involving non-CYP drug-metabolizing enzymes.

Enzyme	Victim	Perpetrator (inhibitor)	Consequence
Acylpeptide hydrolase	Valproic acid glucuronide	Carbapenems, e.g., meropenem	Decreased serum valproic acid levels (92)
Catechol-O- methyltransferase (COMT)	Levodopa	Entacapone	AUC 1.5-fold (93)
Dihydropyrimidine dehydrogenase (DPD)	5-fluorouracil	Sorivudine	Fluoropyrimidine toxicity (94)
Monoamine oxidases (A and B)	Dopamine	Moclobemide	Increased dopamine effects (95)
Thymidine phosphorylase	Trifluridine	Tipiracil	AUC 37-fold (96)
UGT1A1	SN-38 (Irinotecan active metabolite)	Lopinavir-ritonavir	AUC 3-fold (97)
UGT1A9	Dapagliflozin	Mefenamic acid	AUC 1.5-fold (98)
UGT2B7	Lamotrigine	Valproic acid	AUC up to 2-fold (99)
Xanthine oxidase	6-mercaptopurine	Allopurinol	AUC 5-fold (100)

Table 5. Examples of transporter probe substrates and inhibitors for clinical DDI studies.

Transporter	Substrates	Inhibitors
BCRP	rosuvastatin	cyclosporine
	sulfasalazine	eltrombopag
OATP1B1 or OATP1B3	atorvastatin	cyclosporine
	pitavastatin	rifampin (single dose)
	pravastatin	
	repaglinide	
	rosuvastatin	
	simvastatin (acid)	
OATP2B1	celiprolol	apple, orange and grapefruit juices (intestine)
OCT1	sumatriptan	n/a
	metformin	
OCT2	metformin	cimetidine
OAT1	adefovir	pyrimethamine
		probenecid
OAT3	benzylpenicillin	probenecid
P-gp	aliskiren	itraconazole
	dabigatran etexilate	verapamil
	digoxin	
	fexofenadine	

The data are compiled primarily from the UW Metabolism and Transport Drug Interaction Database (Copyright University of Washington 1999-2019. Accessed: February 2019). More information regarding the properties of substrates and inhibitors are given in **Table S3**.

n/a, not available



