



UNIVERSITY OF HELSINKI

<https://helda.helsinki.fi>

Transport of statins by multidrug resistance-associated proteins 1 and 5

Tuomi, Suvi Kukka; Deng, Feng; Neuvonen, Mikko; Niemi, Mikko

2025-06-01

Elsevier B.V.

<http://hdl.handle.net/10138/595025>

Tuomi, S K, Deng, F, Neuvonen, M & Niemi, M 2025, 'Transport of statins by multidrug resistance-associated proteins 1 and 5', *European Journal of Pharmaceutical Sciences*, vol. 209, 107070. <https://doi.org/10.1016/j.ejps.2025.107070>

Downloaded from Helda, University of Helsinki institutional repository. <https://helda.helsinki.fi>
This is an electronic reprint of the original article.
This reprint may differ from the original in pagination and typographic detail.
Please cite the original version.



Transport of statins by multidrug resistance-associated proteins 1 and 5

Suvi-Kukka Tuomi^{a,b}, Feng Deng^{a,b,c}, Mikko Neuvonen^{a,b}, Mikko Niemi^{a,b,c,*} 

^a Department of Clinical Pharmacology, Faculty of Medicine, University of Helsinki, Finland

^b Individualized Drug Therapy Research Program, Faculty of Medicine, University of Helsinki, Finland

^c Department of Clinical Pharmacology, HUS Diagnostic Center, Helsinki University Hospital, Helsinki, Finland

ARTICLE INFO

Keywords:

Skeletal muscle toxicity
MRP1
MRP5
Statin-related myotoxicity
Fluvastatin
Pitavastatin

ABSTRACT

Statins are widely used in the treatment of hypercholesterolemia but also associated with muscle-related adverse effects. Multidrug resistance-associated protein (MRP) 1 and 5 are expressed in the skeletal muscle, where they may regulate intramuscular levels of their substrates. Here, we investigated the transport of various statins by MRP1 and MRP5 with the vesicular transport assay. Statin concentrations in the vesicles were determined with liquid chromatography tandem mass spectrometry. At 6 μM statin concentration, MRP1 transported both 3R,5S-fluvastatin and 3S,5R-fluvastatin with uptake ratios of 2.6 and 2.0. MRP5 transported 3R,5S-fluvastatin, 3S,5R-fluvastatin, and 10 μM pitavastatin with uptake ratios of 2.9, 3.7, and 2.6, respectively. Atorvastatin was only a weak substrate of MRP5 with an uptake ratio of 1.6 and was therefore not investigated further. In concentration-dependent transport experiments, racemic fluvastatin was transported by MRP1 and MRP5 with apparent affinities (K_m) of 225 μM and 23 μM . Pitavastatin was transported by MRP5 with a K_m value of 433 μM . *In vitro* clearance ($CL_{in\ vitro}$) of fluvastatin was 0.36 $\mu\text{l}/\text{min}/\text{mg}$ for MRP1, while MRP5 exhibited a $CL_{in\ vitro}$ value of 1.2 $\mu\text{l}/\text{min}/\text{mg}$ for fluvastatin and 0.21 $\mu\text{l}/\text{min}/\text{mg}$ for pitavastatin. Pravastatin, rosuvastatin, and simvastatin acid were not transported by MRP1 or MRP5. Atorvastatin and pitavastatin were not transported by MRP1. These data indicate that MRP1 transports fluvastatin, while MRP5 transports both fluvastatin and pitavastatin. Because MRP1 and MRP5 are expressed in the skeletal muscle, they may reduce myocyte exposure to fluvastatin and pitavastatin and protect from muscle toxicity.

1. Introduction

Cardiovascular diseases are among the most common causes of death worldwide (Roth et al., 2020). 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors, also called statins, are commonly used for the primary and secondary prevention of cardiovascular diseases (Fig. 1). Statins are effective in reducing low-density lipoprotein cholesterol levels, generally safe, and well-tolerated, with mostly mild adverse effects (Yebo et al., 2019). However, they can cause musculoskeletal adverse effects ranging from common and mild myalgia to rare and possibly life-threatening rhabdomyolysis (Harper and Jacobson, 2007). Importantly, statin-related myotoxicity is dose- and concentration-dependent (Alfirevic et al., 2014; Neuvonen et al., 2006). Several factors, including drug interactions, pharmacogenetic variability, and advanced age, may increase the systemic statin exposure, leading to elevated statin concentrations in myocytes and skeletal

muscle toxicity (Cooper-DeHoff et al., 2022; Lönnberg et al., 2023; Turner and Pirmohamed, 2019).

Solute carrier (SLC) and ATP-binding cassette (ABC) transporters are membrane proteins which have a crucial role in the cellular influx and efflux of drugs, affecting their plasma concentrations and tissue distribution (Galetin et al., 2024; Giacomini et al., 2010; Koepsell, 2013; Vasiliou et al., 2008). Organic anion transporting polypeptides (OATP) 1B1, 1B3, and 2B1 mediate the influx of statins into the liver (Bi et al., 2019; Elsby et al., 2012; Mykkänen et al., 2022, 2024; Niemi et al., 2011). Breast cancer resistance protein (BCRP) and the P-glycoprotein, on the other hand, may restrict the intestinal absorption of statins (Deng et al., 2021; Elsby et al., 2012; Kesitalo, et al., 2009a, 2009b; Lehtisalo et al., 2023; Neuvonen et al., 2006). Several drug transporters are expressed in the skeletal muscle (Knauer et al., 2010). They may participate in regulating statin concentrations in myocytes and, because myotoxicity is concentration-dependent, contribute to statin-related

The authors declare no competing interest for this work. This study was supported by grants from the European Research Council (Grant Agreement 725249), Emil Aaltonen Foundation, Finnish Cultural Foundation, Orion Research Foundation, Sigrid Jusélius Foundation, and State Funding for University-level health research.

* Corresponding author at: Department of Clinical Pharmacology, University of Helsinki, PO Box 20, 00014, Finland.

E-mail address: mikko.niemi@helsinki.fi (M. Niemi).

<https://doi.org/10.1016/j.ejps.2025.107070>

Received 18 October 2024; Received in revised form 30 January 2025; Accepted 12 March 2025

Available online 13 March 2025

0928-0987/© 2025 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

myotoxicity.

OATP2B1 is an influx transporter expressed in the skeletal muscle and transports atorvastatin, fluvastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin acid (Grube et al., 2006; Ho et al., 2006; Kopplov et al., 2005; Nozawa et al., 2004; Prieto Garcia et al., 2024; Varma et al., 2011; Lehtisalo et al., 2023), but its role in the statin influx in skeletal muscle and myotoxicity is unclear. Additionally, monocarboxylate transporters (MCT) 1 and 4 are expressed in the skeletal muscle, and inhibition of MCT4 by statins has been linked with statin-induced cytotoxicity (Bonen et al., 1998; Kikutani et al., 2016; Kobayashi et al., 2006; Pilegaard et al., 1999). Multidrug resistance-associated protein (MRP) 1, 4, and 5 efflux transporters are highly expressed in the skeletal muscle, unlike other well-characterized ABC efflux transporters such as MRP2, BCRP, and P-gp (Knauer et al., 2010). Their physiological role in skeletal muscle cell is not well-characterized, but they may protect cells from endogenous metabolites and xenobiotics. MRP1 transports organic anions, conjugates, and anticancer drugs (Jedlitschky et al., 1996; Renes et al., 1999), whereas MRP4 and MRP5 transport cyclic nucleotides, antivirals, and anticancer drugs (Chen et al., 2001; Jedlitschky et al., 2000; Sampath et al., 2002; Schuetz et al., 1999). In addition, previous studies suggest that MRP1 and MRP5 may transport various statins (Afrouzian et al., 2018; ; Hoste et al., 2023; Kanamitsu et al., 2017; Knauer et al., 2010; Prieto Garcia et al., 2024). However, the studies have been conducted in different laboratories employing different methods making the comparison between statins challenging.

Statin-related myotoxicity can be life-threatening, but even when mild, the toxicity can be clinically important as it often leads to discontinuation of therapy (Alfirevic et al., 2014; Cooper-DeHoff et al., 2022; Voora et al., 2009). Therefore, we considered it important to investigate how statins are transported by efflux transporters expressed in the skeletal muscle. Previously, we characterized and compared the transport of atorvastatin, fluvastatin, pitavastatin, pravastatin, rosuvastatin, simvastatin acid, and four atorvastatin metabolites by six ABC transporters (BCRP, MRP2, MRP3, MRP4, MRP5, P-glycoprotein) (Deng et al., 2021). Here, we investigate the transport of atorvastatin, two fluvastatin enantiomers, pitavastatin, pravastatin, rosuvastatin, and

simvastatin acid by MRP1 and MRP5.

2. Materials and methods

2.1. Materials

Atorvastatin, atorvastatin-d5, fluvastatin-d8, pitavastatin-d5, pravastatin, pravastatin-d9, rosuvastatin, rosuvastatin-d6, and simvastatin acid-d6 were purchased from Toronto Research Chemicals (Toronto, Ontario, Canada). Racemic fluvastatin, 3R,5S-fluvastatin, 3S,5R-fluvastatin, and pitavastatin were obtained from Santa Cruz Biotechnology (Dallas, TX, USA). Simvastatin acid was acquired from SynFine Research (Richmond Hill, Ontario, Canada). MRP1, MRP5, and mock-transfected control vesicles derived from human embryonic kidney 293 cells were purchased from PharmTox (Radboud UMC, Nijmegen, The Netherlands). Solvents used in assays and analytical methods were of analytical quality and purchased from Sigma-Aldrich (St. Louis, MO, USA). Ultrapure water for assays and analyses was purified with Milli-Q water purification system (Merck Millipore, Burlington, MA, USA).

2.2. Vesicular transport assay

The vesicular transport assay was used to investigate statin transport by MRP1 and MRP5 *in vitro*. The assays were carried out with inside-out oriented membrane vesicles over-expressing transporters, and performed as described previously (Deng et al., 2021). In short, membrane vesicles (7.5 μ g) were preincubated at 37 °C for 10 min in the reaction buffer containing various concentrations of statin, assay buffer (PharmTox), and 10 mM MgCl₂. The transport was initiated by adding prewarmed ATP or AMP (4 mM) to the mixture. Samples were incubated at 37 °C for the indicated time. The reactions were terminated with ice-cold stop buffer (PharmTox) and samples were quickly transferred to a MultiScreenHTS FB Filter Plate 1.0 mm/0.65 mm (Merck KGaA, Darmstadt, Germany). The vesicles were immediately washed three times with the stop buffer and twice with ice-cold wash buffer (40 mM 4-morpholinepropanesulfonic acid/Tris pH 7.0 and 70 mM KCl) and dried with airflow. The accumulated statins were eluted from the




Fig 1. Chemical structures of the investigated statins.

vesicles with the lysis solution (50 % methanol and 25 ng/ml isotope-labeled statin as internal standard). All assays were performed in triplicates on 96-well plates.

The functionality of the membrane vesicles and the vesicular transport assay was confirmed by examining the transport of estradiol-17-glucuronide and 5(6)-carboxy-2',7'-dichlorofluorescein by MRP1 and MRP5, respectively. Initially, the transport of 10 and 3 μM of atorvastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin acid and 6 μM 3R,5S-fluvastatin and 3S,5R-fluvastatin into MRP1 and MRP5 vesicles were screened with a 10 min incubation time. Statins that exhibited statistically significantly higher transport into the vesicles in the presence of ATP compared to AMP, and had an uptake ratio of 2.0 or higher, were considered as substrates of MRP1 or MRP5, and were studied further.

The time-dependent transport of 10 μM atorvastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin acid and 6 μM fluvastatin enantiomers were investigated with incubation times of 3, 6, and 10 min. Finally, the concentration-dependent transport was determined with statin concentrations ranging from 6 to 150 μM , using an incubation time of 10 min. The appropriate incubation time for the concentration-dependent transport experiments was selected based on time-dependent transport experiments. Additionally, in time-dependent and concentration-dependent transport experiments, statin transport into control vesicles was investigated.

2.3. Analytical methods

Statins were analyzed using a SCIEX 5500 quadrupole linear ion trap liquid chromatography-tandem mass spectrometry system (Sciex, Toronto, Ontario, Canada) with an electrospray ionization ion source. Chromatographic separation utilized a Luna Omega polar C18 column (100 \times 2.1 mm, 1.6 μm ; Phenomenex, Torrance, CA, USA) and a corresponding guard column. Mobile phases A and B consisted of 5 mM ammonium formate (pH 3.9) and acetonitrile, with a flow rate of 300 $\mu\text{L}/\text{min}$ and a column temperature of 40 $^{\circ}\text{C}$. The mobile phase gradient included a hold at 20 % B for 1 min, a linear increase to 40 % B over 3 min, then to 90 % B over 2 min, and 1 min at 90 % B, followed by re-equilibration to 20 % B. The mass spectrometer operated in multiple reaction monitoring (MRM) mode, with specific ion transitions previously documented (Deng et al., 2021).

2.4. Data and statistical analysis

Data are presented as mean \pm standard deviation (SD) or mean with 95 % confidence of interval (CI). ATP-dependent transport was determined by subtracting the statin uptake into vesicles in the presence of AMP from the uptake in the presence of ATP. The uptake ratio of transport was determined by dividing the uptake of statin in the presence of ATP by the uptake in the presence of AMP. Statistical comparisons in the 6 and 10 μM screening and time-dependent transport experiments were carried out using analysis of variance (GraphPad Prism Software version 10.12, San Diego, CA). A *P* value of below 0.05 was considered statistically significant.

When determining the kinetic parameters of statin transport, we pooled the mean ATP-dependent transport values for each concentration data point from separate experiments. These pooled values were then fitted to the Michaelis-Menten equation Eq. (1), where v represents the velocity of ATP-dependent transport, V_{max} represents the maximum transport rate, $[S]$ represents the substrate concentration, and K_m represents the Michaelis-Menten constant.

$$v = \frac{V_{\text{max}} [S]}{K_m + [S]} \quad (1)$$

The *in vitro* clearance ($CL_{\text{in vitro}}$) of statins was calculated by using Michaelis-Menten parameters, as described in Eq. (2).

$$CL_{\text{in vitro}} = \frac{V_{\text{max}}}{K_m} \quad (2)$$

Molecular descriptors were calculated in R version 4.4.2 with RStudio (R Core Team, 2024). The Simplified Molecular Input Line Entry Specifications (SMILES) of the investigated statins were retrieved from the PubChem database (<https://pubchem.ncbi.nlm.nih.gov/>). Molecular descriptors were calculated based on the retrieved SMILES using the rcdk package, which allows the access to the Chemistry development kit (CDK) Java libraries for chemoinformatics (Guha, 2007).

3. Results

3.1. Transport of known substrate of MRP1 and MRP5

The functionality of the vesicles was confirmed using 15 μM estradiol-17-glucuronide as a positive control for MRP1 and 5 μM 5(6)-carboxy-2',7'-dichlorofluorescein for MRP5. The uptake ratios of these probe substrates in MRP1 and MRP5 were 7.14 and 4.44, respectively, validating both the vesicle functionality and the vesicular transport assay (Supplementary Figure S1).

3.2. Screening for statin efflux transport

MRP1 transported 3R,5S-fluvastatin and 3S,5R-fluvastatin (6 μM) with uptake ratios of 2.6 ± 0.5 and 2.0 ± 0.1 , whereas MRP5 transported 3R,5S-fluvastatin, 3S,5R-fluvastatin, and pitavastatin (10 μM) with uptake ratios of 2.9 ± 0.1 , 3.7 ± 0.1 , and 2.6 ± 1.0 , respectively (Fig. 2; Supplementary Table S1). Pravastatin, rosuvastatin, and simvastatin acid were not transported by either MRP1 or MRP5 (Fig. 2; Supplementary Figure S2; Supplementary Tables S1-2). Additionally, atorvastatin and pitavastatin were not transported by MRP1. Atorvastatin uptake by MRP5 in the presence of ATP differed statistically significantly from that in the presence of AMP. However, the uptake ratio was below two (Supplementary Tables S1-2), indicating that atorvastatin is only a weak substrate of MRP5 and was therefore not investigated further.

3.3. Time-dependent transport

The ATP-dependent transport of fluvastatin enantiomers into MRP1 and MRP5 vesicles and pitavastatin into MRP5 vesicles was time-dependent and differed statistically significantly from control vesicles latest at the 10 min time point (Fig. 3; Supplementary Table S3).

3.4. Concentration-dependent transport

To determine the kinetic parameters of statin transport, we next investigated the concentration-dependent transport of fluvastatin into MRP1, MRP5, and control vesicles, and pitavastatin into MRP5 and control vesicles (Fig. 4 and Table 1). Due to the similar levels of transport observed for the fluvastatin enantiomers, racemic fluvastatin was used to study concentration-dependent transport. Fluvastatin was transported into MRP1 and MRP5 vesicles with K_m values of 225 and 23 μM , and with V_{max} values of 81 and 27 pmol/min/mg. The $CL_{\text{in vitro}}$ values of fluvastatin by MRP1 and MRP5 were 0.36 and 1.2 $\mu\text{L}/\text{min}/\text{mg}$. Pitavastatin was transported by MRP5 with a K_m value of 433 μM , a V_{max} of 92 pmol/min/mg and a $CL_{\text{in vitro}}$ of 0.21 $\mu\text{L}/\text{min}/\text{mg}$.

3.5. Molecular descriptors of the investigated statins

According to molecular descriptors, fluvastatin and pitavastatin have smaller total polar surface areas (81 – 90 vs 104 – 148), number of rotatable bonds (8 vs 10 – 13), and number of hydrogen bond acceptors (5 vs 6 – 9) than the rest of the investigated statins (Supplementary Table S4).




Fig. 2. The uptake of statins into MRP1 and MRP5 vesicles. The statin concentrations were 10 μM , except 6 μM for fluvastatin enantiomers. The incubation time was 10 min and amount of protein 7.5 μg . Dark and light bars represent the mean of two or three experiments in the presence of ATP and AMP, respectively. The means of single experiments are indicated by triangles and circles. * $P < 0.05$, ** $P < 0.01$ compared to AMP.




Fig. 3. The time-dependent transport of 6 μM fluvastatin enantiomers into MRP1, MRP5, and control vesicles, and 10 μM pitavastatin into MRP5 and control vesicles. The amount of protein was 7.5 μg . Results were obtained from two (three for 3S,5R-fluvastatin and MRP5) separate experiments with triplicate samples. * $P < 0.05$, ** $P < 0.01$ compared to control vesicles.




Fig. 4. The concentration-dependent transport of fluvastatin into MRP1, MRP5, and control vesicles, and pitavastatin into MRP5 and control vesicles. The incubation time was 10 min and amount of protein 7.5 μg , except for the transport of fluvastatin by MRP1, which was studied with a protein amount of 11.25 μg . Results were obtained from two to four separate experiments with triplicate samples.

4. Discussion

Skeletal muscle ABC transporters, such as MRP1 and MRP5, may regulate the intracellular concentrations of statins in myocytes and contribute to statin-related myotoxicity. In this study, we investigated the transport of atorvastatin, 3R,5S-fluvastatin, 3S,5R-fluvastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin acid by MRP1 and MRP5 *in vitro* with the vesicular transport assay. Since all the statins

were investigated with the same method in a single laboratory, the results are directly comparable between statins and with the MRP4 data from our previous study, which employed the same method in the same laboratory (Deng et al., 2021). Our findings indicate that MRP1 transports fluvastatin, whereas MRP5 transports both fluvastatin and pitavastatin.

In addition to MRP1 and MRP5, MRP4 has been detected in the skeletal muscle (Knauer et al., 2010). In our previous study employing

Table 1

Estimated kinetic parameters of fluvastatin, pitavastatin, and rosuvastatin transport by MRP1, MRP4, and MRP5.

Statin	Transporter	V _{max} (95 % CI) (pmol/min/mg)	K _m (95 % CI) (μM)	CL _{in vitro} (μl/min/mg)	Number of experiments
Racemic fluvastatin	MRP1	81.2 (54.1-176)	225 (118-620)	0.361	2-3
Racemic fluvastatin (Deng et al., 2021)	MRP4	54.7 (23.4-4053)	31.8 (0-∞)	1.72	3
Racemic fluvastatin	MRP5	26.9 (18.1-45.0)	22.8 (4.89-91.0)	1.18	3
Pitavastatin	MRP5	92.1 (24.1-∞)	433 (52.4-∞)	0.213	3-4
Rosuvastatin (Deng et al., 2021)	MRP4	12.3 (5.13-∞)	39.3 (0.3-∞)	0.312	3

CL_{in vitro}, in vitro clearance; K_m, Michaelis-Menten constant; V_{max}, maximum transport rate.

the same methodology, MRP4 transported fluvastatin and rosuvastatin, but not other statins (Deng et al., 2021) (Table 2). The CL_{in vitro} values of MRP4 for fluvastatin and rosuvastatin were 1.7 and 0.31 μl/min/mg. In the current study, the CL_{in vitro} for MRP1-mediated fluvastatin transport was 0.36 μl/min/mg, while MRP5 transported fluvastatin and pitavastatin with CL_{in vitro} values of 1.2 and 0.21 μl/min/mg. In the previous study, the clinically important statin efflux transporter BCRP transported fluvastatin and pitavastatin with substantially higher CL_{in vitro} values (16 and 7.3 μl/min/mg) than any of the MRPs investigated in our present and previous study (Deng et al., 2021). While the chemical structures of fluvastatin and pitavastatin are quite similar, the observed K_m and CL_{in vitro} values of fluvastatin and pitavastatin in MRP5 vesicles were strikingly different. The nitrogen atom of the quinoline ring in pitavastatin, capable of accepting a hydrogen atom, may contribute to the different transport kinetics in MRP5. Moreover, fluvastatin and

pitavastatin have smaller polar surface areas, and less rotatable bonds and hydrogen bond acceptors than other statins. This suggests that the rigid structure and low hydrogen bonding in addition to lipophilicity could be beneficial for the interaction between statins and MRP1 and MRP5. Overall, the absolute protein abundances of MRP1, MRP4, and MRP5 remain unclear, thus restricting the assessment of their relative contributions to statin efflux in the skeletal muscle.

Unlike previous studies, we observed no transport of atorvastatin, pravastatin, or simvastatin acid by MRP1 nor that of rosuvastatin by MRP5. Two studies using HeLa and HEK293 cell accumulation assays suggested that atorvastatin is an MRP1 substrate (Hoste et al., 2023; Knauer et al., 2010). Because atorvastatin is relatively lipophilic (Turner and Pirmohamed, 2019), passive diffusion and non-specific binding of atorvastatin may have masked small ATP-dependent uptake of atorvastatin into MRP1 vesicles in our study. Additionally, Knauer and co-authors showed that rosuvastatin is a good substrate for MRP1 and MRP4, and relatively poor substrate for MRP5 (Knauer et al., 2010). In our study, the uptake of rosuvastatin into MRP1 vesicles in the presence of ATP was 1.9-fold higher than that in the presence of AMP, but the difference was not statistically significant. In a recent study, MRP1 was suggested to transport pravastatin and simvastatin acid in *Spodoptera frugiperda* (Sf9) membrane vesicles (Prieto Garcia et al., 2024). However, control vesicle and uptake ratio data were not reported, which complicates the interpretation of the results. Nevertheless, another study showed that 1 μM pravastatin was transported in Sf9-MRP1 vesicles with an uptake ratio of 5.0 (Afrouzian et al., 2018).

Instead of whole cell-based methods, we chose to use membrane vesicles as they enable investigating the direct interaction between the drug and transporter, and precise determination of transport kinetics. Moreover, we chose human cell line-derived vesicles since they exhibit higher activity than the insect cell-derived Sf9 vesicles (Sáfár et al., 2021). In addition, post-translational protein modification differs between insect and human cells and the amount of membrane cholesterol is lower in insect cells (Pál et al., 2007). Cholesterol can be loaded to the Sf9 vesicles to increase transporter activity, but this also activates endogenous ABC transporters, possibly SfABCC2 and SfABCC3 (Sjöstedt et al., 2019).

Atorvastatin, fluvastatin, pitavastatin, and simvastatin acid are relatively lipophilic, whereas pravastatin and rosuvastatin are relatively hydrophilic (Neuvonen et al., 2006; Turner and Pirmohamed, 2019). It appears that lipophilic statins are more myotoxic than the hydrophilic

Table 2

Summary of atorvastatin, fluvastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin acid transport by MRP1, MRP4, MRP5, and OATP2B1.

		MRP1 (K _m)	MRP4 (K _m)	MRP5 (K _m)	OATP2B1 (K _m)	References
Atorvastatin	Current study	-	n/a	+	n/a	
	Literature	+	+	-	2.8 μM	(Hoste et al., 2023; Karlgren et al., 2012; Knauer et al., 2010)
Fluvastatin	Current study	225 μM	n/a	23 μM	n/a	
	Literature	n/a	32 μM	n/a	0.7 - 0.8 μM	(Bi et al., 2019; Deng et al., 2021; Kopplow et al., 2005; Noé et al., 2007; Varma et al., 2011)
Pitavastatin	Current study	-	n/a	443 μM	n/a	
	Literature	n/a	+	n/a	1.2 - 25 μM	(Hirano et al., 2006; Kanamitsu et al., 2017; Kanda et al., 2018; Ölander et al., 2016)
Pravastatin	Current study	-	n/a	-	n/a	
	Literature	2.5 - 8.7 μM	+	n/a	+	(Afrouzian et al., 2018; Bi et al., 2019; Kanamitsu et al., 2017; Nozawa et al., 2004; Prieto Garcia et al., 2024; Varma et al., 2011)
Rosuvastatin	Current study	-	n/a	-	n/a	
	Literature	+	21 - 39 μM	+	2.4 - 6.4 μM	(Ho et al., 2006; Deng et al., 2021; Kanda et al., 2018; Kitamura et al., 2008; Knauer et al., 2010; Lehtisalo et al., 2023; Pfeifer et al., 2013)
Simvastatin acid	Current study	-	n/a	-	n/a	
	Literature	179 μM	-	n/a	384 μM	(Deng et al., 2021; Prieto Garcia et al., 2024)

+, transported, but K_m value not available; -, not transported; n/a, not available.

statins *in vitro*. For example, pravastatin and rosuvastatin were less cytotoxic in C2C12 myoblasts than atorvastatin, fluvastatin, and simvastatin acid (Schirris et al., 2015). In addition, in satellite cells isolated from human skeletal muscle biopsies, pravastatin showed lower myotoxicity than atorvastatin, fluvastatin, and simvastatin acid (Skotthjem et al., 2008). Since MRP1 and MRP5 are expressed in the skeletal muscle, they may play a protective role against fluvastatin- and pitavastatin-induced muscle toxicity. Supporting this hypothesis, MK-571, dipyridamole, quercetin, and verapamil, known inhibitors of MRP1, MRP4, MRP5, and OATP2B1, have increased the intracellular retention of atorvastatin and rosuvastatin in cultured human skeletal muscle myoblasts (Knauer et al., 2010). Moreover, overexpression of OATP2B1 has elevated atorvastatin and rosuvastatin accumulation and toxicity in myoblasts (Knauer et al., 2010). In addition, coadministration of the MRP1 inhibitor probenecid with rosuvastatin in rats led to increased skeletal muscle toxicity (Dorajoo et al., 2008).

MRP1 and MRP5 expression have been detected in skeletal muscle by western blot analyses (Knauer et al., 2010). It has been estimated that MRP1 expression in skeletal muscle is 60-fold higher than that of MRP5 (Prieto Garcia et al., 2024), but absolute quantitative evidence, such as liquid chromatography-tandem mass spectrometry-based proteomics, are lacking. This absence of quantitative data makes it challenging to estimate the *in vivo* significance of these transporters in humans. MRP1 and MRP5 are also expressed in various other tissues across the body, for example in the brain and the placenta (Meyer zu Schwabedissen et al., 2005; Nies et al., 2004; Stage et al., 2020; St-Pierre et al., 2000). In barrier tissues such as the placenta, efflux transporters may restrict the entry of harmful compounds. Moreover, MRP5 may play a role in intracellular signaling, since it transports cyclic nucleotides (Jedlitschky et al., 2000; Meyer zu Schwabedissen et al., 2005).

Interestingly, MRP5 is also expressed in the human heart (Dazert et al., 2003). The absence of the skeletal muscle influx transporter MCT4 in cardiomyocytes has been suggested to protect against statin-induced cardiomyotoxicity (Sirvent et al., 2005). The presence of efflux transporters like MRP5, might provide additional protection against statin-induced cardiotoxicity by transporting statins out of the cardiomyocytes. In addition to statin-induced muscle toxicity, statins dose-dependently increase the risk of type 2 diabetes (Lotta et al., 2016; Sattar et al., 2010). However, the underlying mechanism of the risk increase is not known. Some studies have reported individual cases of polyneuropathy associated with statin treatment (Law and Rudnica, 2006). As MRP1 is expressed in neurons (Stage et al., 2020), it is tempting to suggest that it might play a protective role against this condition.

The influx and efflux of statins by skeletal muscle transporters can vary due to drug interactions or genetic variability. For example, genetic variability in *SLCO2B1* has been associated with altered rosuvastatin exposure (Lehtisalo et al., 2023). In a recent study, simvastatin acid and pravastatin were found as *in vitro* OATP2B1 and MRP1 substrates (Prieto Garcia et al., 2024). The authors used proteomics-informed physiologically based pharmacokinetic modelling to simulate the effect of 90 % reduced OATP2B1 or MRP1 activity on the statin plasma and tissue exposure. The modelling suggested that the reduced OATP2B1 activity would decrease muscle-to-plasma ratio of simvastatin acid by 8.5-fold, while the reduced MRP1 activity would increase muscle-to-plasma ratio of simvastatin acid by 5-fold. Moreover, the presence of MRP1 was estimated to reduce the muscle simvastatin acid and pravastatin exposure by 13- and 2-fold compared with passive distribution. This model implies that MRP1 may affect the statin concentration in skeletal muscle and thus protect from statin-induced myotoxicity.

5. Conclusion

In conclusion, our present and previous findings demonstrate that MRP1, MRP4, and MRP5, which are expressed in the skeletal muscle, transport fluvastatin *in vitro*, while MRP5 additionally transports

pitavastatin, and MRP4 transports rosuvastatin. These data indicate that statins differ in their skeletal muscle efflux transport profiles. Moreover, the data can be utilized to improve prediction of statin-induced adverse effects and to enhance physiologically based pharmacokinetic models. In the future, comprehensive investigations into the expression of skeletal muscle drug transporters and the transport of statins by other skeletal muscle transporters, such as MCTs and OATP2B1, are essential to understand the factors influencing statin concentrations in the skeletal muscle. In addition, linking statin-induced muscle toxicity to transporter expression is important for understanding the transporter-related mechanisms of muscle toxicity. Recognizing and considering the role of transporters in adverse effects becomes crucial in the clinical use of statins to proactively manage and anticipate statin-induced muscle toxicity.

CRedit authorship contribution statement

Suvi-Kukka Tuomi: Writing – original draft, Methodology, Formal analysis, Conceptualization. **Feng Deng:** Writing – review & editing, Supervision, Methodology, Formal analysis, Conceptualization. **Mikko Neuvonen:** Writing – review & editing, Methodology, Formal analysis. **Mikko Niemi:** Writing – review & editing, Supervision, Funding acquisition, Conceptualization.

Acknowledgements

The authors thank the FIMM Technology Centre supported by the HiLIFE and Biocenter Finland for providing access to the plate reader instrumentation services. The graphical abstract was created in Bio-Rend. This work was supported by the European Research Council Consolidator Grant (grant agreement 725249), the Sigrid Jusélius Foundation (Helsinki, Finland), and State Funding for University-level Health Research. Feng Deng was supported by the Finnish Cultural Foundation (Helsinki, Finland) and Orion Research Foundation (Helsinki, Finland), and Suvi-Kukka Tuomi by the Finnish Cultural Foundation and the Emil Aaltonen Foundation (Tampere, Finland).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.ejps.2025.107070.

Data availability

Data will be made available on request.

References

- Afrouzian, M., Al-Lahham, R., Patrikeeva, S., Xu, M., Fokina, V., Fischer, W.G., Abdel-Rahman, S.Z., Costantine, M., Ahmed, M.S., Nanovskaya, T., 2018. Role of the efflux transporters BCRP and MRP1 in human placental bio-disposition of pravastatin. *Biochem. Pharmacol.* 156, 467–478. <https://doi.org/10.1016/j.bcp.2018.09.012>.
- Alfirevic, A., Neely, D., Armitage, J., Chinoy, H., Cooper, R.G., Laaksonen, R., Carr, D.F., Bloch, K.M., Fahy, J., Hanson, A., Yue, Q.-Y., Wadelius, M., Maitland-van Der Zee, A. H., Vooora, D., Psaty, B.M., Palmer, C.N.A., Pirmohamed, M., 2014. Phenotype standardization for statin-induced myotoxicity. *Clin. Pharmacol. Ther.* 96, 470–476. <https://doi.org/10.1038/clpt.2014.121>.
- Bi, Y.-A., Costales, C., Mathialagan, S., West, M., Eatemadpour, S., Lazzaro, S., Tyllaska, L., Scialis, R., Zhang, H., Umland, J., Kimoto, E., Tess, D.A., Feng, B., Tremaine, L.M., Varma, M.V.S., Rodrigues, A.D., 2019. Quantitative contribution of six major transporters to the hepatic uptake of drugs: “SLC-Phenotyping” using primary human hepatocytes. *J. Pharmacol. Exp. Ther.* 370, 72–83. <https://doi.org/10.1124/jpet.119.257600>.
- Bonen, A., McCullagh, K.J.A., Putman, C.T., Hultman, E., Jones, N.L., Heigenhauser, G.J. F., 1998. Short-term training increases human muscle MCT1 and femoral venous lactate in relation to muscle lactate. *Am. J. Physiol.* 274, E102–E107. <https://doi.org/10.1152/ajpendo.1998.274.1.E102>.
- Chen, Z.-S., Lee, K., Kruh, G.D., 2001. Transport of cyclic nucleotides and estradiol 17-β-d-glucuronide by multidrug resistance protein 4. *J. Biol. Chem.* 276, 33747–33754. <https://doi.org/10.1074/jbc.M104833200>.

- Cooper-DeHoff, R.M., Niemi, M., Ramsey, L.B., Luzum, J.A., Tarkiainen, E.K., Straka, R. J., Gong, L., Tuteja, S., Wilke, R.A., Wadelius, M., Larson, E.A., Roden, D.M., Klein, T.E., Yee, S.W., Krauss, R.M., Turner, R.M., Palaniappan, L., Gaedigk, A., Giacomini, K.M., Caudle, K.E., Voora, D., 2022. The clinical pharmacogenetics implementation consortium guideline for *SLCO1B1*, *ABCG2*, and *CYP2C9* genotypes and statin-associated musculoskeletal symptoms. *Clin. Pharmacol. Ther.* 111, 1007–1021. <https://doi.org/10.1002/cpt.2557>.
- Dazert, P., Meissner, K., Vogelgesang, S., Heydrich, B., Eckel, L., Böhm, M., Warzok, R., Kerb, R., Brinkmann, U., Schaeffeler, E., Schwab, M., Cascorbi, I., Jedlitschky, G., Kroemer, H.K., 2003. Expression and localization of the multidrug resistance protein 5 (MRP5/ABCC5), a cellular export pump for cyclic nucleotides, in human heart. *Am. J. Pathol.* 163, 1567–1577. [https://doi.org/10.1016/S0002-9440\(10\)63513-4](https://doi.org/10.1016/S0002-9440(10)63513-4).
- Deng, F., Tuomi, S.-K., Neuvonen, M., Hirvensalo, P., Kulju, S., Wenzel, C., Oswald, S., Filppula, A.M., Niemi, M., 2021. Comparative hepatic and intestinal efflux transport of statins. *Drug Metab. Dispos.* 49, 750–759. <https://doi.org/10.1124/dmd.121.000430>.
- Dorajoo, R.S., Pereira, B.P., Yu, Z., Gopalakrishnakone, P., Leong, C.C., Wee, A., Lee, E., 2008. Role of multidrug resistance-associated protein-1 transporter in statin-induced myopathy. *Life Sci* 82, 823–830. <https://doi.org/10.1016/j.lfs.2008.01.021>.
- Elsby, R., Hilgendorf, C., Fenner, K., 2012. Understanding the critical disposition pathways of statins to assess drug–drug interaction risk during drug development: It's not just about OATP1B1. *Clin. Pharmacol. Ther.* 92, 584–598. <https://doi.org/10.1038/clpt.2012.163>.
- Galetin, A., Brouwer, K.L.R., Tweedie, D., Yoshida, K., Sjöstedt, N., Aleksunes, L., Chu, X., Evers, R., Hafey, M.J., Lai, Y., Matsson, P., Riselli, A., Shen, H., Sparreboom, A., Varma, M.V.S., Yang, J., Yang, X., Yee, S.W., Zamek-Gliszczynski, M.J., Fenner, K., Hillgren, K.M., Huang, S.-M., Sugiyama, Y., Giacomini, K.M., 2024. Membrane transporters in drug development and as determinants of precision medicine. *Nat. Rev. Drug Discov.* 23, 255–280. <https://doi.org/10.1038/s41573-023-00877-1>.
- Giacomini, K.M., Huang, S.-M., Tweedie, D.J., Benet, L.Z., Brouwer, K.L.R., Chu, X., Dahlin, A., Evers, R., Fischer, V., Hillgren, K.M., Hoffmaster, K.A., Ishikawa, T., Keppler, D., Kim, R.B., Lee, C.A., Niemi, M., Polli, J.W., Sugiyama, Y., Swaan, P.W., Ware, J.A., Wrighton, S.A., Xiao, G., Yee, S.W., Zamek-Gliszczynski, M.J., Zhang, L., 2010. Membrane transporters in drug development. *Nat. Rev. Drug Discov.* 9, 215–236. <https://doi.org/10.1038/nrd3028>.
- Guha, R., 2007. Chemical informatics functionality in R. *J. Stat. Softw.* 18. <https://doi.org/10.18637/jss.v018.i05>.
- Grube, M., Kock, K., Oswald, S., Draber, K., Meissner, K., Eckel, L., Böhm, M., Felix, S., Vogelgesang, S., Jedlitschky, G., 2006. Organic anion transporting polypeptide 2B1 is a high-affinity transporter for atorvastatin and is expressed in the human heart. *Clin. Pharmacol. Ther.* 80, 607–620. <https://doi.org/10.1016/j.clpt.2006.09.010>.
- Harper, C.R., Jacobson, T.A., 2007. The broad spectrum of statin myopathy: from myalgia to rhabdomyolysis. *Curr. Opin. Lipidol.* 18, 401–408. <https://doi.org/10.1097/MOL.0b013e32825a6773>.
- Hirano, M., Maeda, K., Shitara, Y., Sugiyama, Y., 2006. Drug-drug interaction between pitavastatin and various drugs via OATP1B1. *Drug Metab. Dispos.* 34, 1229–1236. <https://doi.org/10.1124/dmd.106.009290>.
- Ho, R.H., Tirona, R.G., Leake, B.F., Glaeser, H., Lee, W., Lemke, C.J., Wang, Y., Kim, R.B., 2006. Drug and bile acid transporters in rosuvastatin hepatic uptake: function, expression, and pharmacogenetics. *Gastroenterol* 130, 1793–1806. <https://doi.org/10.1053/j.gastro.2006.02.034>.
- Hoste, E., Paquot, A., Panin, N., Horion, S., El Hamdaoui, H., Muccioli, G.G., Haufroid, V., Elens, L., 2023. Genetic polymorphisms in *SLCO2B1* and *ABCC1* jointly modulate atorvastatin intracellular accumulation in HEK293 recombinant cell lines. *Ther. Drug Monit.* 45, 400–408. <https://doi.org/10.1097/FTD.0000000000001043>.
- Jedlitschky, G., Burchell, B., Keppler, D., 2000. The multidrug resistance protein 5 functions as an ATP-dependent export pump for cyclic nucleotides. *J. Biol. Chem.* 275, 30069–30074. <https://doi.org/10.1074/jbc.M005463200>.
- Jedlitschky, G., Leier, I., Buchholz, U., Barnouin, K., Kurz, G., Keppler, D., 1996. Transport of glutathione, glucuronate, and sulfate conjugates by the MRP gene-encoded conjugate export pump. *Cancer Res* 56, 988–994.
- Kanamitsu, K., Kusahara, H., Schuetz, J.D., Takeuchi, K., Sugiyama, Y., 2017. Investigation of the importance of multidrug resistance-associated protein 4 (Mrp4/Abcc4) in the active efflux of anionic drugs across the blood–brain barrier. *J. Pharm. Sci.* 106, 2566–2575. <https://doi.org/10.1016/j.xphs.2017.04.040>.
- Kanda, K., Takahashi, R., Yoshikado, T., Sugiyama, Y., 2018. Total hepatocellular disposition profiling of rosuvastatin and pitavastatin in sandwich-cultured human hepatocytes. *Drug Metab. Pharmacokinet.* 33, 164–172. <https://doi.org/10.1016/j.dmpk.2018.04.001>.
- Karlgren, M., Vildhede, A., Norinder, U., Wisniewski, J.R., Kimoto, E., Lai, Y., Haglund, U., Artursson, P., 2012. Classification of inhibitors of hepatic organic anion transporting polypeptides (OATPs): influence of protein expression on drug–drug interactions. *J. Med. Chem.* 55, 4740–4763. <https://doi.org/10.1021/jm300212s>.
- Keskitalo, J.E., Pasanen, M.K., Neuvonen, P.J., Niemi, M., 2009a. Different effects of the *ABCG2* C.421C>A SNP on the pharmacokinetics of fluvastatin, pravastatin and simvastatin. *Pharmacogenomics* 10, 1617–1624. <https://doi.org/10.2217/pgs.09.85>.
- Keskitalo, J.E., Zolk, O., Fromm, M.F., Kurkinen, K.J., Neuvonen, P.J., Niemi, M., 2009b. *ABCG2* polymorphism markedly affects the pharmacokinetics of atorvastatin and rosuvastatin. *Clin. Pharmacol. Ther.* 86, 197–203. <https://doi.org/10.1038/clpt.2009.79>.
- Kikutani, Y., Kobayashi, M., Konishi, T., Sasaki, S., Narumi, K., Furugen, A., Takahashi, N., Iseki, K., 2016. Involvement of monocarboxylate transporter 4 expression in statin-induced cytotoxicity. *J. Pharm. Sci.* 105, 1544–1549. <https://doi.org/10.1016/j.xphs.2016.01.014>.
- Kitamura, S., Maeda, K., Wang, Y., Sugiyama, Y., 2008. Involvement of multiple transporters in the hepatobiliary transport of rosuvastatin. *Drug Metab. Dispos.* 36, 2014–2023. <https://doi.org/10.1124/dmd.108.021410>.
- Knauer, M.J., Urquhart, B.L., Meyer Zu Schwabedissen, H.E., Schwarz, U.I., Lemke, C.J., Leake, B.F., Kim, R.B., Tirona, R.G., 2010. Human skeletal muscle drug transporters determine local exposure and toxicity of statins. *Circ. Res.* 106, 297–306. <https://doi.org/10.1161/CIRCRESAHA.109.203596>.
- Kobayashi, M., Otsuka, Y., Itagaki, S., Hirano, T., Iseki, K., 2006. Inhibitory effects of statins on human monocarboxylate transporter 4. *Int. J. Pharm.* 317, 19–25. <https://doi.org/10.1016/j.ijpharm.2006.02.043>.
- Koepsell, H., 2013. The SLC22 family with transporters of organic cations, anions and zwitterions. *Mol. Aspects Med.* 34, 413–435. <https://doi.org/10.1016/j.mam.2012.10.010>.
- Kopplow, K., Letschert, K., König, J., Walter, B., Keppler, D., 2005. Human hepatobiliary transport of organic anions analyzed by quadruple-transfected cells. *Mol. Pharmacol.* 68, 1031–1038. <https://doi.org/10.1124/mol.105.014605>.
- Law, M., Rudnicka, A.R., 2006. Statin safety: a systematic review. *Am. J. Cardiol.* 97, S52–S60. <https://doi.org/10.1016/j.amjcard.2005.12.010>.
- Lehtisalo, M., Taskinen, S., Tarkiainen, E.K., Neuvonen, M., Viinamäki, J., Paile-Hyvärinen, M., Lilius, T.O., Tapaninen, T., Backman, J.T., Tornio, A., Niemi, M., 2023. A comprehensive pharmacogenomic study indicates roles for *SLCO1B1*, *ABCG2* and *SLCO2B1* in rosuvastatin pharmacokinetics. *Br. J. Clin. Pharmacol.* 89, 242–252. <https://doi.org/10.1111/bcp.15485>.
- Lönnerberg, K.L., Tornio, A., Hirvensalo, P., Keskitalo, J., Mustaniemi, A.-L., Kiiski, J.I., Filppula, A.M., Niemi, M., 2023. Real-world pharmacogenetics of statin intolerance: Effects of *SLCO1B1*, *ABCG2*, and *CYP2C9* variants. *Pharmacogenet. Genomics* 33, 153–160. <https://doi.org/10.1097/FPC.0000000000000504>.
- Lotta, L.A., Sharp, S.J., Burgess, S., Perry, J.R.B., Stewart, I.D., Willems, S.M., Luan, J., Ardanaz, E., Arriola, L., Balkau, B., Boeing, B., Deloukas, P., Forouhi, N.G., Franks, P.W., Grioni, S., Kaaks, R., Key, T.J., Navarro, C., Nilsson, P.M., Overvad, K., Palli, D., Panico, S., Quirós, J.-R., Riboli, E., Rolandsson, O., Sacerdote, C., Salamanca-Fernandez, E., Slimani, N., Spijkerman, A.M.W., Tjønneland, A., Tumino, R., Van Der, A., 2016. Association between low-density lipoprotein cholesterol-lowering genetic variants and risk of type 2 diabetes: a meta-analysis. *JAMA* 316, 1383. <https://doi.org/10.1001/jama.2016.14568>.
- Meyer Zu Schwabedissen, H.E.U., Grube, M., Heydrich, B., Linnemann, K., Fusch, C., Kroemer, H.K., Jedlitschky, G., 2005. Expression, localization, and function of MRP5 (ABCC5), a transporter for cyclic nucleotides, in human placenta and cultured human trophoblasts. *Am. J. Pathol.* 166, 39–48.
- Mykkänen, A.J.H., Tarkiainen, E.K., Taskinen, S., Neuvonen, M., Paile-Hyvärinen, M., Lilius, T.O., Tapaninen, T., Klein, K., Schwab, M., Backman, J.T., Tornio, A., Niemi, M., 2024. Genome-wide association study of atorvastatin pharmacokinetics: associations with *SLCO1B1*, *UGT1A3*, and *LPP*. *Clin. Pharmacol. Ther.* 115 (6), 1428–1440. <https://doi.org/10.1002/cpt.3236>.
- Mykkänen, A.J.H., Taskinen, S., Neuvonen, M., Paile-Hyvärinen, M., Tarkiainen, E.K., Lilius, T., Tapaninen, T., Backman, J.T., Tornio, A., Niemi, M., 2022. Genome-wide association study of simvastatin pharmacokinetics. *Clin. Pharmacol. Ther.* 112, 676–686. <https://doi.org/10.1002/cpt.2674>.
- Neuvonen, P.J., Niemi, M., Backman, J.T., 2006. Drug interactions with lipid-lowering drugs: mechanisms and clinical relevance. *Clin. Pharmacol. Ther.* 80, 565–581. <https://doi.org/10.1016/j.clpt.2006.09.003>.
- Niemi, M., Pasanen, M.K., Neuvonen, P.J., 2011. Organic anion transporting polypeptide 1B1: a genetically polymorphic transporter of major importance for hepatic drug uptake. *Pharmacol. Rev.* 63, 157–181. <https://doi.org/10.1124/pr.110.002857>.
- Nies, A.T., Jedlitschky, G., König, J., Herold-Mende, C., Steiner, H.H., Schmitt, H.-P., Keppler, D., 2004. Expression and immunolocalization of the multidrug resistance proteins, MRP1–MRP6 (ABCC1–ABCC6), in human brain. *Neuroscience* 129, 349–360. <https://doi.org/10.1016/j.neuroscience.2004.07.051>.
- Noé, J., Portmann, R., Brun, M.E., Funk, C., 2007. Substrate-dependent drug–drug interactions between gemfibrozil, fluvastatin and other organic anion-transporting peptide (OATP) substrates on OATP1B1, OATP2B1, and OATP1B3. *Drug Metab. Dispos.* 35, 1308–1314. <https://doi.org/10.1124/dmd.106.012930>.
- Nozawa, T., Imai, K., Nezu, J.-I., Tsuji, A., Tamai, I., 2004. Functional characterization of pH-sensitive organic anion transporting polypeptide OATP-B in human. *J. Pharmacol. Exp. Ther.* 308, 438–445. <https://doi.org/10.1124/jpet.103.060194>.
- Ölander, M., Wiśniewski, J.R., Matsson, P., Lundquist, P., Artursson, P., 2016. The proteome of filter-grown Caco-2 cells with a focus on proteins involved in drug disposition. *J. Pharm. Sci.* 105, 817–827. <https://doi.org/10.1016/j.xphs.2015.10.030>.
- Pál, Á., Méhn, D., Molnár, É., Gedey, S., Mészáros, P., Nagy, T., Glavinas, H., Janáky, T., Von Richter, O., Báthori, G., Sente, L., Krajcsi, P., 2007. Cholesterol potentiates *ABCG2* activity in a heterologous expression system: improved in vitro model to study function of human *ABCG2*. *J. Pharmacol. Exp. Ther.* 321, 1085–1094. <https://doi.org/10.1124/jpet.106.119289>.
- Pfeifer, N.D., Yang, K., Brouwer, K.L.R., 2013. Hepatic basolateral efflux contributes significantly to rosuvastatin disposition I: characterization of basolateral versus biliary clearance using a novel protocol in sandwich-cultured hepatocytes. *J. Pharmacol. Exp. Ther.* 347, 727–736. <https://doi.org/10.1124/jpet.113.207472>.
- Pilegaard, H., Terzis, G., Halestrap, A., Juel, C., 1999. Distribution of the lactate/H⁺ transporter isoforms MCT1 and MCT4 in human skeletal muscle. *Am. J. Physiol.* 276, E843–E848. <https://doi.org/10.1152/ajpendo.1999.276.5.E843>.
- Prieto Garcia, L., Vildhede, A., Nordell, P., Ahlström, C., Montaser, A.B., Terasaki, T., Lennernäs, H., Sjögren, E., 2024. Physiologically based pharmacokinetics modeling and transporter proteomics to predict systemic and local liver and muscle disposition

- of statins. *CPT Pharmacometrics Syst. Pharmacol.* 13, 1029–1043. <https://doi.org/10.1002/psp4.13139>.
- R Core Team, 2024. R: A language and environment for statistical computing. R Found. Stat. Comput., Vienna, Austria. <https://www.R-project.org/>.
- Renes, J., De Vries, E.G.E., Nienhuis, E.F., Jansen, P.L.M., Müller, M., 1999. ATP- and glutathione-dependent transport of chemotherapeutic drugs by the multidrug resistance protein MRP1. *Br. J. Pharmacol.* 126, 681–688. <https://doi.org/10.1038/sj.bjp.0702360>.
- Roth, G.A., Mensah, G.A., Johnson, C.O., Addolorato, G., Ammirati, E., Baddour, L.M., Barengo, N.C., Beaton, A.Z., Benjamin, E.J., Benziger, C.P., Bonny, A., Brauer, M., Brodmann, M., Cahill, T.J., Carapetis, J., Catapano, A.L., Chugh, S.S., Cooper, L.T., Coresh, J., Criqui, M.H., Ezekowitz, M.D., Fernandez, R.G., Gakidou, T., Guerrero, E., Habib, R., Narula, S.H., Neal, J., Patel, B., Sliwa, R.M., Thrift, K., Vaduganathan, A. G., Vos, M., Watkins, T., Zuhlke, D.A., Murray, L.J., Fuster, C.J.L., 2020. Global burden of cardiovascular diseases and risk factors, 1990–2019. *J. Am. Coll. Cardiol.* 76, 2982–3021. <https://doi.org/10.1016/j.jacc.2020.11.010>.
- Sáfar, Z., Kecskeméti, G., Molnár, J., Kurunczi, A., Szabó, Z., Janáky, T., Kis, E., Krajcsi, P., 2021. Inhibition of ABCG2/BCRP-mediated transport–correlation analysis of various expression systems and probe substrates. *Eur. J. Pharm. Sci.* 156, 105593. <https://doi.org/10.1016/j.ejps.2020.105593>.
- Sampath, J., Adachi, M., Hatse, S., Naesens, L., Balzarini, J., Flatley, R., Matherly, L., Schuetz, J., 2002. Role of MRP4 and MRP5 in biology and chemotherapy. *AAPS J* 4, 22–30. <https://doi.org/10.1208/ps040314>.
- Sattar, N., Preiss, D., Murray, H.M., Welsh, P., Buckley, B.M., de Craen, A.J., Seshasai, S. R.K., McMurray, J.J., Freeman, D.J., Jukema, J.W., Macfarlane, P.W., Packard, C.J., Stott, D.J., Westendorp, R.G., Shepherd, J., Davis, B.R., Pressel, S.L., Marchioni, R., Marfisi, R.M., Maggioni, A.P., Tavazzi, L., Tognoni, G., Kjekshus, J., Pedersen, T.R., Cook, T.J., Gotto, A.M., Clearfield, M.B., Downs, J.R., Nakamura, H., Ohashi, Y., Mizuno, K., Ray, K.K., Ford, I., 2010. Statins and risk of incident diabetes: a collaborative meta-analysis of randomised statin trials. *Lancet* 375, 735–742. [https://doi.org/10.1016/S0140-6736\(09\)61965-6](https://doi.org/10.1016/S0140-6736(09)61965-6).
- Schirris, T.J.J., Renkema, G.H., Ritschel, T., Voermans, N.C., Bilos, A., van Engelen, B.G. M., Brandt, U., Koopman, W.J.H., Beyrath, J.D., Rodenburg, R.J., Willems, P.H.G.M., Smeitink, J.A.M., Russel, F.G.M., 2015. Statin-induced myopathy is associated with mitochondrial complex iii inhibition. *Cell Metab.* 22, 399–407. <https://doi.org/10.1016/j.cmet.2015.08.002>.
- Schuetz, J.D., Connelly, M.C., Sun, D., Paibir, S.G., Flynn, P.M., Srinivas, R.V., Kumar, A., Fridland, A., 1999. MRP4: a previously unidentified factor in resistance to nucleoside-based antiviral drugs. *Nat. Med.* 5, 1048–1051. <https://doi.org/10.1038/12487>.
- Sirvent, P., Bordenave, S., Vermaelen, M., Roels, B., Vassort, G., Mercier, J., Raynaud, E., Lacampagne, A., 2005. Simvastatin induces impairment in skeletal muscle while heart is protected. *Biochem. Biophys. Res. Commun.* 338, 1426–1434. <https://doi.org/10.1016/j.bbrc.2005.10.108>.
- Sjöstedt, N., Salminen, T.A., Kidron, H., 2019. Endogenous, cholesterol-activated ATP-dependent transport in membrane vesicles from *Spodoptera frugiperda* cells. *Eur. J. Pharm. Sci.* 137, 104963. <https://doi.org/10.1016/j.ejps.2019.104963>.
- Skottheim, I.B., Gedde-Dahl, A., Hejazifar, S., Hoel, K., Åsberg, A., 2008. Statin induced myotoxicity: the lactone forms are more potent than the acid forms in human skeletal muscle cells in vitro. *Eur. J. Pharm. Sci.* 33, 317–325. <https://doi.org/10.1016/j.ejps.2007.12.009>.
- Stage, T.B., Mortensen, C., Khalaf, S., Steffensen, V., Hammer, H.S., Xiong, C., Nielsen, F., Poetz, O., Svenningsen, Å.F., Rodriguez-Antona, C., Kroetz, D.L., 2020. P-glycoprotein inhibition exacerbates paclitaxel neurotoxicity in neurons and patients with cancer. *Clin. Pharmacol. Ther.* 108, 671–680. <https://doi.org/10.1002/cpt.1847>.
- St-Pierre, M.V., Serrano, M.A., Macias, R.I.R., Dubs, U., Hoechli, M., Lauper, U., Meier, P. J., Marin, J.J.G., 2000. Expression of members of the multidrug resistance protein family in human term placenta. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 279, R1495–R1503. <https://doi.org/10.1152/ajpregu.2000.279.4.R1495>.
- Turner, R.M., Pirmohamed, M., 2019. Statin-related myotoxicity: a comprehensive review of pharmacokinetic, pharmacogenomic and muscle components. *J. Clin. Med.* 9, 22. <https://doi.org/10.3390/jcm9010022>.
- Varma, M.V., Rotter, C.J., Chupka, J., Whalen, K.M., Duignan, D.B., Feng, B., Litchfield, J., Goosen, T.C., El-Kattan, A.F., 2011. pH-Sensitive interaction of HMG-CoA reductase inhibitors (Statins) with organic anion transporting polypeptide 2B1. *Mol. Pharm.* 8, 1303–1313. <https://doi.org/10.1021/mp200103h>.
- Vasilioi, V., Vasilioi, K., Nebert, D.W., 2008. Human ATP-binding cassette (ABC) transporter family. *Hum. Genom.* 3, 281. <https://doi.org/10.1186/1479-7364-3-3-281>.
- Voorra, D., Shah, S.H., Spasojevic, I., Ali, S., Reed, C.R., Salisbury, B.A., Ginsburg, G.S., 2009. The SLCO1B1*5 genetic variant is associated with statin-induced side effects. *J. Am. Coll. Cardiol.* 54, 1609–1616. <https://doi.org/10.1016/j.jacc.2009.04.053>.
- Yebo, H.G., Aschmann, H.E., Kaufmann, M., Puhani, M.A., 2019. Comparative effectiveness and safety of statins as a class and of specific statins for primary prevention of cardiovascular disease: A systematic review, meta-analysis, and network meta-analysis of randomized trials with 94,283 participants. *Am. Heart J.* 210, 18–28. <https://doi.org/10.1016/j.ahj.2018.12.007>.