



UNIVERSITY OF HELSINKI



<https://helda.helsinki.fi>

Helda

---

# Metformin and Combined Oral Contraceptive Pills in the Management of Polycystic Ovary Syndrome: A Systematic Review and Meta-analysis

Melin, Johanna

Endocrine Society

2024-02

---

Melin, J, Forslund, M, Alesi, S, Piltonen, T, Romualdi, D, Spritzer, P M, Tay, C T, Pena, A, Witchel, S F, Mousa, A & Teede, H 2024, 'Metformin and Combined Oral Contraceptive Pills in the Management of Polycystic Ovary Syndrome: A Systematic Review and Meta-analysis', *Journal of Clinical Endocrinology and Metabolism*, vol. 109, no. 2, pp. e817-e836. <https://doi.org/10.1210/clinem/dga>

---

<http://hdl.handle.net/10138/574674>

[10.1210/clinem/dgad465](https://doi.org/10.1210/clinem/dgad465)

---

cc\_by\_nc\_nd

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

# Metformin and Combined Oral Contraceptive Pills in the Management of Polycystic Ovary Syndrome: A Systematic Review and Meta-analysis

Johanna Melin,<sup>1,2,\*</sup> Maria Forslund,<sup>1,3,\*</sup> Simon Alesi,<sup>1</sup> Terhi Piltonen,<sup>4</sup> Daniela Romualdi,<sup>5</sup> Poli Mara Spritzer,<sup>6</sup> Chau Thien Tay,<sup>1</sup> Alexia Pena,<sup>7</sup> Selma Feldman Witchel,<sup>8</sup> Aya Mousa,<sup>1,\*</sup> and Helena Teede<sup>1,9,\*</sup>

<sup>1</sup>Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Clayton, Victoria 3168, Australia

<sup>2</sup>Department of Obstetrics and Gynecology, University of Helsinki, Helsinki University Hospital, 00014 Helsinki, Finland

<sup>3</sup>Department of Obstetrics and Gynecology, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, 40530 Gothenburg, Sweden

<sup>4</sup>Department of Obstetrics and Gynecology, Research Unit of Clinical Medicine and Medical Research Centre Oulu, Oulu University Hospital, University of Oulu, 90014 Oulu, Finland

<sup>5</sup>Department of Woman and Child Health and Public Health, Fondazione Policlinico Universitario Agostino Gemelli, IRCCS, 00168 Rome, Italy

<sup>6</sup>Gynecological Endocrinology Unit, Division of Endocrinology, Hospital de Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS 90035-003, Brazil

<sup>7</sup>Discipline of Paediatrics, The University of Adelaide and Robinson Research Institute, Adelaide, South Australia 5005, Australia

<sup>8</sup>Division of Pediatric Endocrinology, Department of Pediatrics, UPMC Children's Hospital of Pittsburgh, Pittsburgh, PA 15224, USA

<sup>9</sup>Endocrine and Diabetes Units, Monash Health, Clayton, Victoria 3169, Australia

**Correspondence:** Johanna Melin, MD, PhD, Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Level 1, 43-51 Kanooka Grove, Clayton, Victoria 3168, Australia. Email: [johanna.melin@hus.fi](mailto:johanna.melin@hus.fi).

\*Equal contributions as first and senior authors.

## Abstract

**Context:** Polycystic ovary syndrome (PCOS) affects more than 1 in 10 women.

**Objective:** As part of the 2023 International PCOS Guidelines update, comparisons between combined oral contraceptive pills (COCP), metformin, and combination treatment were evaluated.

**Data Sources:** Ovid Medline, Embase, PsycINFO, All EBM, and CINAHL were searched.

**Study Selection:** Women with PCOS included in randomized controlled trials (RCTs).

**Data Extraction:** We calculated mean differences and 95% CIs regarding anthropometrics, metabolic, and hyperandrogenic outcomes. Meta-analyses and quality assessment using GRADE were performed.

**Data Synthesis:** The search identified 1660 publications; 36 RCTs were included. For hirsutism, no differences were seen when comparing metformin vs COCP, nor when comparing COCP vs combination treatment with metformin and COCP. Metformin was inferior on free androgen index (FAI) (7.08; 95% CI 4.81, 9.36), sex hormone binding globulin (SHBG) (−118.61 nmol/L; 95% CI −174.46, −62.75) and testosterone (0.48 nmol/L; 95% CI 0.32, 0.64) compared with COCP. COCP was inferior for FAI (0.58; 95% CI 0.36, 0.80) and SHBG (−16.61 nmol/L; 95% CI −28.51, −4.71) compared with combination treatment, whereas testosterone did not differ. Metformin lowered insulin (−27.12 pmol/L; 95% CI −40.65, −13.59) and triglycerides (−0.15 mmol/L; 95% CI −0.29, −0.01) compared with COCP. COCP was inferior for insulin (17.03 pmol/L; 95% CI 7.79, 26.26) and insulin resistance (0.44; 95% CI 0.17, 0.70) compared with combination treatment.

**Conclusions:** The choice of metformin or COCP treatment should be based on symptoms, noting some biochemical benefits from combination treatment targeting both major endocrine disturbances seen in PCOS (hyperinsulinemia and hyperandrogenism).

**Key Words:** polycystic ovary syndrome, metformin, combined oral contraceptive pill, meta-analysis, hirsutism, weight

Polycystic ovary syndrome (PCOS) affects approximately 10% of women globally (1) and is characterized by chronic anovulation, oligo/amenorrhea, hyperandrogenism, polycystic ovarian morphology, and metabolic and psychological manifestations. PCOS is associated with hyperinsulinemia, insulin resistance, and increased body mass index (BMI). Higher

BMI can increase risks of gestational diabetes, type 2 diabetes, and cardiovascular disease (2, 3, 4). Insulin enhances ovarian and adrenal androgen secretion, exaggerating hyperandrogenic symptoms in PCOS (5). Androgen excess favors visceral fat deposition and lipid dysfunction with a vicious cycle of worsening of hyperinsulinemia and insulin resistance (6). To

break the vicious cycle, both insulin resistance and hyperandrogenism are key treatment targets in PCOS.

Metformin is an insulin-sensitizing drug, known to inhibit gluconeogenesis and lipogenesis (7) and prevent weight gain by influencing appetite regulatory pathways in the brain (8). Several studies have shown that metformin may improve biochemical hyperandrogenism and anovulation (9, 10). Generally combined oral contraceptive pills (COCP) are regarded as more effective than metformin for the management of hyperandrogenism and menstrual regulation in PCOS (11, 12).

COCP, containing estrogen and progestin components, are primarily used as contraception. Due to their direct endometrial effects, COCPs are also used among women with PCOS to regulate the menstrual cycle. A recent systematic review by our group found COCP to be effective in improving menstrual cyclicity and hirsutism in PCOS, with limited evidence for hirsutism (13). COCPs containing cyproterone acetate (CPA) might be more effective in reducing hirsutism compared to conventional COCPs (14, 15); however, the use of COCPs containing CPA is not recommended as first-line treatment due to the increased risk of venous thromboembolism in the general population (12, 16). Overall, finding the most favorable medical treatment for PCOS remains challenging. To treat insulin resistance, hyperandrogenism, irregular cycles, and metabolic features, a medical treatment combining COCP and metformin has been suggested (17).

We performed a systematic review and meta-analysis of the most up-to-date evidence regarding metformin and COCP in the management of clinical and hormonal outcomes in women with PCOS. Results will directly inform the 2023 International Evidence-based Guideline for the Assessment and Management of PCOS. Subgroup analyses based on BMI and age (adolescent and adult), as well as type of COCP (with or without CPA) were performed to further inform recommendations regarding the optimal treatment for the individual patient.

## Materials and Methods

This systematic review and meta-analysis was conducted as part of the 2023 update of the International Evidence-based Guidelines for the Assessment and Management of PCOS (18). We addressed the efficacy of (i) metformin compared to COCP; (ii) COCP monotherapy compared with metformin and COCP; and (iii) metformin monotherapy compared with metformin and COCP in women with PCOS for improving anthropometric, biochemical, clinical, and psychological outcomes.

### Selection Criteria

The Population, Intervention, Comparison and Outcome (PICO) framework for this systematic review is outlined in Table 1 and was determined by an experienced, multidisciplinary, clinical research team (A.M., T.P., D.R., P.M.S., C.T.T., A.P., S.W., H.T.). Core outcomes were based on a Delphi process involving 700 clinicians, academic opinion leaders, and consumers (19). Outcomes included anthropometric, metabolic, androgenicity, and psychological outcomes, as well as adverse events (detailed in Table 1).

### Data Sources

This systematic review and meta-analysis provide an update of a previous systematic review (12) conducted in 2017 to inform

**Table 1. Population, intervention, comparison and outcome (PICO) of the systematic review and meta-analysis**

PICO elements	Keywords
Patients	Females with PCOS (diagnosed by Rotterdam, NIH, or AES) of any ethnicity and weight. Subgroups: Adolescents (10-19 years), adults, BMI < 25 kg/m <sup>2</sup> , BMI ≥ 25 kg/m <sup>2</sup> Exclusion criteria: Females without PCOS. Less than 2 years post menarche. Women with type 2 diabetes, comorbidities, or major depression
Intervention	All types of COCPs, metformin, or COCP + metformin For hirsutism, at least 6 months treatment, for other outcomes at least 3 months treatment. Subgroups: Progestins based on different androgenicity (containing or not containing CPA) Exclusion criteria: Nonoral formulation of contraceptives
Comparisons	As stated in intervention
Outcomes	Anthropometric: Weight, BMI, WHR Androgenicity: Hirsutism (FG score), FAI, testosterone and SHBG, DHEAS, androstenedione, irregular cycles Metabolic: Insulin resistance (HOMA-IR, Clamp, OGTT), cholesterol, LDL, HDL, triglycerides, CRP, PAI-1 levels Psychological: QoL, depression Adverse effects

Abbreviations: BMI, body mass index; COCP, combined oral contraceptive pills; CPA, cyproterone acetate; CRP, C-reactive protein; DHEAS, dehydroepiandrosterone sulfate; FAI, free androgen index; FG score, Ferriman Gallwey score; HDL, high-density lipoprotein; HOMA-IR, homeostatic model assessment for insulin resistance; LDL, low-density lipoprotein; OGTT, oral glucose tolerance test; PAI-1, plasminogen activator inhibitor; PCOS, polycystic ovary syndrome; QoL, quality of life; WHR, waist to hip ratio; SHBG, sex hormone binding globulin.

the International Evidence-based Guidelines for the Assessment and Management of PCOS (20). This review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (21, 22). The study protocol was registered prior to full-text screening in PROSPERO (Registration number CRD42022345640).

Ovid Medline, Embase, PsycINFO, All EBM, and CINAHL were searched to identify relevant literature from 2017 until July 7, 2022. We also re-evaluated all articles, which were found in the search performed during the previous systematic review ranging from 1946 to 2017, as well as additional references identified from relevant systematic reviews. Details of the search strategy are presented in Supplementary Fig. S1 (23).

### Study Selection

Two authors (J.M., M.F., or S.A.) independently screened each potential study on title and abstract with the use of COVIDENCE. The same authors performed the full-text screening to determine the final included studies. If there was any doubt about inclusion, the study was reviewed and discussed in a larger group (J.M., M.F., S.A., and C.T.T.).

### Quality Appraisal of the Evidence

Quality appraisal of the included studies, in terms of risk of bias (RoB), was performed by J.M. and M.F. using RoB2 (24). Individual quality items included methods of

randomization and group allocation; blinding of patients, carers, investigators, or outcome assessors; methods of outcome assessment and reporting; methods of data analysis; statistical issues; and trial pre-registration. Disagreements were resolved by discussion and re-inspection of the full-text article. Each study was adjudged as having an overall low RoB, some concerns, or high RoB. We used the Grading of Recommendations, Assessment, Development and Evaluations (GRADE) method to estimate the certainty of the evidence (25). GRADE assessments were conducted in duplicate by J.M. and M.F.

## Data Extraction

Two independent reviewers (J.M. and M.F.) extracted data from included studies, using a specially developed data extraction form according to selection criteria. Extracted data included a description of the study (authors, country, year of publication, setting, diagnostic criteria for PCOS), participants (mean age and BMI), intervention (dose and duration of metformin and COCP) and study results according to outcome. All units were recalculated to SI units and SD.

## Statistical Analysis

Meta-analyses were performed using Review Manager 5.4.1. For continuous outcomes we calculated mean differences (MD) and 95% CI and for dichotomous outcomes odds ratios (ORs) and 95% CI. Due to clinical heterogeneity from differences in metformin dose, type of COCP, and duration of treatment, random effects models were used for the meta-analyses.

We performed subgroup analyses according to BMI, age (adults and adolescents), and type of COCP. Regarding BMI, studies were sorted into 3 categories; those with participants with a BMI < 25 kg/m<sup>2</sup>, those with a BMI ≥ 25 kg/m<sup>2</sup>, and studies using other BMI cutoffs (BMI not classified). Participants aged 10 to 19 years were classified as adolescents. Subgroup analyses according to type of COCP were classified into COCP containing CPA and other COCP.

Funnel plots were inspected for the assessment of publication bias.

## Results

### Study Selection

From 1660 search results, 450 articles were chosen for full-text review. In addition, we screened 176 full texts from the previous search in 2017 (12). After full-text review, 36 randomized controlled trials (RCTs) in 46 publications were identified (Fig. 1). Of these, 25 RCTs compared metformin with COCP, 17 compared COCP with metformin and COCP, and 6 compared metformin with metformin and COCP. Study characteristics are presented in Table 2. The RCTs were performed in Europe (n = 9) (26-40) North America (n = 4) (41-44), the Middle East (n = 14) (45-58), Asia (n = 8) (59-66), and Australia (n = 1) (67-70) between 2000 and 2021. Intervention durations ranged from 3 to 24 months, with 6 months of follow-up being the most commonly reported data. Metformin doses ranged from 500 mg to 2000 mg daily, with 1500 to 2000 mg daily being the most common dose (in 28 of the 36 studies). Regarding COCP, all studies used 30 to 35 µg ethinylestradiol (EE), 20 studies used CPA as the progestin compound, 14 studies used other progestins, and 2 studies had one study arm with participants using EE/drospirenone and another arm with those using EE/

CPA (26, 40). Four studies involved adolescents (41, 42, 44, 49) and the remainder focused on adults.

### Risk of Bias

In our systematic review, 2 studies fulfilled the criteria for low RoB, 14 had some concerns, and 20 had a high RoB. All studies adequately described the PCOS diagnostic criteria and inclusion and exclusion criteria. The most common reason for a high RoB was bias arising from the randomization process. The second most common reason was missing outcome data. RoB assessments of the included studies are shown in Fig. 2.

### Meta-analyses

Results on metformin compared with COCP are shown in Table 3. Subgroup analyses are shown in Supplementary Tables S1 and S2 (23), also illustrated in Supplementary Fig. S2 (subgroups according to BMI) (23), Supplementary Fig. S3 (subgroups according to adults/adolescents) (23), and Supplementary Fig. S4 (subgroups according to COCP with or without CPA) (23).

Combination treatment (COCP and metformin) compared with COCP monotherapy is presented in Table 4, with subgroup analysis reported in Supplementary Table S3 and Supplementary Fig. S5 (23). Results on combination treatment vs metformin monotherapy are presented in Table 5, with subgroup analysis in Supplementary Table S4 and Supplementary Fig. S6 (23).

We found no evidence of publication bias in the funnel plots.

### Anthropometric Outcomes

#### Metformin vs COCP

Regarding overall results on weight, BMI, and waist to hip ratio (WHR), no differences were found (Table 3).

In subgroup analyses according to BMI and adults/adolescents no differences were seen on weight, BMI and WHR (Supplementary Table S1; Supplementary Figs. S2 and S3) (23).

In subgroup analyses comparing metformin with COCP with and without CPA (Supplementary Table S2; Supplementary Fig. S4) (23) there were no differences in weight. Nevertheless, BMI was lower with metformin when compared with EE/CPA (MD -0.99 kg/m<sup>2</sup>; 95% CI, -1.74 to -0.23), but not when compared with COCP without CPA.

#### COCP vs combination

COCP alone compared with combination treatment with COCP and metformin showed no differences in weight, WHR, or BMI (Table 4).


#### Metformin vs combination

Metformin was superior in lowering WHR (MD -0.03; 95% CI, -0.06 to -0.01) compared with metformin, with no difference in BMI (Table 5).

### Clinical and Biochemical Hyperandrogenism

#### Metformin vs COCP

No difference was seen in hirsutism in the overall analysis but regarding biochemical hyperandrogenism, metformin was



**Figure 1.** Included studies are shown in the PRISMA flowchart. The search was performed for 3 medical treatments (combined oral contraceptive pills [COCP], metformin, and anti-androgens) as part of the update of the PCOS guidelines. Results comparing COCP, metformin, and combined COCP and metformin were included in this systematic review.

inferior to COCP on free androgen index (FAI) (MD 7.08; 95% CI, 4.81 to 9.36), sex hormone binding globulin (SHBG) (MD  $-118.61$  nmol/L; 95% CI,  $-174.46$  to  $-62.75$ ), and testosterone (MD  $0.48$  nmol/L; 95% CI, 0.32 to 0.64) (Table 3).

Subgroup analysis categorized by BMI (Supplementary Table S1; Supplementary Fig. S2) (23) showed that in women with PCOS and a BMI  $< 25$  kg/m<sup>2</sup>, metformin was less effective than COCP in treating hirsutism (MD 1.73; 95% CI, 0.07 to 3.40) and improving FAI (MD 5.78; 95% CI, 2.82 to 8.73), total testosterone (MD  $0.56$  nmol/L; 95% CI, 0.29 to

0.83), and SHBG (MD  $-168$  nmol/L; 95% CI,  $-211$  to  $-124$ ).

In women with a BMI  $\geq 25$  kg/m<sup>2</sup>, there was no difference in hirsutism when comparing metformin with COCP. For biochemical hyperandrogenism, metformin was inferior for FAI (MD 9.05; 95% CI, 6.44 to 11.66); total testosterone (MD  $0.40$  nmol/L; 95% CI, 0.15 to 0.66), and SHBG (MD  $-96$  nmol/L; 95% CI,  $-121$  to  $-72$ ).

The type of COCP did not influence the overall results regarding clinical and biochemical hyperandrogenism (Supplementary Table S2; Supplementary Fig. S4) (23).

**Table 2. Characteristics of included studies**

Study ID	RoB	Comparisons	Country, duration	N	Mean age (years)	Mean BMI	PCOS	Menarche age	Smokers	Comments
Aghamohammadzadeh 2010	High	C: 35 µg EE + CPA 2 mg M: Metformin 1000 mg/day	Iran 3 + 6 months	C = 30 M = 30	All 23.4 ± 8.1	All 25.57 ± 5.4	NIH	NR	NR	Adults BMI not classified
Allen 2005	Mod	C: 35 µg EE + 0.25 mg NOR/day M: Metformin 1000 mg/day	USA 6 months	C = 15 M = 16	C: 15.3 (12.5-21) M: 15.4 (13.1-18.4)	C: 40.1 ± 2.1 M: 37.3 ± 1.3	Author defined			Adolescents BMI > 25
Al-Zubeidi 2015	High	C: 30 µg EE + 1 mg NOR/day M: Metformin 2000 mg/day	USA 6 months	C = 10 M = 12	C: 16 (15-17) M: 16 (14-18)	C: 33.4 ± 9 M: 33.7 ± 6	NIH	>2 years post menarche	NR	Adolescents BMI not classified
Bilgir 2009	High	C: 35 µg EE + 2 mg CPA C + M: 35 µg EE + 2 mg CPA + Metformin 1700 mg/day	Turkey 3 months	C = 20 C + M = 20	1: 24.3 ± 5.7 2: 25.2 ± 4.6	C: 28.2 ± 6.0 M: 28.2 ± 4.3	Rott	NR	Nonsmokers	Adults BMI not classified
Bodur 2018	High	C: 30 µg EE + 3 mg DRSP M: Metformin 1700 mg C + M: 30 µg EE + 3 mg DRSP + Metformin 1700 mg	Turkey 6 months	C = 17 M = 17 C + M = 12	1: 26.62 ± 4.92 2: 26.24 ± 3.96 3: 27.35 ± 5.65	C: 23.45 ± 3.40 M: 25.06 ± 3.08 C + M: 25.11 ± 3.75	Rott	NR	NR	Adults BMI not classified
Cetinkalp 2009	High	C: 35 µg EE + 2 mg CPA M: Metformin 2000 mg/day	Turkey 4 months	C = 33 M = 47	NR	C: 24.72 ± 4.1 M: 25.82 ± 6.1	Rott	NR	NR	Adults BMI not classified
Christakou 2014	Mod	C1: 35 µg EE + 2 mg CPA C2: 30 µg EE + 3 mg DRSP M: Metformin 1700 mg/day	Greece 3 + 6 months	C1 = 38 C2 = 36 M = 35	C1: 22 ± 0.6 C2: 23.2 ± 0.6 M: 21.5 ± 0.5	C1: 21.80 ± 0.4 C2: 22.37 ± 0.5 M: 23.03 ± 0.7	NIH	NR	All nonsmokers	Adults BMI < 25 EE/DRSP used in meta-analysis
Cibula 2005	Mod	C: 35 µg EE + 250 µg NOR 21/7 CM: 35 µg EE + 250 µg NOR + Metformin 1500 mg/day	Czech Republic 6 months	C = 15 CM = 13	C: 23.2 ± 4.6 CM: 23.8 ± 5.4	C: 22.1 ± 3.1 CM: 24.7 ± 4.9	Author defined			Adults BMI not classified
Dardzinska 2014	Mod	C: 35 µg EE + 2 mg CPA M: Metformin 1700 mg/day 2 months washout	Poland 4 months	Phase 1 C = 21 M = 14 Phase 2 C = 14 M = 7	C 1st: 24.9 [23.5;26.4] M 1st: 24.6 [23.5;26.4] [23.0;26.3]	C 1st: 24.9 ± 4.4 M 1st: 25.1 ± 9.8C 1st: 24.9 [23.5;26.4] M 1st: 24.6 [23.0;26.3]	Rott	NR	20% of all	Adults BMI < 25 Crossover

(continued)

Table 2. Continued

Study ID	RoB	Comparisons	Country, duration	N	Mean age (years)	Mean BMI	PCOS	Menarche age	Smokers	Comments
Dorgham 2021	High	1: only laser 2: laser + metformin 500 mg 3: laser + 35 µg EE + 2 mg CPA	Egypt 6 months (results available also for 3 months)	1 = 50 2 = 50 3 = 50	NR	NR	Rott	NR	NR	Adults BMI not classified Facial hirsutism required
El Maghraby 2015	High	C: 30 µg EE + 15 mg progesterin/day M: Metformin 1700 mg/day	Egypt 24 months (results available for 6, 12, 18, 24 m)	C = 33 M = 32	C: 16.90 ± 1.60 M: 17.20 ± 2.00	NR	Rott	NR	NR	Adolescents BMI not classified
Elter 2002	Mod	C: 35 µg EE + 2 mg CPA 21/7 C + M: 35 µg EE + 2 mg CPA + Metformin 1500 mg/day	Turkey 4 months	C = 20 C + M = 20	C: 23.45 ± 6.07 C + M: 24.90 ± 6.62	C: 21.83 ± 1.40 C + M: 22.74 ± 2.66	Author defined	NR	NR	Adults BMI < 25
Essah 2011	Low	C: 35 µg EE + 0.18/0.215/0.25 mg NOR (+ placebo) C + M: 35 µg EE + 0.18/0.215/0.25 mg NOR + Metformin 1500 mg/day	USA 3 months	C = 10 C + M = 9	NR (adults)	C: 32.6 ± 2.3 C + M: 36.2 ± 2.5	Rott	NR	All nonsmokers	Adults BMI not classified
Feng 2016	Mod	C: 35 µg EE + 2 mg CPA C + M: 35 µg EE + 2 mg CPA + Metformin 900-1700 mg/day	China 3 months	C = 41 C + M = 41	C: 28.57 ± 3.04 C + M: 27.86 ± 3.79	C: 27.77 ± 4.23 C + M: 29.46 ± 4.43	Rott	NR	NR	Adults BMI not classified
Fonseka 2020	Low	C1: EE/CPA (Diane-35) C2: EE/DES (Fermion) C1 + M: EE/CPA + Metformin (dose not specified) C2 + M: EE/DES + Metformin (dose not specified)	Sri Lanka 12 months	Implied C1 = 20 C2 = 23 C1 + M = 26 C2 + M = 30	C1: 23.35 ± 5.10 C2: 22.39 ± 6.45 C1 + M: 24.81 ± 6.24 C2 + M: 27.90 ± 6.89	C1: 28.27 ± 6.94 C2: 26.74 ± 4.88 C1 + M: 27.93 ± 4.89 C2 + M: 27.20 ± 4.28	Rott	NR	NR	Adults BMI not classified
Glintborg 2014a Glintborg 2014b Alinok 2018 Glintborg 2017	High	C: 30 µg EE + 150 mg DSG M: Metformin 2000 mg/day C + M: 30 µg EE + 150 mg DSG + Metformin 2000 mg/day	Denmark 12 months	C = 23 M = 19 C + M = 23	Median, 25-75%ile: C: 28 (24; 32) M: 31 (24; 33) C + M: 30 (24; 31)	Median, 25-75%ile: C: 28.0 (22.9; 31.8) M: 25.9 (24.1; 29.6) C + M: 27.6 (24.3; 31.2)	Rott	NR	NR	Adults BMI not classified

(continued)

**Table 2. Continued**

Study ID	RoB	Comparisons	Country, duration	N	Mean age (years)	Mean BMI	PCOS	Menarche age	Smokers	Comments
Harborne 2003	High	C: 35 µg EE + 2 mg CPA 21/7 M: Metformin 1500 mg/day	Scotland 12 months	C = 26 M = 26	C: 31.7 (26.8-36.5) M: 31.3 (27.9-34.7)	C: 31.8 (28.4-34.4) M: 31.7 (29.5-35.5)	Rott	NR	NR	Adults BMI > 2.5
Hoeger 2008	High	C: 30 µg EE + 0.15 mg DSG M: Metformin 1700 mg/day	USA 6 months	C = 10 M = 6	C: 15.4 ± 1.4 M: 15.4 ± 1.7	C: 37.8 ± 5.1 M: 36.1 ± 7.5	Rott	NR	All nonsmokers	Adolescents BMI > 2.5
Kaya 2012	High	C: 30 µg EE + 3 mg DRSP C + M: 30 µg EE + 3 mg DRSP + Metformin 1700 mg/day	Turkey 6 months	C = 19 C + M = 18	C: 23.2 ± 5.4 C + M: 23.0 ± 4.5	C: 26.4 ± 6.2 C + M: 31.7 ± 7.3	AES	NR	NR	Adults BMI not classified
Kaya 2015	High	C: 30 µg EE + 3 mg DRSP C + M: 30 µg EE + 3 mg DRSP + Metformin 1700 mg/day	Turkey 6 months	C = 25 C + M = 25	C: 23 ± 5 C + M: 24 ± 4	C: 26.7 ± 5.7 C + M: 29.8 ± 6.9	AES	NR	excluded	Adults BMI not classified
Kebapcilar 2009	High	C: 35 µg EE + 2 mg CPA C + M: 35 µg EE + 2 mg CPA + Metformin 1700 mg/day	Turkey 3 months	C = 22 C + M = 21	C: 24.1 ± 5.6 C + M: 25.1 ± 4.4	C: 27.2 ± 6.2 C + M: 28.7 ± 4.4	Rott	NR	excluded	Adults BMI not classified
Kebapcilar 2010	High	C: 35 µg EE + 2 mg CPA C + M: 35 µg EE + 2 mg CPA + Metformin 1700 mg/day M: Metformin 1700 mg/day	Turkey 3 months	12/group	24.0 ± 5.4 years; for all	C: 28.7 ± 6 C + M: 27.6 ± 3 M: 27.8 ± 4	Rott	NR	excluded	Adults BMI not classified
Kilic 2011	Mod	C: 30 µg EE + 0.15 mg DSG M: Metformin 1700 mg/day	Turkey 6 months	Obese C = 25 M = 24 Nonobese C = 24 M = 23	Obese C: 29.0 ± 3.5 M: 28.7 ± 3.7 Nonobese C: 26.7 ± 3.8 M: 26.3 ± 3.0	Obese C: 27.7 ± 0.9 M: 31.5 ± 2.2 Nonobese C: 21.6 ± 1.4 M: 23.3 ± 1.6	Rott	NR	excluded	Adults BMI < 2.5 BMI > 2.5
Kumar 2018	Mod	C: 35 µg EE/ + 2 mg CPA M: Metformin 2000 mg/day C + M: 35 µg EE + 2 mg CPA + Metformin 2000 mg/ day	India 6 months	C = 28 M = 30 C + M = 29	C: 22.9 (5) M: 22 (5.2) C + M: 24.1 (5.9)	C: 26.15 (4.9) M: 27.14 (6) C + M: 30.10 (5.5)	Rott	NR	NR	Adults BMI < 2.5
Luque-Ramírez 2009	Mod	C: 35 µg EE/2 mg CPA M: Metformin 1700 mg/day	Spain 6 months	C = 15 M = 19	C: 23.4 ± 5.6 M: 25.1 ± 6.6	C: 29.2 ± 5.7 M: 30.5 ± 6.9	NIH	NR	C: 40% M: 42%	Adults BMI not classified

(continued)

Table 2. Continued

Study ID	RoB	Comparisons	Country, duration	N	Mean age (years)	Mean BMI	PCOS	Menarche age	Smokers	Comments
Lv 2005	High	C: 35 µg EE/2 mg CPA C + M: 35 µg EE/2 mg CPA + Metformin 500 mg/day	China 6 months	C = 25 C + M = 25	C: 24.4 ± 5.1 C + M: 24.5 ± 5.6	C: 21.8 ± 1.4 C + M: 22.1 ± 20.2	Author defined	NR	NR	Adults BMI < 2.5
Meyer 2007	Mod	C1: 35 µg EE + 2 mg CPA (high) C2: 20 µg EE + 100 µg LVG + 50 mg SPL (low dose)*** M: Metformin 2000 mg/day	Australia 6 months	C1 = 31 C2 = 33 M = 36	Average: 31 years	C1: 36.5 no SD C2: 35.5 no SD M: 36.3 no SD	NIH	NR	Excluded	Adults BMI not classified
Burchall 2017										
Hutchison 2008										
Moran 2010										
Mhao 2016	High	C: EE 30 µg/CMA 2mg M: Metformin 1000 mg/day	Iraq 3 months	C = 10 M = 16	Age range, 14-40 years	C: 30.5 ± 5.3 M: 27.2 ± 5.4	NR	NR	NR	Adults BMI not classified
Morin-Papunen 2003a	High	C: 35 µg EE + 2 mg CPA 21/7	Finland 6 months	Nonobese: C = 10 M = 10 Obese: C = 10 M = 8	Nonobese: C: 28.5 ± 1.7 (SE) M: 28.2 ± 1.4 Obese: C: 29.8 ± 1.0 (SE) M: 29.9 ± 1.5	Nonobese: C: 21.8 ± 0.7 (SE) M 22.5 ± 0.8 Obese: C: 37.2 ± 1.8 (SE) M 32.5 ± 1.1	Aligns with Rott		Adults BMI < 2.5 BMI > 2.5	
Morin-Papunen 2000	High	M: Metformin 1000 mg/day for 3 months, then 2000 mg/day								
Morin-Papunen 2003b										
Rautio 2005										
Moro 2013	Mod	C: 30 µg EE + 3 mg DRSP M: Metformin 1500 mg/day C + M: 30 µg EE + 3 mg DRSP + Metformin 1500 mg/day	Italy 6 months	C: 25 M: 25 C + M: 26	C: 26 ± 3 M: 25 ± 5 C + M: 25 ± 4	Median (range) C: 23.7 (20.8-28.6) M: 25.1 (21.9-28.3) C + M: 26.5 (21.3-30)	Rott	NR	C: 36% M: 40% C + M: 38%	Adults BMI < 2.5
Ozgartas 2008	High	C: 35 µg EE + 2 mg CPA M: Metformin 1700 mg/day	Turkey 3 months	C = 21 M = 20	NR (≥18 years)	C: 21.72 ± 1.24 M: 21.81 ± 1.27	Rott	NR	All nonsmokers	Adults BMI < 2.5
Panidis 2011	High	C1: 35 µg EE + 2 mg CPA C2: 30 µg EE + 3 mg DRSP M: Metformin 1700 mg/day	Greece 6 months	C1 = 15 C2 = 15 M = 15	C1: 20.67 ± 4.13 C2: 22.00 ± 2.07 M: 20.53 ± 3.09	C1: 21.04 + 1.97 C2: 21.69 + 2.33M: 21.83 + 1.73	NIH	NR	NR	Adults BMI < 2.5 EE/DRSP used in meta-analysis
Sahu 2019	Mod	C: 35 µg EE + 2 mg CPA 21/7 M: Metformin 1000 mg/day	India 6 months	C = 44 M = 42	C: 26.8 ± 4.2 M: 27.0 ± 5.2	C: 25.6 ± 2.7 M: 25.7 ± 2.6	Rott	NR	All nonsmokers	Adults BMI not classified
Song 2017	High	C: 35 µg EE + 2 mg CPA M: 35 µg EE + 2 mg CPA + Metformin 1500 mg/day	China 3 months	C = 60 M = 60	C: 27.68 ± 4.99 M: 28.63 ± 5.12	C: 28.64 ± 4.89 M: 27.00 ± 3.47	Rott	NR	C: 22% M: 25%	Adults BMI not classified
Ruan 2018										

(continued)

Table 2. Continued

Study ID	RoB	Comparisons	Country, duration	N	Mean age (years)	Mean BMI	PCOS	Menarche age	Smokers	Comments
Wei 2012	Mod	C: 35 µg EE + 2 mg CPA + placebo C + M: 35 µg EE + 2 mg CPA + Metformin 1500 mg/day	China 3 months	C = 28 C + M = 30	“All female were reproductive-aged”	C: 24.9 ± 1.7 C + M: 24.7 ± 1.9	Rott	NR	All nonsmokers	Adults BMI not classified
Wu 2008 2015	Mod	C: 35µg EE + 2 mg CPA M: Metformin 1500 mg/day C + M: 35µg EE + 2 mg CPA + Metformin 1500 mg/day	China 3 months	Obese: C: 7 M: 7 C + M: 6 Nonobese: C: 12 M: 11 C + M: 10	Obese: C: 25.0 ± 4.3 M: 25.6 ± 3.6 C + M: 24.5 ± 2.4 Nonobese: C: 26.1 ± 4.6 M: 25.6 ± 4.2 C + M: 25.8 ± 4.0	Obese: C: 25.3 ± 0.8 M: 25.6 ± 0.6 CM: 25.2 ± 1.0 Nonobese: C: 21.4 ± 1.6 M: 21.5 ± 1.8 CM: 21.6 ± 1.4	Rott	NR	Excluded	Adults BMI > 25 BMI < 25

Abbreviations: AES, Androgen Excess Society; BMI, body mass index; C, Oral contraceptive pill; CPA, cyproterone acetate; DRSP, drospirenone; DSG, desogestrel; EE, ethinylestradiol; LVG, levonorgestrel; M, metformin; NIH, National Institutes of Health; NOR, norgestimate; NR, Not reported; PCOS, polycystic ovary syndrome; RoB, risk of bias; Rott, Rotterdam; SPL, spironolactone.

**COCP vs combination**

No differences were noted in hirsutism between COCP alone and combination treatment. COCP alone was inferior to combination treatment for FAI (MD 0.58; 95% CI, 0.36 to 0.80), SHBG (MD -16.61 nmol/L; 95% CI, -28.51 to -4.71), dehydroepiandrosterone sulfate (DHEAS) (MD 0.93 µmol/L; 95% CI, 0.54 to 1.31), and androstenedione (MD 1.37 nmol/L; 95% CI, 0.14 to 2.60), whereas no differences were noted regarding total and free testosterone (Table 4; Supplementary Fig. S5 (23)).

**Metformin vs combination**

Metformin was inferior for DHEAS (MD 82.38 µmol/L; 95% CI, 15.43 to 149.36) and total testosterone (MD 0.64 nmol/L; 95% CI, 0.26 to 1.02). None of the identified studies reported hirsutism as an outcome for this comparison (Table 5; Supplementary Fig. S6 (23)).

**Metabolic Outcomes**

**Metformin vs COCP**

Metformin was more effective in lowering overall fasting insulin levels (MD -27.12 pmol/L; 95% CI, -40.65 to -13.59), total cholesterol (MD -0.40 mmol/L; 95% CI, -0.66 to -0.14), triglycerides (MD -0.15 mmol/L; 95% CI, -0.29 to -0.01), and C-reactive protein (CRP) levels (MD -11.31 nmol/L; 95% CI, -19.78 to -2.85) compared with COCP (Table 3). For fasting glucose, homeostatic model assessment for insulin resistance (HOMA-IR), high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol and plasminogen activator inhibitor 1 (PAI-1), there were no differences.

In subgroup analyses on women with a BMI ≥ 25 kg/m<sup>2</sup> (Supplementary Fig. S2; Supplementary Table S1) (23), the metformin-treated group had lower CRP (MD -33.09 nmol/L; 95% CI, -47.33 to -18.84) and HDL (MD -0.24 mmol/L; 95% CI, -0.38 to -0.09) compared with COCP. Fasting insulin was lower with metformin regardless of BMI (Supplementary Table S1; Supplementary Fig. S2) (23).

Metformin was superior to COCP for fasting insulin, both in adults (MD -22.17 pmol/L; 95% CI, -29.93 to -14.42) and adolescents (MD -30.03 pmol/L; 95% CI, -78.63 to -18.56). Metformin was also superior for total cholesterol, both in adults (MD -0.34 mmol/L; 95% CI, -0.60 to -0.08) and for adolescents (MD -1.12 mmol/L; 95% CI, -1.74 to -0.50) (Supplementary Table S1) (23). The type of COCP showed no major differences compared with the overall result (Supplementary Table S2) (23).

**COCP vs combination**

COCP alone increased fasting insulin (MD 17.03 pmol/L; 95% CI, 7.79 to 26.26), HOMA-IR (MD 0.44; 95% CI, 0.17 to 0.70), and CRP (MD 1.94 nmol/L; 95% CI, 0.05 to 3.84) compared with combination treatment (Table 4; Supplementary Fig. S5 (23)). In women with a BMI < 25 kg/m<sup>2</sup>, COCP alone resulted in marginally higher glucose (MD 0.25 mmol/L; 95% CI, 0.07 to 0.43) compared with combination treatment.

**Metformin vs combination**

Metformin alone showed marginally lower fasting glucose (MD -0.33 mmol/L; 95% CI, -0.64 to -0.01) and CRP (MD

Study	Risk of bias domains					Overall
	D1	D2	D3	D4	D5	
Aghamohammadzadeh 2010	⊗	⊗	⊕	⊕	⊖	⊗
Allen 2005	⊕	⊖	⊕	⊕	⊖	⊖
Al-Zubeidi 2015	⊕	⊖	⊗	⊖	⊗	⊗
Bilgir 2009	⊗	⊖	⊕	⊖	⊖	⊗
Bodur 2018	⊕	⊖	⊗	⊕	⊖	⊗
Cetinkalp 2009	⊗	⊗	⊖	⊖	⊖	⊗
Christakou 2014	⊖	⊖	⊕	⊕	⊖	⊖
Cibula 2005	⊕	⊖	⊖	⊖	⊖	⊖
Dardzinska 2014	⊖	⊖	⊖	⊕	⊕	⊖
Dorgham 2021	⊖	⊖	⊗	⊖	⊖	⊗
El Maghraby 2015	⊗	⊖	⊖	⊗	⊖	⊗
Elter 2002	⊕	⊖	⊕	⊕	⊖	⊖
Essah 2011	⊕	⊕	⊕	⊕	⊕	⊕
Feng 2016	⊕	⊕	⊖	⊖	⊖	⊖
Fonseka 2020	⊕	⊕	⊕	⊕	⊕	⊕
Glintborg 2014a	⊗	⊖	⊗	⊖	⊕	⊗
Harborne 2003	⊕	⊖	⊗	⊕	⊖	⊗
Hoeger 2008	⊖	⊖	⊗	⊖	⊖	⊗
Kaya 2012	⊗	⊗	⊕	⊖	⊖	⊗
Kaya 2015	⊗	⊖	⊕	⊖	⊖	⊗
Kebapcilar 2009	⊗	⊖	⊕	⊖	⊖	⊗
Kebapcilar 2010	⊗	⊖	⊕	⊖	⊖	⊗
Kilic 2011	⊕	⊖	⊕	⊕	⊖	⊖
Kumar 2018	⊖	⊖	⊕	⊕	⊖	⊖
Luque-Ramirez 2009	⊕	⊖	⊖	⊕	⊕	⊖
Lv 2005	⊗	⊖	⊗	⊖	⊕	⊗
Meyer 2007	⊖	⊖	⊕	⊕	⊕	⊖
Mhao 2016	⊗	⊗	⊗	⊗	⊗	⊗
Morin-Papunen 2000 (obese)	⊗	⊖	⊖	⊕	⊖	⊗
Morin-Papunen 2003a (non-obese)	⊗	⊖	⊕	⊕	⊖	⊗
Moro 2013	⊕	⊖	⊕	⊖	⊖	⊖
Ozgurtas 2008	⊗	⊗	⊕	⊖	⊖	⊗
Panidis 2011	⊗	⊗	⊕	⊕	⊖	⊗
Sahu 2019	⊖	⊖	⊕	⊖	⊕	⊖
Song 2017	⊗	⊖	⊕	⊖	⊖	⊗
Wei 2012	⊕	⊖	⊕	⊖	⊖	⊖
Wu 2009	⊖	⊖	⊖	⊕	⊖	⊖

Domains:  
D1: Bias arising from the randomization process.  
D2: Bias due to deviations from intended intervention.  
D3: Bias due to missing outcome data.  
D4: Bias in measurement of the outcome.  
D5: Bias in selection of the reported result.

Judgement  
⊗ High  
⊖ Some concerns  
⊕ Low

Figure 2. Risk of bias assessments of the included studies.

**Table 3. Grade assessments and evidence profile on outcomes comparing metformin with combined oral contraceptive pills**

No. studies	Risk of bias	Inconsistency	Indirectness	Imprecision	Met (n)	COCP (n)	Effect, random, MD (95% CI)	Favors	Certainty
Weight (kg)									
Overall 7	Serious <sup>d</sup>	Very serious <sup>e</sup>	No serious	Serious <sup>b</sup>	188	177	-1.25 kg (-12.95 to 10.44)	No difference	⊕○○○ VERY LOW
Waist-hip ratio (WHR)									
Overall 7	Very serious <sup>b</sup>	Very serious <sup>e</sup>	Serious <sup>f</sup>	No serious	88	85	-0.02 (-0.03 to 0.00)	No difference	⊕○○○ VERY LOW
BMI (kg/m <sup>2</sup> )									
Overall 20	Serious <sup>c</sup>	Very serious <sup>e</sup>	No serious	No serious	401	390	-0.74 kg/m <sup>2</sup> (-1.58 to 0.09)	No difference	⊕○○○ VERY LOW
Hirsutism (FG score)									
Overall 6	Very serious <sup>b</sup>	Very serious <sup>e</sup>	No serious	Serious <sup>b</sup>	120	127	0.72 (-1.40 to 2.85)	No difference	⊕○○○ VERY LOW
Free androgen index (FAI)									
Overall 8	Very serious <sup>b</sup>	Very serious <sup>e</sup>	No serious	No serious	107	114	7.08 (4.81 to 9.36)	COCP	⊕⊕○○ LOW <sup>†</sup>
Sex hormone binding globulin (SHBG) (nmol/L)									
Overall 9	Very serious <sup>b</sup>	Very serious <sup>e</sup>	No serious	No serious	171	168	-118.61 (-174.46 to -62.75)	COCP	⊕⊕○○ LOW <sup>†</sup>
Dehydroepiandrosterone sulfate (DHEAS) (umol/L)									
Overall 7	Very serious <sup>b</sup>	Serious <sup>d</sup>	Serious <sup>f</sup>	No serious	170	160	1.18 (0.17 to 2.18)	COCP	⊕○○○ VERY LOW
Androstendione (nmol/L)									
Overall 4	Very serious <sup>b</sup>	Serious <sup>d</sup>	No serious	No serious	66	70	3.25 (1.46 to 5.04)	COCP	⊕○○○ VERY LOW
Total testosterone (nmol/L)									
Overall 17	Serious <sup>c</sup>	Serious <sup>d</sup>	No serious	No serious	323	321	0.48 (0.32 to 0.64)	COCP	⊕⊕○○ LOW
Free testosterone (pmol/L)									
Overall 4	Very serious <sup>b</sup>	Serious <sup>d</sup>	No serious	Serious <sup>b</sup>	107	91	3.59 (1.41 to 5.78)	COCP	⊕○○○ VERY LOW
Fasting insulin (pmol/L)									
Overall 14	Serious <sup>c</sup>	Serious <sup>d</sup>	No serious	No serious	292	282	-27.12 (-40.65 to -13.59)	Metformin	⊕⊕○○ LOW
Fasting glucose (mmol/L)									
Overall 10	Very serious <sup>b</sup>	Serious <sup>d</sup>	No serious	No serious	210	201	-0.08 (-0.23 to 0.06)	No difference	⊕○○○ VERY LOW
Homeostatic model assessment for insulin resistance (HOMA-IR)									
Overall 8	Serious <sup>d</sup>	Very serious <sup>e</sup>	Serious <sup>f</sup>	No serious	202	184	-0.60 (-1.24 to 0.05)	No difference	⊕○○○ VERY LOW

(continued)

**Table 3. Continued**

	No. studies	Risk of bias	Inconsistency	Indirectness	Imprecision	Met (n)	COCP (n)	Effect, random, MD (95% CI)	Favors	Certainty
Total cholesterol (mmol/L)										
Overall	13	Serious <sup>c</sup>	Very serious <sup>e</sup>	No serious	No serious	281	273	-0.40 (-0.66 to -0.14)	Metformin	⊕○○○ VERY LOW
High-density lipoprotein (HDL) cholesterol (mmol/L)										
Overall	13	Serious <sup>d</sup>	Very serious <sup>e</sup>	No serious	No serious	283	267	-0.05 (-0.18 to 0.07)	No difference	⊕○○○ VERY LOW
Low-density lipoprotein (LDL) cholesterol (mmol/L)										
Overall	13	Serious <sup>d</sup>	Serious <sup>d</sup>	No serious	No serious	281	273	-0.08 (-0.29 to 0.13)	No difference	⊕○○○ LOW
Triglycerides (mmol/L)										
Overall	13	Serious <sup>e</sup>	Serious <sup>d</sup>	No serious	No serious	277	264	-0.15 (-0.29 to -0.01)	Metformin	⊕○○○ LOW
C-reactive protein (CRP) (mmol/L)										
Overall	7	Serious <sup>e</sup>	Serious <sup>d</sup>	No serious	No serious	165	156	-11.31 (-19.78 to -2.85)	Metformin	⊕○○○ LOW
Plasminogen activator inhibitor-1 levels (PAI-1) (ng/mL)										
Overall	2	Very serious <sup>b</sup>	Serious <sup>d</sup>	Serious <sup>f</sup>	Very serious <sup>g</sup>	23	27	-1.05 (-24.65 to 22.54)	No difference	⊕○○○ VERY LOW
Girls with restored menses										
Overall	5	Very serious <sup>b</sup>	No serious	No serious	No serious	85	82	0.17 (0.05 to 0.57)	COCP	⊕○○○ LOW

Bolding indicates a statistically significant MD ( $P < .05$ ).  
 Abbreviations: COCP, combined oral contraceptive pills; MD, mean difference; RCT; randomized controlled trials; RoB; risk of bias.  
<sup>a</sup>Downgraded once as all studies high or moderate RoB.  
<sup>b</sup>Downgraded twice as all studies or all but one are high RoB.  
<sup>c</sup>Downgraded once as studies high to moderate RoB.  
<sup>d</sup>Downgraded once as  $I^2$  is close to or > 50% but CI partly overlapping.  
<sup>e</sup>Downgraded twice as  $I^2$  very high and CI not overlapping.  
<sup>f</sup>Downgraded once as no adolescents in the overall group.  
<sup>g</sup>Downgraded twice as there are very few studies.  
<sup>h</sup>Downgraded once as the CI is wide.  
<sup>i</sup>Upgraded once due to large effect.

**Table 4. Grade assessments and evidence profile of outcomes in PCOS comparing COCP with metformin and COCP**

No studies	Quality assessment					No. participants		Effect, random [95% CI]	Favors	Certainty
	Risk of bias	Inconsistency	Indirectness	Imprecision	Other	COCP	COCP + Met			
Weight (kg)										
Overall	6 Serious <sup>a</sup>	No serious	No serious	No serious	None	125	124	MD 0.06 (-1.99 to 2.11)	No difference	⊕⊕⊕ MODERATE
Waist-hip ratio (WHR)										
Overall	7 Serious <sup>a</sup>	No serious	No serious	No serious	None	143	141	MD 0.00 (-0.02 to 0.01)	No difference	⊕⊕⊕ MODERATE
BMI (kg/m <sup>2</sup> )										
Overall	14 Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	284	279	MD -0.22 (-0.98 to 0.55)	No difference	⊕⊕⊕ LOW
Hirsutism (FG score)										
Overall	3 Serious <sup>a</sup>	No serious	No serious	No serious	None	71	86	MD 0.99 (-0.35 to 2.32)	No difference	⊕⊕⊕ MODERATE
Free androgen index (FAI)										
Overall	6 Serious <sup>a</sup>	No serious	No serious	No serious	None	172	171	MD 0.58 (0.36 to 0.80)	COCP + met	⊕⊕⊕ MODERATE
Sex hormone binding globulin (SHBG) (nmol/L)										
Overall	7 Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	177	175	MD -16.61 (-28.51 to -4.71)	COCP + met	⊕⊕⊕ LOW
Dehydroepiandrosterone sulfate (DHEAS) (umol/L)										
Overall	9 Very serious <sup>c</sup>	No serious	No serious	No serious	None	231	231	MD 0.93 (0.54 to 1.31)	COCP + met	⊕⊕⊕ LOW
Total testosterone (nmol/L)										
Overall	12 Serious <sup>a</sup>	No serious	No serious	No serious	None	290	287	MD 0.08 (-0.00 to 0.15)	No difference	⊕⊕⊕ MODERATE
Free testosterone (pmol/L)										
Overall	6 Very serious <sup>c</sup>	No serious	No serious	No serious	None	103	100	MD 0.96 (-0.35 to 2.27)	No difference	⊕⊕⊕ LOW
Androstendione (nmol/L)										
Overall	6 Very serious <sup>c</sup>	Very serious <sup>d</sup>	No serious	No serious	None	159	157	MD 1.37 (0.14 to 2.60)	COCP + met	⊕⊕⊕ VERY LOW
Fasting insulin (pmol/L)										
Overall	14 Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	319	314	MD 17.03 (7.79 to 26.26)	COCP + met	⊕⊕⊕ LOW
Fasting glucose (mmol/L)										
Overall	9 Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	244	239	MD 0.15 (-0.01 to 0.30)	No difference	⊕⊕⊕ LOW
Homeostatic model assessment for insulin resistance (HOMA)										

(continued)

**Table 4. Continued**

No studies	Quality assessment				No. participants		Effect, random [95% CI]	Favors	Certainty
	Risk of bias	Inconsistency	Indirectness	Imprecision	Other	COCP			
Overall 10	Very serious <sup>c</sup>	No serious	No serious	No serious	None	272	268	COCP + met	⊕⊕○○ LOW
Total cholesterol (mmol/L)									
Overall 14	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	350	349	No difference	⊕⊕○○ LOW
Low-density lipoprotein (LDL) cholesterol (mmol/L)									
Overall 14	Serious <sup>a</sup>	No serious	No serious	No serious	None	350	348	No difference	⊕⊕○○ MODERATE
High-density lipoprotein (HDL) cholesterol (mmol/L)									
Overall 14	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	350	349	No difference	⊕⊕○○ LOW
Triglycerides (mmol/L)									
Overall 14	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	None	350	348	No difference	⊕⊕○○ LOW
C-reactive protein (CRP) (nmol/L)									
Overall 4	Serious <sup>a</sup>	No serious	No serious	No serious	None	74	68	COCP + met	⊕○○○ VERY LOW
Plasminogen activator inhibitor-1 levels (PAI-1) (ng/mL)									
Overall 2	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	Very serious <sup>e</sup>	None	27	21	No difference	⊕○○○ VERY LOW
Health-related quality of life (HR-QoL)									
Overall 1	Serious <sup>a</sup>	Not applicable	Not applicable	Very serious <sup>e</sup>	None	23	23	Narrative review	⊕○○○ VERY LOW

Bolding indicates a statistically significant MD ( $P < .05$ ).  
 Abbreviations: COCP, combined oral contraceptive pills; MD, mean difference; Met, metformin; PCOS, polycystic ovary syndrome; RCT, randomized controlled trials; RoB, risk of bias.  
<sup>a</sup>Downgraded once as the majority of evidence is at moderate or high RoB.  
<sup>b</sup>Downgraded once due to I2 > 50%.  
<sup>c</sup>Downgraded twice due to high RoB.  
<sup>d</sup>Downgraded twice due to I2 > 50% and CI not overlapping.  
<sup>e</sup>Downgraded twice due to very few patients.

**Table 5. Grade assessments and evidence profile of outcomes in PCOS comparing metformin with metformin and combined oral contraceptive pills**

No. studies	Risk of bias	Inconsistency	Indirectness	Imprecision	Met	Met + COCP	Effect, random, MD (95% CI)	Favors	Certainty	
Waist-hip ratio (WHR)										
Overall	2	Serious <sup>a</sup>	No serious	No serious	Very serious <sup>e</sup>	18	16	<b>−0.03</b> (−0.06 to −0.01)	<b>Metformin</b>	⊕○○○ VERY LOW
BMI (kg/m <sup>2</sup> )										
Overall	4	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	Serious <sup>d</sup>	60	57	−1.31 (−2.65 to 0.03)	No difference	⊕○○○ VERY LOW
Dehydroepiandrosterone sulfate (DHEAS) (umol/L)										
Overall	2	Serious <sup>a</sup>	No serious	No serious	Serious <sup>d</sup>	60	58	<b>82.39</b> (15.43 to 149.36)	<b>Metformin + COCP</b>	⊕⊕○○ LOW
Total testosterone (nmol/L)										
Overall	4	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	73	71	<b>0.64</b> (0.26 to 1.02)	<b>Metformin + COCP</b>	⊕⊕○○ LOW
Fasting insulin pmol/L)										
Overall	4	Serious <sup>a</sup>	No serious	No serious	Serious <sup>d</sup>	60	57	−3.35 pmol/L (−17.67 to 10.97)	No difference	⊕⊕○○ LOW
Fasting glucose (mmol/L)										
Overall	2	Serious <sup>a</sup>	Serious <sup>b</sup>	Serious <sup>e</sup>	Very serious <sup>e</sup>	47	41	<b>−0.33 mmol/L</b> (−0.64 to −0.01)	<b>Metformin</b>	⊕○○○ VERY LOW
Homeostatic model assessment for insulin resistance (HOMA-IR)										
Overall	3	Serious <sup>a</sup>	Very serious <sup>c</sup>	No serious	Serious <sup>d</sup>	59	53	−1.28 (−2.92 to 0.35)	No difference	⊕○○○ VERY LOW
Total cholesterol (mmol/L)										
Overall	3	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	67	67	−0.61 mmol/L (−1.23 to 0.01)	No difference	⊕⊕○○ LOW
High-density lipoprotein (HDL) cholesterol (mmol/L)										
Overall	3	Serious <sup>a</sup>	No serious	No serious	No serious	67	67	−0.06 mmol/L (−0.17 to 0.04)	No difference	⊕⊕⊕○ MODERATE
Low-density lipoprotein (LDL) cholesterol (mmol/L)										
Overall	3	Serious <sup>a</sup>	Serious <sup>b</sup>	No serious	No serious	67	67	−0.14 mmol/L (−0.46 to 0.17)	No difference	⊕⊕○○ LOW
Triglycerides (mmol/L)										
Overall	3	Serious <sup>a</sup>	Very serious <sup>c</sup>	No serious	No serious	67	67	−0.20 mmol/L (−0.72 to 0.31)	No difference	⊕○○○ VERY LOW
C-reactive protein (CRP) (nmol/L)										
Overall	2	Serious <sup>a</sup>	No serious	No serious	Serious <sup>d</sup>	47	41	<b>−4.08 nmol/L</b> (−6.01 to −2.16)	<b>Metformin</b>	⊕⊕○○ LOW

Bolding indicates a statistically significant MD ( $P < .05$ ).

Abbreviations: COCP, combined oral contraceptive pills; MD, mean difference; Met, metformin; PCOS, polycystic ovary syndrome; RCT; randomized controlled trials; RoB; risk of bias.

<sup>a</sup>Downgraded once as the studies are at moderate or high RoB.

<sup>b</sup>Downgraded once as  $I^2$  is close to or  $>50\%$  but CI partly overlapping.

<sup>c</sup>Downgraded twice as  $I^2$  very high and CI not overlapping.

<sup>d</sup>Downgraded once as there are only a few studies.

<sup>e</sup>Downgraded twice as there are very few participants.

−4.08 nmol/L; 95% CI, −6.01 to −2.16), compared with a combination treatment (Table 5; Supplementary Fig. S6 (23)).

### Cycle Regularity and Quality of Life Outcomes

Metformin was inferior to COCP on restoring regular menses overall (OR 0.17; 95% CI, 0.05 to 0.57), as well as in adults (OR 0.19; 95% CI, 0.05 to 0.72) and adolescents (OR 0.10; 95% CI, 0.01 to 1.92).

We were not able to perform a meta-analysis on health-related quality of life. Results from 3 identified studies (32, 42, 50) showed conflicting results and our overall assessment is that there is no difference in quality of life.

### Side Effects

Due to lack of systematic reporting, where many studies do not report adverse effects at all or do not report in a similar manner, we were not able to perform a meta-analysis. However, the reports suggest more gastrointestinal side effects with metformin.

### Discussion

#### Main Findings

This extensive systematic review and meta-analysis, including 36 RCTs, was performed to directly inform recommendations on the use of metformin and COCP in women with PCOS, as

part of the 2023 update of the International Evidence-based Guidelines on the Assessment and Treatment of PCOS. Our findings showed that metformin was superior to COCP for metabolic outcomes, especially in women with PCOS and a BMI  $\geq 25$  kg/m<sup>2</sup>, whereas COCP was superior for improving cycle regularity and, in women with a BMI  $< 25$  kg/m<sup>2</sup>, for improving hirsutism. The combination of metformin and COCP was more effective for improving biochemical hyperandrogenism, insulin levels, and insulin resistance than COCP monotherapy.

## Interpretation

### Clinical and biochemical hyperandrogenism

For many patients with PCOS, medical treatment is indicated to treat clinical hyperandrogenism, including acne and hirsutism. For these symptoms, COCPs have been recommended as first-line treatment. In addition, COCPs regulate menstrual cycles and provide contraception (12, 15). In this study we confirm that COCP was superior to metformin for treatment of hirsutism in women with BMI  $< 25$  kg/m<sup>2</sup>, whereas evidence for other BMI groups was of very low quality. However, we found that COCP was superior in improving biochemical hyperandrogenism compared with metformin. A systematic review on COCP treatment in women with PCOS, including both RCTs and non-RCTs, suggested that COCP containing CPA might be more effective in improving hirsutism. However, no direct comparisons between different COCP were made, limiting the conclusions (15). General population studies have shown that CPA increases the risk of venous thromboembolism compared with other COCP (71, 72); hence, COCPs containing CPA are currently not recommended as first-line treatments for PCOS (20).

No differences were found regarding hirsutism between COCP and combination treatment. Nevertheless, combination treatment was more effective in improving FAI, SHBG, and DHEAS compared with COCP alone. One systematic review (17) studied hirsutism and acne in PCOS, comparing monotherapy with metformin or COCP with combination treatment, with monotherapy being less effective for hirsutism compared with combination treatment. However, that review also included studies with shorter treatment durations (3 months) hindering interpretation of their findings.

As high insulin levels increase luteinizing hormone-mediated ovarian androgen synthesis (73), combination treatment presumably targets several distinct mechanisms leading to improved clinical and biochemical hyperandrogenism. Thus, combination treatment theoretically offers several benefits. Importantly, additional high-quality prospective studies are needed to better ascertain the efficacy of these treatment regimens. We also recognize the importance of self-assessment of severity and impact on quality of life, over and above clinical assessment.

### Anthropometric outcomes

Women with PCOS have an increased prevalence of the metabolic syndrome (74, 75). In our meta-analysis, there were no differences in weight, WHR, and BMI, but evidence was only available with very low certainty. Previously, metformin has been shown to improve BMI compared with placebo (76). A recent systematic review which included nonrandomized trials found that combined metformin, COCP, and anti-

androgen treatment improved BMI and glucose tolerance (11). However, this study did not compare metformin monotherapy to a combination treatment with metformin and COCP. In our study the combination of COCP and metformin showed no benefit for anthropometric measures compared with COCP alone.

### Metabolic outcomes

Hyperinsulinemia and insulin resistance play important roles in the pathophysiology of PCOS for both normal-weight women and women with obesity (2, 3, 73, 77). Metformin decreases insulin resistance and insulin levels. COCP does not have any major effects on carbohydrate metabolism in healthy women, whereas the effect on lipid metabolism depends on the level of estrogen and type of progestin, with potential negative effects (78, 79, 80, 81). Our systematic review confirms that metformin is superior in lowering fasting insulin levels, total cholesterol, and triglycerides compared with COCP, in line with findings reported previously (12).

Metformin can be used where COCPs are contradicted, in older women, women with obesity or medical conditions (such as migraine with aura, risk of venous thromboembolism or severe hypertension) or where pregnancy is desired. The major disadvantage of metformin treatment is the generally mild, usually self-limited, gastrointestinal side effects, limiting patient acceptance of appropriate doses. We also report that metformin lowered CRP compared with COCP treatment. CRP is often used as a metabolic risk marker and is associated with an increased risk of cardiovascular disease (82). Metformin has previously been shown to decrease CRP, both in obese and nonobese women, compared with placebo (83). COCP appears to increase CRP (79, 84, 85), yet the significance of these effects remains unknown.

Importantly, the addition of metformin to COCP improved insulin levels and insulin resistance, as compared with COCP alone. This is of special interest for women with PCOS with additional risk factors for type 2 diabetes, such as a high BMI (86, 87). Two systematic reviews found that adding metformin to COCP and anti-androgens improved BMI and glucose tolerance (11, 88).

### Strengths and Limitations

This review presents the most up-to-date evidence on COCP and metformin treatment in women with PCOS. Strengths of the report include the rigorous processes, including PICO developed by clinicians, researchers, and patients; and the large number of RCTs included. Because critical appraisal is inherently subjective, both RoB and GRADE assessments were conducted by 2 authors independently. Several subgroup analyses were included to highlight benefits in population subgroups of interest. Limitations include those inherent in the included studies, with many studies having a high RoB, mainly due to often poorly described randomization processes or lack of blinding. Certainty of evidence was also affected by inconsistency of effect sizes and small sample sizes. Several studies did not provide data on BMI.

Identified research gaps include the lack of larger studies on adolescents with PCOS and on the comparison of metformin with combination treatment. Additional high-quality studies are needed to assess symptoms, especially related to clinical hyperandrogenism rather than biochemical markers of hyperandrogenism. Future directions include understanding

mechanisms of action and factors that may impact on medication responses, such as genotype and phenotype of PCOS.

## Conclusion

Results from this extensive systematic review and meta-analysis will advise the pending 2023 PCOS guideline update. The guideline will recommend COCP to be used over metformin for management of irregular cycles and hirsutism, and metformin over COCP for metabolic indications in PCOS. While our meta-analyses indicated that the combined treatment with metformin and COCP improved biochemical hyperandrogenism, insulin levels, and insulin resistance more than COCP alone, no difference was seen in clinical outcomes.

For women with PCOS, the choice of treatment should be based on clinical symptoms. Combination treatment appears to be beneficial in high metabolic risk groups, targeting the 2 major endocrine disturbances seen: hyperinsulinemia and hyperandrogenism. Our results, including several sub-analyses, add to the current evidence base and contribute toward reaching the ultimate goal of shared decision-making and effective, tailored, and individualized treatment for patients with PCOS.

## Funding

J.M. received funding from The Medical Society of Finland and Orion Research foundation. M.F. received funding from the Iris foundation and Hjalmar Svensson Foundation. A.M. is supported by a biomedical research fellowship from the National Health and Medical Research Council (NHMRC) of Australia.

## Author Contributions

This systematic review will inform the forthcoming 2023 update of the International Evidence-based Guideline on Assessment and Management of PCOS, led by H.T. Senior experts within the guideline working group included T.P., D.R., P.M.S., C.T.T., A.P., S.W., A.M. and H.T. C.C.T. and A.M. were part of the evidence team. The search, screening, and RoB and GRADE assessments were performed by J.M., M.F., and S.A. J.M. and M.F. did the data extraction. J.M. and M.F. drafted the first version of the manuscript, which was revised by the other authors. All authors approved of the final version of the manuscript.

## Disclosures

J.M. reports a postdoc research grant from Orion Research Foundation. D.R. reports an honorarium from Novo Nordisk for a lecture on PCOS and obesity. Neither Orion nor Novo Nordisk had any influence on this work. The other authors have no conflicts to declare.

## Data Availability

The data underlying this article are secondary, aggregated from already published work.

## References

- Bozdag G, Mumusoglu S, Zengin D, Karabulut E, Yildiz BO. The prevalence and phenotypic features of polycystic ovary syndrome: a systematic review and meta-analysis. *Hum Reprod.* 2016;31(12):2841-2855.
- Dunaif A, Segal KR, Futterweit W, Dobrjansky A. Profound peripheral insulin resistance, independent of obesity, in polycystic ovary syndrome. *Diabetes.* 1989;38(9):1165-1174.
- Cassar S, Misso ML, Hopkins WG, Shaw CS, Teede HJ, Stepto NK. Insulin resistance in polycystic ovary syndrome: a systematic review and meta-analysis of euglycaemic-hyperinsulinaemic clamp studies. *Hum Reprod.* 2016;31(11):2619-2631.
- Kakoly NS, Khomami MB, Joham AE, *et al.* Ethnicity, obesity and the prevalence of impaired glucose tolerance and type 2 diabetes in PCOS: a systematic review and meta-regression. *Hum Reprod Update.* 2018;24(4):455-467.
- Wu J, Zhu Y, Jiang Y, Cao Y. Effects of metformin and ethinyl estradiol-cyproterone acetate on clinical, endocrine and metabolic factors in women with polycystic ovary syndrome. *Gynecol Endocrinol.* 2008;24(7):392-398.
- Diamanti-Kandarakis E, Dunaif A. Insulin resistance and the polycystic ovary syndrome revisited: an update on mechanisms and implications. *Endocr Rev.* 2012;33(6):981-1030.
- Rena GP ER, Sakamoto K. Molecular mechanism of action of metformin: old or new insights? *Diabetologia.* 2013;56(9):1898-1906.
- Malin SK, Kashyap SR. Effects of metformin on weight loss: potential mechanisms. *Curr Opin Endocrinol Diabetes Obes.* 2014;21(5):323-329.
- Naderpoor N, Shorakae S, de Courten B, Misso ML, Moran LJ, Teede HJ. Metformin and lifestyle modification in polycystic ovary syndrome: systematic review and meta-analysis. *Hum Reprod Update.* 2015;21(5):560-574.
- RCOG. Metformin therapy for the management of infertility in women with polycystic ovary syndrome: scientific impact paper No. 13. *BJOG.* 2017;124(12):e306-e313.
- Luque-Ramírez M, Nattero-Chávez L, Ortiz Flores AE, Escobar-Morreale HF. Combined oral contraceptives and/or anti-androgens versus insulin sensitizers for polycystic ovary syndrome: a systematic review and meta-analysis. *Hum Reprod Update.* 2018;24(2):225-241.
- Teede H, Tassone EC, Piltonen T, *et al.* Effect of the combined oral contraceptive pill and/or metformin in the management of polycystic ovary syndrome: a systematic review with meta-analyses. *Clin Endocrinol (Oxf).* 2019;91(4):479-489.
- Forslund M, Melin J, Alesi S, *et al.* Combined oral contraceptive pill compared with no medical treatment in the management of polycystic ovary syndrome: a systematic review. *Clin Endocrinol.* 2023;99(1):79-91.
- De Leo V, Musacchio MC, Cappelli V, Piomboni P, Morgante G. Hormonal contraceptives: pharmacology tailored to women's health. *Hum Reprod Update.* 2016;22(5):634-646.
- Amiri M, Kabir A, Nahidi F, Shekofteh M, Ramezani Tehrani F. Effects of combined oral contraceptives on the clinical and biochemical parameters of hyperandrogenism in patients with polycystic ovary syndrome: a systematic review and meta-analysis. *Eur J Contracept Reprod Health Care.* 2018;23(1):64-77.
- World Health Organization. 2015 quick reference chart for the WHO medical eligibility criteria for contraceptive use. 5th ed.; 2015. Accessed April 26, 2023. <https://www.fhi360.org/sites/default/files/media/documents/chart-medical-eligibility-contraceptives-english.pdf>
- Fraison E, Kostova E, Moran LJ, *et al.* Metformin versus the combined oral contraceptive pill for hirsutism, acne, and menstrual pattern in polycystic ovary syndrome. *Cochrane Database Syst Rev.* 2020;8(8):Cd005552.
- Joham AE, Norman RJ, Stener-Victorin E, *et al.* Polycystic ovary syndrome. *Lancet Diabetes Endocrinol.* 2022;10(9):668-680.
- Al Wattar BH, Teede H, Garad R, *et al.* Harmonising research outcomes for polycystic ovary syndrome: an international multi-stakeholder core outcome set. *Hum Reprod.* 2021;36(3):523-528.
- Teede HJ, Misso ML, Costello MF, *et al.* Recommendations from the international evidence-based guideline for the assessment and

- management of polycystic ovary syndrome. *Fertil Steril*. 2018;110(3):364-379.
21. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg*. 2010;8(5):336-341.
  22. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
  23. Melin J, Forslund M, Alesi S, et al. Supplementary Material. Metformin and combined oral contraceptive pills in the management of polycystic ovary syndrome: a systematic review and meta-analysis. Monash University. Dataset. 2023. <https://doi.org/10.26180/23632254.v1>
  24. Sterne JAC, Savović J, Page MJ, et al. Rob 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:l4898.
  25. Schünemann HBJ, Guyatt G, Oxman A, eds. *GRADE Handbook for Grading Quality of Evidence and Strength of Recommendations*. The GRADE Working Group; 2022.
  26. Christakou C, Kollias A, Piperi C, Katsikis I, Panidis D, Diamanti-Kandarakis E. The benefit-to-risk ratio of common treatments in PCOS: effect of oral contraceptives versus metformin on atherogenic markers. *Hormones*. 2014;13(4):488-497.
  27. Cibula D, Fanta M, Vrbikova J, et al. The effect of combination therapy with metformin and combined oral contraceptives (COC) versus COC alone on insulin sensitivity, hyperandrogenism, SHBG and lipids in PCOS patients. *Hum Reprod*. 2005;20(1):180-184.
  28. Dardzinska J, Rachon D, Kuligowska-Jakubowska M, et al. Effects of metformin or an oral contraceptive containing cyproterone acetate on serum C-reactive protein, interleukin-6 and soluble vascular cell adhesion molecule-1 concentrations in women with polycystic ovary syndrome. *Exp Clin Endocrinol Diabetes*. 2014;122(2):118-125.
  29. Glintborg D, Altinok M, Mumm H, Hermann A, Ravn P, Andersen M. Body composition is improved during 12 months' treatment with metformin alone or combined with oral contraceptives compared with treatment with oral contraceptives in polycystic ovary syndrome. *J Clin Endocrinol Metab*. 2014;99(7):2584-2591.
  30. Glintborg D, Mumm H, Altinok M, Richelsen B, Bruun J, Andersen M. Adiponectin, interleukin-6, monocyte chemoattractant protein-1, and regional fat mass during 12-month randomized treatment with metformin and/or oral contraceptives in polycystic ovary syndrome. *J Endocrinol Invest*. 2014;37(8):757-764.
  31. Glintborg D, Mumm H, Holst J, Andersen M. Effect of oral contraceptives and/or metformin on GLP-1 secretion and reactive hypoglycaemia in polycystic ovary syndrome. *Endocr Connect*. 2017;6(4):267-277.
  32. Altinok M, Ravn P, Andersen M, Glintborg D. Effect of 12-month treatment with metformin and/or oral contraceptives on health-related quality of life in polycystic ovary syndrome. *Gynecol Endocrinol*. 2018;34(10):859-863.
  33. Harborne L, Fleming R, Lyall H, Sattar N, Norman J. Metformin or antiandrogen in the treatment of hirsutism in polycystic ovary syndrome. *J Clin Endocrinol Metab*. 2003;88(9):4116-4123.
  34. Luque-Ramírez M, Mendieta-Azcona C, Alvarez-Blasco F, Escobar-Morreale H. Effects of metformin versus ethinyl-estradiol plus cyproterone acetate on ambulatory blood pressure monitoring and carotid intima media thickness in women with the polycystic ovary syndrome. *Fertil Steril*. 2009;91(6):2527-2536.
  35. Morin-Papunen L, Vauhkonen I, Koivunen R, Ruokonen A, Martikainen H, Tapanainen J. Endocrine and metabolic effects of metformin versus ethinyl estradiol-cyproterone acetate in obese women with polycystic ovary syndrome: a randomized study. *J Clin Endocrinol Metab*. 2000;85(9):3161-3168.
  36. Morin-Papunen L, Rautio K, Ruokonen A, Hedberg P, Puukka M, Tapanainen J. Metformin reduces serum C-reactive protein levels in women with polycystic ovary syndrome. *J Clin Endocrinol Metab*. 2003;88(10):4649-4654.
  37. Morin-Papunen L, Vauhkonen I, Koivunen R, Ruokonen A, Martikainen H, Tapanainen J. Metformin versus ethinyl estradiol-cyproterone acetate in the treatment of nonobese women with polycystic ovary syndrome: a randomized study. *J Clin Endocrinol Metab*. 2003;88(1):148-156.
  38. Rautio K, Tapanainen J, Ruokonen A, Morin-Papunen L. Effects of metformin and ethinyl estradiol-cyproterone acetate on lipid levels in obese and non-obese women with polycystic ovary syndrome. *Eur J Endocrinol*. 2005;152(2):269-275.
  39. Moro F, Morciano A, Tropea A, et al. Effects of drospirenone-ethinylestradiol and/or metformin on CD4(+) CD28(null) T lymphocytes frequency in women with hyperinsulinemia having polycystic ovary syndrome: a randomized clinical trial. *Reprod Sci*. 2013;20(12):1508-1517.
  40. Panidis D, Georgopoulos NA, Piouka A, et al. The impact of oral contraceptives and metformin on anti-müllerian hormone serum levels in women with polycystic ovary syndrome and biochemical hyperandrogenemia. *Gynecol Endocrinol*. 2011;27(8):587-592.
  41. Allen HF, Mazzoni C, Heptulla RA, et al. Randomized controlled trial evaluating response to metformin versus standard therapy in the treatment of adolescents with polycystic ovary syndrome. *J Pediatr Endocrinol Metab*. 2005;18(8):761-768.
  42. Al-Zubeidi H, Klein KO. Randomized clinical trial evaluating metformin versus oral contraceptive pills in the treatment of adolescents with polycystic ovarian syndrome. *J Pediatr Endocrinol Metab*. 2015;28(7-8):853-858.
  43. Essah P, Arrowood J, Cheang K, Adawadkar S, Stovall D, Nestler J. Effect of combined metformin and oral contraceptive therapy on metabolic factors and endothelial function in overweight and obese women with polycystic ovary syndrome. *Fertil Steril*. 2011;96(2):501-505.
  44. Hoeger K, Davidson K, Kochman L, Cherry T, Kopin L, Guzik DS. The impact of metformin, oral contraceptives, and lifestyle modification on polycystic ovary syndrome in obese adolescent women in two randomized, placebo-controlled clinical trials. *J Clin Endocrinol Metab*. 2008;93(11):4299-4306.
  45. Aghamohammadzadeh N, Aliasgarzadeh A, Baglar L, et al. Comparison of metformin and cyproterone-estrodial compound effect on HS C-reactive protein and serum androgen levels in patients with polycystic ovary syndrome. *Pak J Med Sci*. 2010;2(26):347-351.
  46. Bilgir O, Kebapcilar L, Taner C, et al. The effect of ethinylestradiol (EE)/cyproterone acetate (CA) and EE/CA plus metformin treatment on adhesion molecules in cases with polycystic ovary syndrome (PCOS). *Intern Med*. 2009;48(14):1193-1199.
  47. Bodur S, Dundar O, Kanat-Pektas M, Kinci MF, Tutuncu L. The effects of different therapeutic modalities on cardiovascular risk factors in women with polycystic ovary syndrome: a randomized controlled study. *Taiwan J Obstet Gynecol*. 2018;57(3):411-416.
  48. Cetinkalp S, Karadeniz M, Erdogan M, Ozgen G, Saygili F, Yilmaz C. The effects of rosiglitazone, metformin, and estradiol-cyproterone acetate on lean patients with polycystic ovary syndrome. *Endocrinologist*. 2009;19(3):94-97.
  49. Maghraby HA E, Nafee T, Guiziry D, Elnashar A. Randomized controlled trial of the effects of metformin versus combined oral contraceptives in adolescent PCOS women through a 24 month follow up period. *Middle East Fertil Soc J*. 2015;20(3):131-137.
  50. Dorgham N, Sharobim A, Haggag H, El-Kalioby M, Dorgham D. Adding combined oral contraceptives or metformin to laser treatment in polycystic ovarian syndrome hirsute patients. *J Drugs Dermatol*. 2021;20(3):302-306.
  51. Elter K, Imir G, Durmusoglu F. Clinical, endocrine and metabolic effects of metformin added to ethinyl-estradiol-cyproterone acetate in non-obese women with polycystic ovarian syndrome: a randomized controlled study. *Hum Reprod*. 2002;17(7):1729-1737.
  52. Kaya M, Calapkorur B, Karaca Z, et al. The effects of treatment with drospirenone/ethinyl oestradiol alone or in combination with metformin on elastic properties of aorta in women with polycystic ovary syndrome. *Clin Endocrinol*. 2012;77(6):885-892.

53. Kaya M, Yildirim S, Calapkorur B, Akpek M, Unluhizarci K, Kelestimur F. Metformin improves endothelial function and carotid intima media thickness in patients with PCOS. *Gynecol Endocrinol.* 2015;31(5):401-405.
54. Kebapcilar L, Yuksel A, Bozkaya G, et al. Effects of an EE/CA compared with EE/CA-metformin on serum ADMA levels in women with polycystic ovary syndrome. *Cent Eur J Med.* 2009;4(4):423-427.
55. Kebapcilar L, Taner C, Kebapcilar A, Alacacioglu A, Sari I. Comparison of four different treatment regimens on coagulation parameters, hormonal and metabolic changes in women with polycystic ovary syndrome. *Arch Gynecol Obstet.* 2010;281(1):35-42.
56. Kilic S, Yilmaz N, Zulfikaroglu E, Erdogan G, Aydin M, Batioglu S. Inflammatory-metabolic parameters in obese and nonobese normoandrogenemic polycystic ovary syndrome during metformin and oral contraceptive treatment. *Gynecol Endocrinol.* 2011;27(9):622-629.
57. Mhao N, Al-Hilli S, Hadi N, Jamil D, Al-Aubaidy H. A comparative study to illustrate the benefits of using ethinyl estradiol-cyproterone acetate over metformin in patients with polycystic ovarian syndrome. *Diabetes Metab Syndr.* 2015;10(Suppl 1):95-98.
58. Ozgurtas T, Oktenli C, Dede M, et al. Metformin and oral contraceptive treatment reduced circulating asymmetric dimethylarginine (ADMA) levels in patients with polycystic ovary syndrome (PCOS). *Atherosclerosis.* 2008;200(2):336-344.
59. Feng W J, Zhang D, Shi H. Management of polycystic ovarian syndrome with Diane-35 or Diane-35 plus metformin. *Gynecol Endocrinol.* 2016;32(2):147-150.
60. Fonseka S, Wijeyaratne CN, Gawarammana IB, et al. Effectiveness of low-dose ethinylestradiol/cyproterone acetate and ethinylestradiol/desogestrel with and without metformin on hirsutism in polycystic ovary syndrome: a randomized, double-blind, triple-dummy study. *J Clin Aesthet Dermatol.* 2020;13(7):18-23.
61. Kumar Y, Kotwal N, Singh Y, Upreti V, Somani S, Hari Kumar K. A randomized, controlled trial comparing the metformin, oral contraceptive pills and their combination in patients with polycystic ovarian syndrome. *J Family Med Prim Care.* 2018;7(3):551-556.
62. Lv L, Liu Y, Sun Y, Tan K. Effects of metformin combined with cyproterone acetate on clinical features, endocrine and metabolism of non-obese women with polycystic ovarian syndrome. *J Huazhong Univ Sci Technol.* 2005;25(2):194-197.
63. Sahu A, Tripathy P, Mohanty J, Nagy A. Doppler Analysis of ovarian stromal blood flow changes after treatment with metformin versus ethinyl estradiol-cyproterone acetate in women with polycystic ovarian syndrome: a randomized controlled trial. *J Gynecol Obstet Hum Reprod.* 2019;48(5):335-339.
64. Song J, Ruan X, Gu M, Wang L, Wang H, Mueck A. Effect of orlistat or metformin in overweight and obese polycystic ovary syndrome patients with insulin resistance. *Gynecol Endocrinol.* 2018;34(5):413-417.
65. Ruan X, Song J, Gu M, Wang L, Wang H, Mueck A. Effect of Diane-35, alone or in combination with orlistat or metformin in Chinese polycystic ovary syndrome patients. *Arch Gynecol Obstet.* 2018;297(6):1557-1563.
66. Wei W, Zhao H, Wang A, et al. A clinical study on the short-term effect of berberine in comparison to metformin on the metabolic characteristics of women with polycystic ovary syndrome. *Eur J Endocrinol.* 2012;166(1):99-105.
67. Meyer C, McGrath BP, Teede HJ. Effect of medical therapy on insulin resistance and the cardiovascular system in polycystic ovary syndrome. *Diabetes Care.* 2007;30(3):471-478.
68. Hutchison S, Harrison C, Stepto N, Meyer C, Teede H. Retinol-binding protein 4 and insulin resistance in polycystic ovary syndrome. *Diabetes Care.* 2008;31(7):1427-1432.
69. Burchall GF, Piva TJ, Ranasinha S, Teede HJ. Differential effects on haemostatic markers by metformin and the contraceptive pill: a randomized comparative trial in PCOS. *Thromb Haemost.* 2017;117(11):2053-2062.
70. Moran L, Meyer C, Hutchison S, Zoungas S, Teede H. Novel inflammatory markers in overweight women with or without polycystic ovary syndrome and following pharmacological intervention. *J Endocrinol Invest.* 2010;33(4):258-265.
71. Dragoman MV, Tepper NK, Fu R, Curtis KM, Chou R, Gaffield ME. A systematic review and meta-analysis of venous thrombosis risk among users of combined oral contraception. *Int J Gynaecol Obstet.* 2018;141(3):287-294.
72. Heikinheimo O, Toffol E, Partonen T, But A, Latvala A, Haukka J. Systemic hormonal contraception and risk of venous thromboembolism. *Acta Obstet Gynecol Scand.* 2022;101(8):846-855.
73. De Leo V, Musacchio MC, Cappelli V, Massaro MG, Morgante G, Petraglia F. Genetic, hormonal and metabolic aspects of PCOS: an update. *Reprod Biol Endocrinol.* 2016;14(1):38.
74. Alberti KGMM, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 2009;120(16):1640-1645.
75. Lim SS, Kakoly NS, Tan JWJ, et al. Metabolic syndrome in polycystic ovary syndrome: a systematic review, meta-analysis and meta-regression. *Obes Rev.* 2019;20(2):339-352.
76. Melin J, Forslund M, Alesi S, et al. The impact of metformin with or without lifestyle modification versus placebo on polycystic ovary syndrome: a systematic review and meta-analysis of randomized controlled trials. *Eur J Endocrinol.* 2023;lvad098. doi: [10.1093/ejendo/lvad098](https://doi.org/10.1093/ejendo/lvad098)
77. Stepto NK, Cassar S, Joham AE, et al. Women with polycystic ovary syndrome have intrinsic insulin resistance on euglycaemic-hyperinsulaemic clamp. *Hum Reprod.* 2013;28(3):777-784.
78. Korytkowski MT, Mokan M, Horwitz MJ, Berga SL. Metabolic effects of oral contraceptives in women with polycystic ovary syndrome. *J Clin Endocrinol Metab.* 1995;80(11):3327-3334.
79. Piltonen T, Puurunen J, Hedberg P, et al. Oral, transdermal and vaginal combined contraceptives induce an increase in markers of chronic inflammation and impair insulin sensitivity in young healthy normal-weight women: a randomized study. *Hum Reprod.* 2012;27(10):3046-3056.
80. Lopez L, Grimes D, Schulz K. Steroidal contraceptives: effect on carbohydrate metabolism in women without diabetes mellitus. *Cochrane Database Syst Rev.* 2019;2019(11). doi: [10.1002/14651858.CD006133.pub5](https://doi.org/10.1002/14651858.CD006133.pub5)
81. Bastianelli C, Farris M, Rosato E, Brosens I, Benagiano G. Pharmacodynamics of combined estrogen-progestin oral contraceptives: 1. Effects on metabolism. *Expert Rev Clin Pharmacol.* 2017;10(3):315-326.
82. Romero-Cabrera JL, Ankeny J, Fernández-Montero A, Kales SN, Smith DL. A systematic review and meta-analysis of advanced biomarkers for predicting incident cardiovascular disease among asymptomatic middle-aged adults. *Int J Mol Sci.* 2022;23(21):13540.
83. Morin-Papunen L, Rantala AS, Unkila-Kallio L, et al. Metformin improves pregnancy and live-birth rates in women with polycystic ovary syndrome (PCOS): a multicenter, double-blind, placebo-controlled randomized trial. *J Clin Endocrinol Metab.* 2012;97(5):1492-1500.
84. Kangasniemi MH, Arffman RK, Joenväärä S, et al. Ethinylestradiol in combined hormonal contraceptive has a broader effect on serum proteome compared with estradiol valerate: a randomized controlled trial. *Hum Reprod.* 2023;38(1):89-102.
85. Kangasniemi MH, Haverinen A, Luiro K, et al. Estradiol valerate in COC has more favorable inflammatory profile than synthetic ethinyl estradiol: a randomized trial. *J Clin Endocrinol Metab.* 2020;105(7):e2483-e2490.
86. Ollila ME, West S, Keinänen-Kiukaanniemi S, et al. Overweight and obese but not normal weight women with PCOS are at

- increased risk of type 2 diabetes mellitus—a prospective, population-based cohort study. *Hum Reprod.* 2017;32(2):423-431.
87. Forslund M, Landin-Wilhelmsen K, Trimpou P, Schmidt J, Brannstrom M, Dahlgren E. Type 2 diabetes mellitus in women with polycystic ovary syndrome during a 24-year period: importance of obesity and abdominal fat distribution. *Hum Reprod Open.* 2020;1:hoz042.
88. Wu L, Liu Y, Huang X, *et al.* Oral contraceptives (OCs) in combination with metformin versus OCs alone on metabolism in nonobese polycystic ovary syndrome: a meta-analysis and systematic review of randomized controlled trials. *Clin Endocrinol.* 2023;99(1):3-16.