



UNIVERSITY OF HELSINKI



<https://helda.helsinki.fi>

Helda

---

## The effect of beaver facilitation on Common Teal : pairs and broods respond differently at the patch and landscape scales

Nummi, Petri

Wiley Blackwell

2019-04

---

Nummi, P, Suontakanen, E-M, Holopainen, S & Väänänen, V-M 2019, 'The effect of beaver facilitation on Common Teal : pairs and broods respond differently at the patch and landscape scales', *Ibis*, vol. 161, no. 2, pp. 301-309. <https://doi.org/10.1111/ibi.12626>

---

<http://hdl.handle.net/10138/302629>

10.1111/ibi.12626

---

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

**The effect of beaver facilitation on Common Teal: pairs and broods respond differently at the patch and landscape scales**

PETRI NUMMI\*, EEVA-MARIA SUONTAKANEN, SARI HOLOPAINEN & VELI-MATTI VÄÄNÄNEN

*Department of Forest Sciences, University of Helsinki, P.O. Box 27, 00014 University of Helsinki, Finland*

\*Corresponding author.

Email: [petri.nummi@helsinki.fi](mailto:petri.nummi@helsinki.fi)

+358 05 4486160

17 Avian species respond to ecological variability at a range of spatial scales and  
18 according to life history stage. Beaver dams create wetland systems for waterbirds  
19 that are utilized throughout different stages of the breeding season. We studied  
20 how beaver-induced variability affected mobile pairs and more sedentary broods  
21 along with the production of Common Teal *Anas crecca* at the patch and landscape  
22 scale on their breeding grounds. Beavers *Castor* spp. are ecosystem engineers that  
23 enhance waterfowl habitats by impeding water flow and creating temporary  
24 flooding. Two landscapes in southern Finland with (Evo) and without (Nuksio)  
25 American Beavers *Castor canadensis* were used in this study. To investigate the  
26 patch-scale effect, pair and brood densities along with brood production were first  
27 compared at beaver-occupied lakes and non-beaver lakes in the beaver landscape.  
28 Annual pair and brood densities/km shoreline and brood production were compared  
29 between beaver and non-beaver landscapes. Facilitative effects of beaver activity  
30 were manifest on brood density at both patch and landscape scales: these were  
31 over 90 and 60 percent higher in beaver patches and landscapes, respectively. An  
32 effect of beaver presence on pair density was only seen at the landscape level. Pair  
33 density did not strongly affect brood production, as shown earlier for relatively  
34 mildly density-dependent Teal populations. Because the extent of beaver flooding  
35 was a crucial factor affecting annual Teal production in the study area, we infer  
36 beaver activity has consequences for the local Teal population. Ecosystem  
37 engineering by the beaver could therefore be considered as a restoration tool in  
38 areas where waterfowl are in need of high-quality habitats.

39

40 **Keywords:** *Anas crecca*, brood production, *Castor canadensis*, ecosystem engineer,  
41 experienced scale, Teal density, wetland management

42

43

44 Avian habitat selection, population stability and response to disturbance may differ  
45 according to the scale at which these are investigated (Wiens 1989a, Denny *et al.*  
46 2004). Schneider (2001) identifies three scale-related problems which arise when  
47 ecological processes are studied. First, many ecological problems are often large in  
48 scale. Second, most variables can only be measured in small areas. Third, patterns or  
49 processes at small scales do not necessarily hold or prevail at larger scales (but see  
50 Denny *et al.* 2004). Sometimes the mechanisms underlying observed patterns  
51 operate at different scales than those at which the patterns are detected. Even  
52 investigators addressing the same questions, but at different scales, have been  
53 observed to reach differing conclusions (Wiens 1989b). Examples concerning birds  
54 include Least Flycatchers *Empidonax minimus* negatively influencing the distribution  
55 of American Redstart *Setophaga ruticilla* through interspecific aggression at the  
56 scale of four-hectare forest plots. At the regional level, however, the total  
57 abundances of the two species correlated positively (Sherry & Holmes 1988).  
58 Similarly, variation in habitat use by sedentary Pied-billed Grebe *Podilymbus*  
59 *podiceps* was explained solely by within-patch variation whereas area requirements  
60 for vagile Black Tern *Chlidonias niger*, which forages up to four km away from the  
61 nest wetland, fluctuated in response to landscape structure (Naugle *et al.* 1999).

62 Beavers *Castor* spp. are ecosystem engineers of the boreal forest, creating  
63 disruption to riparian areas by their damming of creeks or ponds and forming of  
64 inundated areas (Naiman *et al.* 1988, Wright *et al.* 2003, Hyvönen & Nummi 2008).  
65 Both American *Castor canadensis* and Eurasian Beaver *C. fiber* are assumed to have  
66 a similar effect on the riparian ecosystem (Danilov & Fyodorov 2015) that would  
67 otherwise not exist in the landscape (Remillard *et al.* 1987, Johnston & Naiman  
68 1990, Nummi & Kuuluvainen 2013). Beaver patches undergo succession from  
69 terrestrial to aquatic habitat and back, thereby increasing landscape heterogeneity  
70 (Remillard *et al.* 1987). Via ecosystem engineering, beavers may also act as  
71 facilitators for various species, such as plants (Wright *et al.* 2002), butterflies (Bartel  
72 *et al.* 2010), waterbirds (Nummi & Holopainen 2014), and bats (Nummi *et al.* 2011).  
73 Therefore using beaver as a restoration tool for freshwater habitats has been  
74 suggested (Törnblom *et al.* 2011, Law *et al.* 2016).

75 Among boreal ducks, patch creation or modification by beavers has been found to  
76 especially benefit Common Teal *Anas crecca*, (hereafter Teal), as both Teal pairs and  
77 broods rapidly occupy newly formed beaver flowages (Nummi & Pöysä 1997,  
78 Nummi & Hahtola 2008). This has been attributed to both habitat modification and  
79 resource enhancement (*sensu* Bruno *et al.* 2003): beaver ponds provide more  
80 invertebrates that are potential food for Teal and have shallower depths – suitable

81 for foraging – than non-disturbed boreal ponds (Longcore *et al.* 2006, Nummi &  
82 Hahtola 2008). Compared to non-flooded lakes, Teal densities overall tend to be  
83 higher on beaver ponds due to greater numbers of pairs and broods, as well as  
84 elevated Teal duckling survival with ample foods there (Nummi & Hahtola 2008).

85 As is the case for most organisms, ducks may experience their environment  
86 differently according to age, i.e. as a mobile flying adult compared to a more  
87 sedentary flightless duckling (Levin 1992). In our study we investigate how Teal pairs  
88 and unfledged broods (which clearly differ in their mobility) respond to  
89 environmental variation at patch (0.1–50 ha) versus landscape scales (40–50 km<sup>2</sup>).  
90 The patches in our study are easily covered by both breeding pairs and broods  
91 within tens of minutes or 1–2 hours of swimming. In contrast, shifting between the  
92 most distant lakes within each landscape of our study would only take a few  
93 minutes for the flying pairs, but at least 15 hours (but most likely 1–3 days) for the  
94 non-flying broods (Martin & Forsyth 1983, Duncan 1993). Duck pairs therefore  
95 typically have home ranges comprising many small wetlands and they travel a few  
96 kilometers between their foraging patches (Gilmer *et al.* 1975, Guillemain & Elmberg  
97 2014). We thus predict that the effect of spatial scale will differ between pairs and  
98 broods.

99 We hypothesize that Teal broods would be affected by beaver inundation already at  
100 the patch scale, whereas the beaver effect would be clearly seen in pairs at the  
101 landscape scale.

102

103

## 104 **METHODS**

105

### 106 **Study area**

107 Data were gathered in an oligotrophic watershed (area 39 km<sup>2</sup>) at Evo inhabited by  
108 American Beaver (hereafter Beaver) (61°12' N, 25°07' E; 51 lakes) and the Nuuksio  
109 lake area (53 km<sup>2</sup>; 60°19'N, 24°28'E; 54 lakes) in southern Finland not inhabited by  
110 Beavers (Arvola *et al.* 2010, Nyberg *et al.* 2010). The study lakes of both areas are  
111 oligotrophic, relatively small (Nuuksio 0.2–94.5 ha, Evo 0.1–49.5 ha), have fish  
112 populations and are closed or headwater lakes (Väänänen *et al.* 2012). The predator  
113 communities of the two areas are similar with Pine Marten *Martes martes* and Red  
114 Fox *Vulpes vulpes* as main predators (Natural Resources Institute Finland 2017)

115 along with two alien predators, the American Mink *Neovison vison* and Raccoon Dog  
116 *Nyctereutes procyonoides*. Ducks are additionally predated by several birds of prey  
117 and the Northern Pike *Esox lucius*.

118 Boreal forest covers most of the two areas, interspersed with lakes and mires.  
119 Agriculture and human settlement are very limited and local. Apart from Beaver-  
120 created variability in Evo, landscape-level habitat structure of the Evo and Nuuksio  
121 lakes has been fairly stable for the last 20 years (Suhonen *et al.* 2011, Thompson *et al.*  
122 *et al.* 2016). In Evo, Beavers change their lake of occupancy on average every three  
123 years, thus new Beaver habitat patches are created continuously while old ones are  
124 abandoned (Hyvönen & Nummi 2008). Lake shores in both study areas are generally  
125 steep, with little emergent vegetation (mainly sedges *Carex* spp. and Common Reed  
126 *Phragmites australis*), although some lakes have stands of Water Horsetail  
127 *Equisetum fluviatile* and cattail *Typha* spp. Emergent vegetation is usually lined with  
128 narrow belts of Yellow Water Lilies *Nuphar lutea* and Water Lilies *Nymphaea*  
129 *candida*; submerged vegetation is very sparse. Rocky shores are also typical in  
130 Nuuksio. In Beaver ponds the shores are significantly shallower than non-Beaver  
131 ponds and contain inundated herbaceous vegetation and bushes (Nummi & Hahtola  
132 2008). Located in a cool continental climate, all wetlands freeze over from  
133 November to April. Each spring, they are thus colonized anew by migrating ducks.  
134 Overall, the lakes of our two study areas were very similar and have been previously  
135 combined in waterbird studies (Nummi *et al.* 2012; Väänänen *et al.* 2012).

136

137

### 138 **Teal data**

139 In the Beaver landscape Evo duck data are from 1988–2014 and in non-Beaver  
140 landscape Nuuksio data are from 1994–2014. The number of pairs was estimated in  
141 both study landscapes once a year between the end of April and the beginning of  
142 May. The number of nesting pairs was interpreted using the survey guidelines of the  
143 Finnish Museum of Natural History (Koskimies & Pöysä 1991). Brood data from the  
144 Beaver landscape were collected from 1988–2008 by conducting five annual brood  
145 surveys from June to August, and from 2009–2014 by conducting two annual  
146 surveys from June to July. In the non-Beaver landscape, the first brood survey was  
147 conducted in June and the second round in July. During each survey, a point count  
148 was first made from the shore, after which a round count was performed by circling  
149 the lakes by foot or boat (Nummi & Pöysä, 1993); the method was chosen based on  
150 lake size and shoreline structure. In the most densely vegetated Beaver ponds all  
151 three methods (point count, circling by foot and circling by boat) were combined to  
152 attain coverage of the entire pond area. In Beaver ponds the ducks are most difficult

153 to detect (see Holopainen *et al.* 2014), thus the number of birds found there are  
154 conservative. In both landscapes, all wetlands inside a certain area were surveyed.

155 Brood surveys included broods, ducklings without a female and also females  
156 emitting alarm signals. Duckling number was counted for every observed brood,  
157 after which the brood age was determined following the seven age classes of Pirkola  
158 and Högmander (1974). Age class was also estimated for ducklings seen without an  
159 adult female.

160 To eliminate the error caused by the irregularity in the number of annual brood  
161 counts, brood density was counted per survey in both study landscapes. In the  
162 comparison of Beaver and non-Beaver landscapes, we only took into account the  
163 two surveys (in June and July) in the Beaver landscape which corresponded to those  
164 of the non-Beaver landscape. Brood production was resolved by counting the  
165 number of broods per adult pair. The different age classes of the ducklings were not  
166 taken into account, but observations of juveniles were excluded. Thus, all sightings  
167 of the six age classes were taken into account, but they were not separated from  
168 one another.

169 Pairs and brood densities are given per km shoreline, thus information on shoreline  
170 length of the lakes and ponds in the study areas were required. Lake shore lengths  
171 in the Beaver study area were already measured by Nummi and Pöysä (1993). The  
172 shore lengths for the non-Beaver lakes and ponds, excluding a few small ponds,  
173 were obtained from the Finnish Environment Institute's open data service  
174 ([syke.fi/avoindata](http://syke.fi/avoindata)). Shoreline length was measured for the remaining ponds using  
175 the same database's map service "Karpalo". Many lakes and ponds in the non-  
176 Beaver study area have a partly rocky shoreline, which is an unsuitable breeding  
177 environment for ducks. This amount of rocky shoreline was removed from the total  
178 shoreline length during analyses. Lengths of the rocky shorelines were measured  
179 using the map service of the Finnish Environment Institute ([syke.fi/avoindata](http://syke.fi/avoindata)).

180

## 181 **Beaver data**

182 Both European and American Beavers were introduced to Evo during the 1930s and  
183 1950s (Lahti & Helminen 1974). Nowadays the Beaver population in Evo is  
184 comprised solely of American Beavers (Parker *et al.* 2012).

185 Since 1976, Beaver data have been collected from the lakes and ponds flooded by  
186 Beavers. The data indicate the years during which Beavers have flooded a lake or a  
187 pond. In Evo, Beaver ponds are most commonly formed by the damming of an

188 existing pond (Nummi & Hahtola 2008). Beavers occupy one site for an average of  
189 three years, and often return to the same areas they have flooded before (Hyvönen  
190 & Nummi 2008).

191

## 192 **Patch-scale comparison of teal abundance before and during flooding**

193 On the patch scale we studied whether the Beaver increases pair and brood  
194 densities and brood production. The study was conducted in the Beaver area by  
195 comparing specific variables before and during a Beaver inundation in a patch. The  
196 goal of the comparison was to collect data two years before and two years during  
197 the flood. The density of only one year was used if Beaver-induced flooding had  
198 occurred for only one year in the patch.

199 Pair density was compared in 19 different lakes before and during Beaver  
200 inundation, and brood densities were similarly examined in 18 lakes. Brood  
201 production was also examined in 12 different lakes. These different figures result  
202 from the fact that in some cases the Beaver flood was only present in a certain lake  
203 during the pair time, and in some cases during the brood time, and in 12 cases both  
204 parameters were available. In each of these three cases, a control lake was included  
205 for each Beaver lake, representing the most similar lake along an environmental  
206 gradient of habitat luxuriance (Nummi & Pöysä 1997). In control lakes especially the  
207 amount of broods remained extremely low which is the normal pattern in boreal  
208 lakes not affected by Beaver (Elmberg *et al.* 2005, Gunnarsson *et al.* 2004). The  
209 environmental gradient takes into account the shore's vegetation type, the amount  
210 of vegetation, water depth at the shore and lake size (Suhonen *et al.* 2011). The  
211 comparison was made using the Wilcoxon signed-rank test, which is a non-  
212 parametric version of the pairwise *t*-test (Ranta *et al.* 2012). The examination was  
213 made using the IBM SPSS Statistics 21 software (IBM Corp. 2012).

214

## 215 **Comparison of teal density in Beaver and non-Beaver lakes at Evo**

216 We also studied whether pair and brood densities and brood production differed in  
217 lakes with and without Beaver in the Beaver landscape. Lakes where data  
218 concerning Beaver floods were available during the study period were included in  
219 the data set. Altogether 24 Beaver-flooded lakes and 18 lakes with no Beaver  
220 activity during years 1988–2014 were included in the examination. We first tried to  
221 perform a glmm analysis using the individual lakes, and after data exploration ended  
222 up trying hurdle models (ZAP for broods and ZANB for pairs) with a random lake-

223 effect. Unfortunately the lake level analysis could not be done with the glmm hurdle  
224 models, apparently because of so many zeros in both the Beaver occurrence column  
225 and teal column. Instead, the comparison was performed using the years when  
226 Beaver-induced floods occurred in the Beaver lakes, while all the years of the study  
227 period were used in the non-Beaver lakes. Annual average pair density was counted  
228 for each lake. Brood density was obtained using the years from 1988 to 2014. The  
229 same years were also used in examining pair density and brood production,  
230 excluding year 1988, from which no data on pair numbers were available. The  
231 comparison was made using the independent samples Mann-Whitney *U*-test by  
232 using the IBM SPSS Statistics 21 software (IBM Corp. 2012).

233

### 234 **Beaver and non-Beaver areas – comparing two landscapes**

235 We compared the annual brood and pair densities and brood production for 1994–  
236 2014 between the Beaver and non-Beaver landscapes. All pair and brood sightings  
237 of one year were added together, and brood density and broods per pair figures  
238 were transformed into figures per one survey. The comparison was made using the  
239 Mann-Whitney *U*-test, because the sightings were not normally distributed.  
240 Comparisons were performed using the IBM SPSS Statistics 21 software (IBM Corp.  
241 2012). In addition to the difference between the study areas, we explored whether  
242 annual population variation differs within these two study areas. This annual  
243 variation was examined by testing the homogeneity of the variances with Fligner-  
244 Killeen's test. Fligner-Killeen's test was chosen, because it also works well with  
245 samples that are not normally distributed (Ranta *et al.* 2012). Fligner-Killeen's test  
246 was conducted using the variation observed by Conover *et al.* (1981), in which the  
247 rank is counted using the median instead of plain values. The test was performed  
248 using R 2.3.2 software (R Core Team 2015).

249

250

## 251 **RESULTS**

252

### 253 **Patch scale**

254 Teal brood density in the ponds increased significantly with Beaver inundation ( $Z = -$   
255  $2.24$ ,  $P = 0.025$ ,  $n = 18$ , Fig. 1a), as did the median number of broods per pair ( $Z = -$   
256  $2.02$ ,  $P = 0.043$ ,  $n = 12$ ). Pair density, however, did not increase significantly ( $Z = -1.6$ ,

257  $P = 0.110$ ,  $n = 19$ ). In the non-flooded control lakes no significant change was found  
258 in either of the three cases ( $P > 0.10$  in all cases, Fig. 1b).

259

### 260 **Within landscape**

261 Within the Beaver landscape, Teal brood density in Beaver ponds was again  
262 significantly higher than that of non-Beaver ponds, as was the number of broods  
263 produced per pair (Table 1). The pair density of Beaver and non-Beaver ponds did  
264 not differ significantly (Table 1).

265

### 266 **Between landscapes**

267 Teal pair density ( $U = -5.18$ ,  $P < 0.001$ , Fig. 2a), brood density ( $U = -4.24$ ,  $P = < 0.001$ ,  
268 Fig. 2b), as well as the number of broods per pair ( $U = -2.25$ ,  $P = 0.012$ , Fig. 2c) were  
269 significantly higher in the Beaver landscape (median: 0.34 pairs/ shoreline km, 0.03  
270 broods/km, 0.11 broods/pair) than in the non-Beaver landscape (0.15 pairs/km, 0.01  
271 broods/km, 0.08 broods per pair). No significant differences were observed in the  
272 variability of pair or brood production per pair between the Beaver and non-Beaver  
273 landscapes (Fligner-Killeen test,  $P > 0.05$  in both cases). Brood density variability was  
274 higher in the Beaver landscape ( $\chi^2 = 7.24$ ,  $P = 0.007$ ).

275

276

## 277 **DISCUSSION**

278

279 We found scale-dependent patterns in the effects of Beaver on breeding Teal. Our  
280 study corroborated earlier observations that Beaver ponds increase the number of  
281 Teal broods at the patch scale (Nummi & Hahtola 2008), the brood densities being  
282 over 90 % higher during Beaver inundation. This finding was coupled with an  
283 increase in the number of broods produced per pair. The increase in Teal pairs upon  
284 Beaver flooding was not significant, contrasting the earlier study by Nummi & Pöysä  
285 (1997). The results from these experimental-like situations of before and during  
286 inundation were supported when Beaver ponds and non-Beaver ponds within the  
287 Beaver landscape were compared: the densities of pairs and broods along with  
288 breeding success were eight to nine times higher in Beaver ponds. These  
289 observations fit the conceptual framework created by Levin (1992). He stated that

290 localized disturbances play an important role in the maintenance of species such as  
291 the Teal which benefit from disrupted, high-quality habitats; in the case of Teal  
292 habitats with abundant invertebrates and shallow vegetated shores suitable for  
293 foraging (Nummi *et al.* 2013, Nummi & Hahtola 2008).

294 When the Beaver and non-Beaver landscapes were compared, a similar picture  
295 emerged. Both the number of pairs and broods were significantly higher in the  
296 Beaver-influenced landscape than in the one which lacked Beavers. In the Beaver  
297 landscape the pair density was approximately 50 % and brood density 60 % higher  
298 than those found in the non-Beaver landscape. Similarly, the average number of  
299 broods per pair was higher in the Beaver landscape. We of course had only one  
300 landscape pair to compare, so possible other differences between the landscapes  
301 could not be effectively controlled. There appears to be no pronounced difference in  
302 general productivity between the two areas, since the densities of two other  
303 common ducks, Mallard *Anas platyrhynchos* and Common Goldeneye *Bucephala*  
304 *clangula* were similar in Evo (Nummi & Pöysä 1995a) and Nuuksio (Taskinen 1997).

305 We found that the Beaver effect does not appear to emerge as clearly at the patch  
306 scale for pairs as it does for broods. However, pair density was also higher with  
307 Beaver influence at the landscape level. Once a pattern is detected, it is possible to  
308 concentrate on measuring its determinants and the mechanisms behind it (Levin  
309 1992). In our case (at least) two mechanisms could bring about the pattern we  
310 observed. Firstly, although pairs might use Beaver ponds for foraging, they very  
311 likely encompass many small wetlands in their home range, and spend only part of  
312 their time in the Beaver flowages. This pattern has been shown for radio-tracked  
313 Mallards with home ranges of approximately 230 ha that include e.g. seasonal  
314 wetlands (Gilmer *et al.* 1975, see also Dwyer *et al.* 1979); individual breeding Teals  
315 have not been followed using radio telemetry but the home ranges of Teal pairs are  
316 approximately the same size as those of Mallard (Nudds & Ankney 1982). Less  
317 mobile broods, again, may spend all of their time in the optimal Beaver ponds, once  
318 they have been discovered (Nummi & Hahtola 2008; the home ranges of  
319 comparable Mallard broods were 10–15 ha; Talent *et al.* 1982, Chouinard *et al.*  
320 2007). Secondly, and possibly more importantly, other flooded areas in riparian  
321 forests, namely seasonal wetlands created by melting snow, have been shown to be  
322 suitable for foraging during the pairing season (Paton 2005, see also Holopainen *et*  
323 *al.* 2014). This may dilute the Beaver flowage effect during springs when large  
324 amounts of flooded shores and vernal pools are available.

325 At the landscape scale Teal broods were concentrated in the few Beaver patches of  
326 the Beaver landscape of Evo (Nummi & Pöysä 1995b), while pairs were less so  
327 (Elmberg *et al.* 2005). The amount of Beaver flooding was also the most important  
328 factor affecting yearly Teal production in the area, among such environmental  
329 factors as food abundance (Holopainen *et al.* 2014). This could also be the reason  
330 why the brood density varied more in the Beaver landscape. Moreover, our finding  
331 that the average number of broods per pair was higher in the higher density Beaver  
332 landscape than in the lower density non-Beaver landscape corroborates our earlier  
333 study indicating that Teal populations may not show very strong spatial density  
334 dependence (Nummi *et al.* 2015). This supports the hypothesis that considering  
335 environmental variability, an association is often visible with a species' life history  
336 adaptation along a "slow-fast continuum" and temporary vs. permanent habitats  
337 (Fowler 1981, Sæther & Engen 2002). "Fast" species typically live in habitats that are  
338 either unpredictable in time or short-lived, and those species themselves are often  
339 short-lived as well (Sæther & Engen 2002).

340 Considering that 60–90 % of European wetlands were lost during the last century  
341 and that similar figures apply to many areas of North America, there is great need  
342 for wetland restoration (Junk *et al.* 2013). From geomorphological, hydrological and  
343 ecological aspects introducing Beavers has the potential to be a mechanism for  
344 wetland restoration (Hey & Philippi 1995, Burchsted 2010, Törnblom *et al.* 2011).  
345 Beavers are known to enhance biodiversity by beneficially affecting numerous  
346 groups of organisms (Stringer & Gaywood 2016), sometimes facilitating whole  
347 species communities (Dalbeck *et al.* 2007, Nummi & Holopainen 2014). Our data  
348 support earlier findings which consider Beavers as a feasible and economic way of  
349 restoring riparian landscapes (Brown & Parsons 1979, Nummi 1989, Thompson *et al.*  
350 2016).

351

352 We thank Tony Fox and two anonymous referees for valuable comments on the manuscript, and  
353 the numerous field workers for their help in the lakes. The grant from the Maj and Tor Nessling  
354 Foundation to S. H. and from the Foundation for Research of Natural Resources in Finland to P. N.  
355 are much appreciated.

356

## 357 REFERENCES

358

- 359 Arvola, L., Rask, M., Ruuhijärvi, J., Tulonen, T., Vuorenmaa, J., Ruoho-Airola, T. &  
360 Tulonen, J. 2010. Long-term patterns in pH and colour in small acidic boreal lakes of  
361 varying hydrological and landscape settings. *Biogeochem.* 101: 269–279.
- 362 Bartel, R.A., Haddad, N.M. & Wright, J.P. 2010. Ecosystem engineers maintain a rare  
363 species of butterfly and increase plant diversity. *Oikos* 119: 883–890.
- 364 Brown, M.K. & Parsons, G.R. 1979. Waterfowl production on beaver flowages in a  
365 part of New York. *New York Fish and Game Journal* 26: 142–153.
- 366 Bruno, J.F., Stachovicz, J.J. & Bertness M.D. 2003. Inclusion of facilitation into  
367 ecological theory. *Trends Ecol. Evol.* 18: 119–125.
- 368 Burchsted, D., Daniels, M., Thorson, R. & Vokoun, J. 2010. The River Discontinuum:  
369 Applying Beaver Modifications to Baseline Conditions for Restoration of Forested  
370 Headwaters. *BioScience* 60: 908–922.
- 371 Chouinard Jr., M.P & Arnold, T.W. 2007. Survival and habitat use of mallard (*Anas*  
372 *platyrhynchos*) broods in the San Joaquin Valley, California. *The Auk* 124: 1305–  
373 1316.
- 374 Conover, W.J., Johnson, M.E. & Johnson, M.M. 1981. A Comparative Study of Tests  
375 for Homogeneity of Variances, with Applications to the Outer Continental Shelf  
376 Bidding Data. *Technometrics* 23: 351–361.
- 377 Dalbeck, L., Lüscher, B. & Ohlhoff, D. 2007. Beaver ponds as habitat of amphibian  
378 communities in a central European highland. *Amphibia-Reptilia* 28: 493–501.
- 379 Danilov, P.I. & Fyodorov, F.V. 2015. Comparative characterization of the building  
380 activity of Canadian and European beavers in northern European Russia. *Russian J.*  
381 *Ecol.* 46: 272–278.
- 382 Denny, M.W., Helmuth, B., Leonard, G.H., Harley, C.D.G., Hunt, L.J.H. & Nelson, E.K.  
383 2004. Quantifying scale in ecology: lessons from a wave-swept shore. *Ecological*  
384 *Monographs* 74: 513–532.
- 385 Duncan, D.C. 1983. Extensive overland movement of Pintail, *Anas acuta*, brood and  
386 attempted predation by hawks. *Can. Field-Naturalist* 97: 216–217.
- 387 Dwyer, T.J., Krapu, G.L. & Janke, D.M. 1979. Use of prairie pothole habitat by  
388 breeding Mallards. *J. Wildl. Manage.* 43: 526–31.

- 389 Elmberg, J., Nummi, P., Pöysä, H., Gunnarsson, G. & Sjöberg, K. 2005. Early breeding  
390 Teal *Anas crecca* use the best lakes and have the highest reproductive success. *Ann.*  
391 *Zool. Fennici* 42: 37–43.
- 392 Fowler, C.W. 1981. Density dependence as related to life history strategy. *Ecology*  
393 62: 602–610.
- 394 Gilmer, D.S., Ball, I.J., Cowardin, L.M., Riechmann, J.H. & Tester, J.R. 1975. Habitat  
395 Use and Home Range of Mallards Breeding in Minnesota. *J. Wildl. Manage.* 39: 781–  
396 789.
- 397 Guillemain, M. & Elmberg, J. 2014. *The Teal*. T & AD Poyser, London.
- 398 Gunnarsson, G., Elmberg, J., Sjöberg, K., Pöysä, H. & Nummi, P. 2004. Why are there  
399 so many empty lakes? Food limits survival of mallard ducklings. *Can. J. Zool.* 82:  
400 1698–1703.
- 401 Hey, D.L. & Philippi, N.S., 1995. Flood reduction through wetland restoration: the  
402 upper Mississippi River Basin as a case history. *Restor. Ecol.* 3: 4–17.
- 403 Holopainen, S., Nummi, P. & Pöysä, H. 2014. Breeding in the stable boreal  
404 landscape: Lake habitat variability drives brood production in the teal (*Anas crecca*).  
405 *Freshwat. Biol.* 59: 2621–2631.
- 406 Hyvönen, T. & Nummi, P. 2008. Habitat dynamics of beaver *Castor canadensis* at  
407 two spatial scales. *Wildl. Biol.* 14: 302–308.
- 408 Johnston, C.A. & Naiman, R.J. 1990. Aquatic patch creation in relation to beaver  
409 population trends. *Ecology* 71: 1617–1621.
- 410 Junk, W.J., An, S., Finlayson, C.M., Gopal, B., Květ, J., Mitchell, S.A., Mitsch, W.J. &  
411 Robarts, R.D. 2013. Current state of knowledge regarding the world's wetlands and  
412 their future under global climate change: a synthesis. *Aquatic Sci.* 75: 151–167.
- 413 Koskimies, P. & Pöysä, H. 1991. Waterfowl point count. In: Koskimies, P. & Väisänen,  
414 R.A. (eds) *Monitoring bird populations* Zoological Museum, Finnish Museum of  
415 Natural History, Helsinki: 41–44.
- 416 Lahti, S. & Helminen, M. 1974. The beaver *Castor fiber* (L.) and *C. canadensis* (Kuhl)  
417 in Finland. *Acta Ther.* 19: 177–189.
- 418 Law, A., Mclean, F. & Willby, N. 2016. Habitat engineering by beaver benefits  
419 aquatic biodiversity and ecosystem processes in agricultural streams. *Freshwat. Biol.*  
420 61: 486–499.

- 421 Levin, S.A. 1992. The problem of pattern and scale in ecology. *Ecology* 73: 1943–  
422 1967.
- 423 Levin, S.A. & Paine, R.T. 1974. Disturbance, patch formation, and community  
424 structure. *Proc. Nat. Acad. Sci.* 71: 2744–2747.
- 425 Longcore, J.R., McAuley, D.G., Pendelton, G.W., Bennatti, C.R., Mingo, T.M. &  
426 Stromborg, K. L. 2006. Macroinvertebrate abundance, water chemistry, and wetland  
427 characteristics affect use of wetlands by avian species in Maine. *Hydrobiologia* 567:  
428 143–167.
- 429 Martin, P.A. & Forsyth, D.J. 1993. Survival and behaviour of captive Mallard broods  
430 exposed to carbofuran-sprayed vegetation: a field experiment. *Ecotoxicology* 2: 79–  
431 92.
- 432 Naiman, R.J., Johnston, C.A. & Kelley, J.C. 1988. Alteration of North American  
433 streams by beaver: the structure and dynamics of streams are changing as beaver  
434 recolonize their historic habitat. *BioScience* 38: 753–762.
- 435 Natural Resources Institute Finland 2017. <https://www.riistakolmiot.fi/raportit>  
436 (read 30th April, 2017).
- 437 Naugle, D.E., Higgins, K.F., Nusser, S.M. Johnson, W.C. 1999. Scale-dependent  
438 habitat use in three species of prairie wetland birds. *Landscape Ecology* 14: 267–  
439 276.
- 440 Nudds, T.D. & Ankney, C.D. 1982. Ecological correlates of territory and home range  
441 size in North American dabbling ducks. *Wildfowl* 33: 58–62.
- 442 Nummi P. 1989. Simulated effects of the beaver on vegetation, invertebrates and  
443 ducks *Ann. Zool. Fennici* 26: 43–52.
- 444 Nummi, P. & Hahtola, A. 2008. The beaver as an ecosystem engineer facilitates teal  
445 breeding. *Ecography* 31: 519–524.
- 446 Nummi, P. & Holopainen, S. 2014. Whole-community facilitation by beaver:  
447 ecosystem engineer increases waterbird diversity. *Aquatic Conserv.* 24: 623–633.
- 448 Nummi, P, Holopainen, S., Rintala, J. & Pöysä, H. 2015. Mechanisms of density  
449 dependence in ducks: importance of space and per capita food. *Oecologia* 177: 679–  
450 688.
- 451 Nummi P., Kattainen S., Ulander P. & Hahtola A. 2011: Bats benefit from beavers: a  
452 facilitative link between aquatic and terrestrial food webs. *Biod. Cons.* 20: 851–859.

- 453 Nummi P. & Kuuluvainen, T. 2013. Forest disturbance by an ecosystem engineer:  
454 beaver in boreal landscapes. *Boreal Environ. Res.* 18: 13–24.
- 455 Nummi, P., Paasivaara, A., Suhonen, S. & Pöysä, H. 2013. Wetland use by brood-  
456 rearing female ducks in a boreal forest landscape: the importance of food and  
457 habitat. *Ibis* 155: 68–79.
- 458 Nummi, P. & Pöysä, H. 1993. Habitat associations of ducks during different phases of  
459 the breeding season. *Ecography* 16: 319–328.
- 460 Nummi, P. & Pöysä, H. 1995a. Breeding success of ducks in relation to different  
461 habitat factors. *Ibis* 137: 145–150.
- 462 Nummi, P. & Pöysä, H. 1995b. Habitat use by different-aged duck broods and  
463 juvenile ducks. *Wildl. Biol.* 1: 181–187.
- 464 Nummi, P. & Pöysä, H. 1997. Population and community level responses in *Anas*-  
465 species to patch disturbance caused by an ecosystem engineer, the beaver.  
466 *Ecography* 20: 580–584.
- 467 Nummi, P., Väänänen, V-M., Rask, M., Nyberg, K. & Taskinen, K. 2012. Competitive  
468 effects of fish in structurally simple habitats: perch, invertebrates, and goldeneye in  
469 small boreal lakes. *Aquatic Sci.* 74: 343–350.
- 470 Nyberg, K., Vuorenmaa, J., Tammi, J., Nummi, P., Väänänen, V-M., Mannio, J. & Rask,  
471 M. 2010. Re-establishment of perch in three lakes recovering from acidification:  
472 rapid growth associated with abundant food resources. *Boreal Environ. Res* 15: 480–  
473 490.
- 474 Parker, H., Nummi, P., Hartman, G. & Rosell, F. 2012. Invasive North American  
475 beaver *Castor canadensis* in Eurasia: a review of potential consequences and a  
476 strategy for eradication. *Wild. Biol.* 18: 354–365.
- 477 Paton, P.W.C. 2005. A review of vertebrate community composition in seasonal  
478 forest pools of the northeastern United States. *Wetlands Ecol. Manage.* 13: 235–  
479 246.
- 480 Pirkola M.K. & Högmander J. 1974. Sorsapoikueiden iänmääritys. (English summary:  
481 The age determination of duck broods in the field). *Suomen Riista* 25: 50–55.
- 482 Remillard, M.M., Gruendling, G.K. & Bogucki, D.J. 1987. Disturbance by beaver  
483 (*Castor canadensis* Kuhl) and increased landscape heterogeneity. In: Turner, M. G.  
484 (ed.) *Landscape Heterogeneity and Disturbance*, *Ecological Studies* 64: 103–122.

- 485 Sæther, B-E. & Engen, S. 2002. Pattern of variation in avian population growth rates.  
486 Phil. Trans. Royal Soc. London B 357: 1185–1195.
- 487 Schneider, D.C. 2001. The rise of the concept of scale in ecology. *BioScience* 51:  
488 545–553.
- 489 Sherry, T.W. & Holmes, R.T. 1988. Habitat selection by breeding American Redstarts  
490 in response to a dominant competitor, the Least Flycatcher. *The Auk* 105: 350–364.
- 491 Stringer, A.P & Gaywood, M.J. 2016. The impacts of beavers *Castor* spp. On  
492 biodiversity and the ecological basis for their reintroduction to Scotland, UK.  
493 *Mammal Review* 46: 270–283.
- 494 Suhonen, S., Nummi, P. & Pöysä, H. 2011. Long term stability of habitats and use by  
495 ducks in boreal lakes. *Boreal Environ. Res.* 16 (suppl. B): 71–80.
- 496 Talent, L.G., Krapu, G.L. & Jarvis, R.L. 1982. Habitat use by mallard broods in South  
497 Central North Dakota. *J. Wildl. Manage.* 46: 629–635.
- 498 Taskinen, K. 1997. Vesilintuyhteisöjen rakenne ja poikastuotto Nuuksion  
499 metsäjärvillä – pH:n, ravinnon ja ahventen määrän vaikutus sinisorsan ja telkän  
500 esiintymiseen alueella. MSc Thesis. University of Helsinki.
- 501 Thompson, S., Vehkaoja, M. & Nummi, P. 2016. Beaver-created deadwood dynamics  
502 in the boreal forest. *Forest Ecol. Manage.* 360: 1–8.
- 503 Törnblom, J., Angelstam, P., Hartman, G., Henrikson, L. & Sjöberg, G. 2011. Toward a  
504 research agenda for water policy implementation: Knowledge about beaver (*Castor*  
505 fiber) as a tool for water management with a catchment perspective. *Baltic For.* 17:  
506 154–161.
- 507 Väänänen, V-M., Nummi, P., Pöysä, H., Rask, M. & Nyberg, K. 2012. Fish-duck  
508 interactions in boreal lakes in Finland as reflected by abundance correlations.  
509 *Hydrobiology* 697: 85–93.
- 510 Wiens, J. A. 1989a. *The Ecology of Bird Communities*. Vol. 2. Processes and  
511 variations. Cambridge University Press, Cambridge.
- 512 Wiens, J. A. 1989b. Spatial scaling in ecology. *Functional Ecol.* 3: 385–397.
- 513 Wright, J.P., Jones, C.G. & Flecker, A.S. 2002. An ecosystem engineer, the beaver,  
514 increases species richness at the landscape scale. *Oecologia* 132: 96–101.

516 Fig. 1. Changes in Teal pair and brood densities as well as broods/pair per census in  
517 Beaver flowages (black) and control ponds (grey). Shown are medians and  
518 interquartile ranges. For pairs,  $n = 19$ ; for broods,  $n = 18$ ; for broods/pair,  $n = 12$ .

519 Fig. 2. Teal pair (a) and brood densities (b) as well as broods/pair (c) at Beaver (Evo)  
520 and non-Beaver (Nuuksio) landscapes during 1994–2014.

521

522

523 Table 1. Teal pair density, brood density and broods produced per pair at the Beaver lakes and  
 524 control, non-Beaver lakes of Evo landscape. Test results indicate the independent Mann-Whitney  
 525 U-tests performed between the duck parameters of the Beaver lakes non-Beaver lakes. SD =  
 526 standard deviation; Z = test value; significant results are bolded ( $P < 0.05$ ). IQR = interquartile  
 527 range.

	Beaver lakes ( $n=24$ )	Non-Beaver lakes ( $n=18$ )	Test results
Pair density			
Median	0.46	0.29	$Z = -1.707; P = 0.088$
IQR	1.31	0.33	
Brood density			
Median	0.06	0.00	$Z = -2.188; P = \mathbf{0.029}$
IQR	0.46	0.04	
Broods/pair			
Median	0.03	0.01	$Z = -2.322; P = \mathbf{0.020}$
IQR	0.15	0.02	

528

529